


Complex Geometry I: Complex Differential Geometry

Bowen Liu¹



Don't just read it; fight it!

--- Paul R. Halmos

¹liubw22@mails.tsinghua.edu.cn

CONTENTS

Part 1. Basic Complex Geometry	4
0. Review of complex analysis	4
0.1. One variable case	4
0.2. Several variables case	5
1. Local theory	8
1.1. Algebraic germ	8
1.2. Analytic germ	10
1.3. Hilbert's Nullstellensatz	12
1.4. Dimension	12
1.5. Meromorphic functions and relatively prime	13
2. Complex manifold	14
2.1. Basic definitions and properties	14
2.2. Analytic subvariety	15
2.3. Examples	15
2.4. Vector bundle	18
2.5. Euler sequence and adjunction formula	23
3. Divisor and line bundle	25
3.1. Divisor	25
3.2. Relations between divisor and line bundle	27
3.3. Ample line bundle	28
4. Tangent and cotangent bundle	29
4.1. Complex and holomorphic tangent bundle	29
4.2. Bidegree forms	30
4.3. Dolbeault operators	31
Part 2. Complex Differential Geometry	34
5. Connections and its curvature	34
5.1. Connections on complex vector bundle	34
5.2. Chern connection	35
5.3. First Chern class	40
5.4. Lefschetz $(1, 1)$ -theorem	41
6. Hermitian geometry	45
6.1. Hermitian manifold and Riemannian manifold	45
6.2. Curvatures of Hermitian manifold	49
6.3. Useful formulas of Hermitian geometry	50
6.4. Gauduchon metric	53
6.5. Second fundamental form	54
7. Kähler geometry	57
7.1. Kähler manifold	57
7.2. Levi-Civita connection encounters Chern connection	58
7.3. Curvatures of Kähler metric	59
7.4. Fubini-Study metric	64

Part 3. Hodge theory	66
8. Hodge theorem	66
8.1. Hodge star and adjoint operators	66
8.2. Hodge theorem	69
9. Hodge decomposition	71
9.1. Kähler identities	71
9.2. Hodge decomposition	74
9.3. Bott-Chern cohomology	75
10. Serre duality	77
10.1. Operators on bundle valued forms	77
10.2. Serre duality	78
11. Lefschetz decomposition	80
11.1. Lefschetz decomposition	80
11.2. Hodge index	82
Part 4. Positivity and vanishing theorems	88
12. Positivity	88
12.1. Positivity of line bundle	88
12.2. Positivity of vector bundles	89
13. Vanishing theorems	90
13.1. Kodaira vanishing theorem	90
13.2. Nakano vanishing	92
13.3. Griffiths vanishing	94
14. Kodaira embedding and Chow theorem	95
14.1. Blow-up	95
14.2. Kodaira embedding	96
14.3. Chow's theorem	99
15. More Bochner techniques	102
15.1. Obstruction to holomorphic vector fields	102
15.2. Obstruction to holomorphic 1-form	103
15.3. Rigidity of complex projective space	103
16. Schwarz lemmas	105
16.1. Schwarz lemmas for holomorphic bisectional curvature	105
16.2. Schwarz lemmas for holomorphic sectional curvature	108
Part 5. Topics	111
17. Calabi-Yau theorem	111
17.1. Introduction	111
17.2. The Monge-Ampère equation and priori estimates	111
17.3. Proof of Calabi-Yau theorem	116
17.4. Aubin-Yau theorem	116
18. Deformations of complex structure	119
18.1. The Maurer-Cartan equation	119
19. Hermitian-Yang-Mills metric	122
19.1. Stable bundle	122

Part 6. Appendix	126
20. Sheaf and Cohomology	127
20.1. Sheaves	127
20.2. Derived functor formulation of Sheaf Cohomology	133
20.3. Acyclic resolution	135
20.4. Examples about acyclic sheaf	136
20.5. Proof of de Rham theorem using sheaf cohomology	142
20.6. Hypercohomology	142
21. Hodge theorem	144
References	145

Part 1. Basic Complex Geometry

In this part, we mainly follows [Huy05] and [Dem12].

0. REVIEW OF COMPLEX ANALYSIS

0.1. One variable case. We first give a quick review about basic results in holomorphic functions of one variable. Fix an open subset $U \subseteq \mathbb{C}$. There are too many ways to define a holomorphic function, and all of them are equivalent.

Definition 0.1.1 (holomorphic). A function $f: U \rightarrow \mathbb{C}$ is called holomorphic at $z_0 \in U$, if there exists an open ball $B_\varepsilon(z_0) \subseteq U$ with $\varepsilon > 0$ such that $f|_{B_\varepsilon(z_0)}$ can be written as convergent power series, that is

$$f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n, \quad z \in B_\varepsilon(z_0)$$

f is holomorphic on U , if f is holomorphic at any point of U .

Remark 0.1.1 (Cauchy-Riemann equation). The second definition is given by Cauchy-Riemann equation. To be explicit, for a function $f: U \rightarrow \mathbb{C}$, we can regard it as a function defined on \mathbb{R}^2 , and write it as $f(x, y) = u(x, y) + \sqrt{-1}v(x, y)$, where u, v are real-valued functions, then f is holomorphic if and only if u, v are continuously differentiable and satisfy the following Cauchy-Riemann equations:

$$\begin{aligned} \frac{\partial u}{\partial x} &= \frac{\partial v}{\partial y} \\ \frac{\partial u}{\partial y} &= -\frac{\partial v}{\partial x} \end{aligned}$$

If we introduce the following two operators

$$\begin{aligned} \frac{\partial}{\partial z} &:= \frac{1}{2} \left(\frac{\partial}{\partial x} - \sqrt{-1} \frac{\partial}{\partial y} \right) \\ \frac{\partial}{\partial \bar{z}} &:= \frac{1}{2} \left(\frac{\partial}{\partial x} + \sqrt{-1} \frac{\partial}{\partial y} \right) \end{aligned}$$

Then Cauchy-Riemann equation is equivalent to $\frac{\partial f}{\partial \bar{z}} = 0$. Indeed,

$$\begin{aligned} \frac{\partial f}{\partial \bar{z}} &= \frac{1}{2} \left(\frac{\partial f}{\partial x} + \sqrt{-1} \frac{\partial f}{\partial y} \right) \\ &= \frac{1}{2} \left(\frac{\partial u}{\partial x} + \sqrt{-1} \frac{\partial v}{\partial x} + \sqrt{-1} \frac{\partial u}{\partial y} - \frac{\partial v}{\partial y} \right) \\ &= 0 \end{aligned}$$

Remark 0.1.2 (Cauchy integral formula). The third definition is given by Cauchy integral formula. To be explicit, a function $f: U \rightarrow \mathbb{C}$ is holomorphic

if and only if f is continuously differentiable and for any $B_\varepsilon(z_0) \subseteq U$, the following formula holds

$$f(z_0) = \frac{1}{2\pi\sqrt{-1}} \int_{\partial B_\varepsilon(z_0)} \frac{f(z)}{z - z_0} dz$$

Here are some standard facts in complex analysis, which can be found in any textbook.

Theorem 0.1.1 (maximum principle). Let $U \subseteq \mathbb{C}$ be open and connected. If $f: U \rightarrow \mathbb{C}$ is holomorphic and non-constant, then $|f|$ has no local maximum in U .

Theorem 0.1.2 (identity theorem). If $f, g: U \rightarrow \mathbb{C}$ are two holomorphic functions a connected open subset $U \subseteq \mathbb{C}$ such that $f(z) = g(z)$ for all z in a non-empty subset V of U , then $f = g$.

Theorem 0.1.3 (Riemann extension theorem). Let $f: B_\varepsilon(z_0) - \{z_0\} \rightarrow \mathbb{C}$ be a bounded holomorphic function, then f can be extended to a holomorphic function $f: B_\varepsilon(z_0) \rightarrow \mathbb{C}$.

Theorem 0.1.4 (Riemann mapping theorem). Let $U \subseteq \mathbb{C}$ be a simply-connected proper open subset. Then U is biholomorphic to the unit ball.

Theorem 0.1.5 (Liouville). Every bounded holomorphic function $f: \mathbb{C} \rightarrow \mathbb{C}$ is constant.

Remark 0.1.3. Liouville theorem implies that \mathbb{C} is not biholomorphic to the unit ball. It's a striking difference to the real case since we know unit ball is homeomorphic to \mathbb{R} .

0.2. Several variables case. Now let U be an open subset of \mathbb{C}^n . For any $w \in U$, a polydisc $B_\varepsilon(w) = \{z : |z^i - w_i| < \varepsilon_i\}$, where $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n)$.

Definition 0.2.1 (holomorphic). A function $f: U \rightarrow \mathbb{C}$ is called holomorphic at point $w \in U$, if there exists a polydisc $B_\varepsilon(w) \subseteq U$ such that the restriction of $f|_{B_\varepsilon(w)}$ is given by power series

$$\sum_{i_1, \dots, i_n=0}^{\infty} a_{i_1 \dots i_n} (z_1 - w_1)^{i_1} \dots (z_n - w_n)^{i_n}$$

Remark 0.2.1 (equivalent definitions of holomorphic function).

(1) A function $f: U \rightarrow \mathbb{C}$ is holomorphic, if it satisfies Cauchy-Riemann equations for all coordinates $z^i = x^i + \sqrt{-1}y^i$, that is

$$\frac{\partial f}{\partial \bar{z}^i} = 0, \quad i = 1, 2, \dots, n$$

where $\frac{\partial}{\partial z^i} := \frac{1}{2}(\frac{\partial}{\partial x^i} + \sqrt{-1}\frac{\partial}{\partial y^i})$

- (2) A function $f: U \rightarrow \mathbb{C}$ is holomorphic if and only if f is continuously differentiable and for any $z_0 \in U$, the following formula holds

$$f(z) = \frac{1}{(2\pi\sqrt{-1})^n} \int_{\partial B_\varepsilon(w)} \frac{f(w)}{(z_1 - w_1) \dots (z_n - w_n)} dw_1 \dots dw_n$$

Remark 0.2.2. Other results such as maximum theorem, identity theorem and Liouville theorem generalize easily to the higher dimension. A version of Riemann extension still holds true. However, Riemann mapping theorem fails.

Exercise 0.2.1. Show that polydisc $B_{(1,1)}(0) \subseteq \mathbb{C}^2$ is not biholomorphic to the unit disk $D = \{z \in \mathbb{C}^2 : \|z\| < 1\}$.

Lemma 0.2.1 (local $\partial\bar{\partial}$ -lemma). Let ω be a real $(1,1)$ -form defined on \mathbb{C}^n . Then ω is d-closed if and only if for any point $z \in \mathbb{C}^n$, there exists an open neighborhood U of z and a smooth function $\varphi: U \rightarrow \mathbb{R}$ such that

$$\omega = \sqrt{-1} \partial\bar{\partial}\varphi.$$

Lemma 0.2.2. Let $\varphi: \mathbb{C}^n \rightarrow \mathbb{R}$ be a smooth function such that $\partial\bar{\partial}\varphi = 0$. Then for any point $z \in \mathbb{C}^n$, there exists an open neighborhood U of z and a holomorphic functions $f: U \rightarrow \mathbb{C}$ such that $\varphi = \text{Re}(f)$ over U .

Theorem 0.2.1 (Hartogs' theorem). Suppose $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n)$ and $\varepsilon' = (\varepsilon'_1, \dots, \varepsilon'_n)$ are given such that for all i one has $\varepsilon'_i < \varepsilon_i$. If $n > 1$, then any holomorphic map $f: B_\varepsilon(0) \setminus \overline{B_{\varepsilon'}(0)} \rightarrow \mathbb{C}$ can be uniquely extended to a holomorphic map $f: B_\varepsilon(0) \rightarrow \mathbb{C}$.

Remark 0.2.3. This is only valid in dimension at least two.

Definition 0.2.2 (holomorphic). A function $f: U \rightarrow \mathbb{C}^n$ is called holomorphic if all coordinate functions f_1, \dots, f_n are holomorphic functions $U \rightarrow \mathbb{C}$.

Definition 0.2.3 (biholomorphic). A holomorphic map $f: U \rightarrow V$ between two open subsets $U, V \subseteq \mathbb{C}^n$ is biholomorphic if f is bijective and its inverse f^{-1} is also holomorphic.

Definition 0.2.4 (complex Jacobian). Let $f: U \rightarrow \mathbb{C}^n$ be a holomorphic map, the complex Jacobian of f at point $z \in U$ is the matrix

$$J_{\mathbb{C}}(f)(z) := \left(\frac{\partial f_i}{\partial z^j}(z) \right)_{\substack{1 \leq i \leq n \\ 1 \leq j \leq m}}$$

where $f_i = z^i \circ f$.

Remark 0.2.4. For each $z \in U$, the smooth map $f: U \subseteq \mathbb{C}^m = \mathbb{R}^{2m} \rightarrow \mathbb{C}^n = \mathbb{R}^{2n}$ induces a \mathbb{R} -linear map, which is denoted by $J_{\mathbb{R}}(f)(z): T_z \mathbb{R}^{2m} \rightarrow T_{f(z)} \mathbb{R}^{2n}$. Suppose $\{\frac{\partial}{\partial x^i}, \frac{\partial}{\partial y^i}\}_{1 \leq i \leq m}$ and $\{\frac{\partial}{\partial r^j}, \frac{\partial}{\partial s^j}\}_{1 \leq j \leq n}$ are local frames of $T_z \mathbb{R}^{2m}$ and $T_{f(z)} \mathbb{R}^{2n}$ respectively, then with respect to these basis one has

$$J_{\mathbb{R}}(f)(z) = \begin{pmatrix} \left(\frac{\partial u_i}{\partial x^j} \right) & \left(\frac{\partial u_i}{\partial y^j} \right) \\ \left(\frac{\partial v_i}{\partial x^j} \right) & \left(\frac{\partial v_i}{\partial y^j} \right) \end{pmatrix}$$

where $u_i = r^i \circ f$ and $v_i = s^i \circ f$. If we consider its \mathbb{C} -linear extension, with respect to basis $\{\frac{\partial}{\partial z^i}, \frac{\partial}{\partial \bar{z}^i}\}_{1 \leq i \leq m}$ and $\{\frac{\partial}{\partial w^j}, \frac{\partial}{\partial \bar{w}^j}\}_{1 \leq j \leq n}$, it can be written as

$$J_{\mathbb{R}}(f)(z) = \begin{pmatrix} \left(\frac{\partial f_i}{\partial z^j}\right) & \left(\frac{\partial f_i}{\partial \bar{z}^j}\right) \\ \left(\frac{\partial \bar{f}_i}{\partial z^j}\right) & \left(\frac{\partial \bar{f}_i}{\partial \bar{z}^j}\right) \end{pmatrix}$$

In particular, if f is holomorphic, then $\det J_{\mathbb{R}}(f) = \det J_{\mathbb{C}}(f) \det \overline{J_{\mathbb{C}}(f)} = |\det J_{\mathbb{C}}(f)|^2 \geq 0$.

Definition 0.2.5 (regular value). Let $U \subseteq \mathbb{C}^m$ be an open subset and let $f: U \rightarrow \mathbb{C}^n$ be a holomorphic map, $z \in U$ is called regular point, if $J_{\mathbb{C}}(f)(z)$ is surjective. If every point $z \in f^{-1}(w)$ is regular point, then w is called a regular value.

Remark 0.2.5. In particular, if $f^{-1}(w) = \emptyset$, then w is also called a regular value.

Theorem 0.2.2 (inverse function theorem). Let $f: U \rightarrow V$ be a holomorphic map between two open subsets $U, V \subseteq \mathbb{C}^n$. If $z \in U$ is a regular point, then there exist open subsets $z \in U' \subseteq U$ and $f(z) \in V' \subseteq V$ such that f induces a biholomorphic map $f: U' \rightarrow V'$.

Theorem 0.2.3 (implicit function theorem). Let $U \subseteq \mathbb{C}^m$ be an open subset and let $f: U \rightarrow \mathbb{C}^n$ be a holomorphic map, where $m \geq n$. Suppose $z_0 \in U$ is a point such that

$$\det(J_{\mathbb{C}}(f)(z_0)) \neq 0$$

Then there exist open subsets $U_1 \subseteq \mathbb{C}^{m-n}, U_2 \subseteq \mathbb{C}^n$ and a holomorphic map $g: U_1 \rightarrow U_2$ such that $U_1 \times U_2 \rightarrow U$ and $f(z) = f(z_0)$ if and only if $g(z_{n+1}, \dots, z_m) = (z_1, \dots, z_n)$.

Corollary 0.2.1. Let $U \subseteq \mathbb{C}^m$ be an open subset and $f: U \rightarrow \mathbb{C}^n$ be a holomorphic map. Suppose that $z_0 \in U$ such that $J_{\mathbb{C}}(f)(z_0)$ has maximal rank. Then

- (1) If $m \geq n$, then there exists a biholomorphic map $h: V \rightarrow U'$, where U' is an open subset of U containing z_0 , and V is an open subset of \mathbb{C}^n containing $f(z_0)$, such that $f(h(z_1, \dots, z_n)) = (z_1, \dots, z_n)$.
- (2) If $m \leq n$, then there exists a biholomorphic map $g: V \rightarrow V'$, where V, V' are open subsets of \mathbb{C}^n containing $f(z_0)$, such that $g(f(z)) = (z_1, \dots, z_m, 0, \dots, 0)$.

1. LOCAL THEORY

1.1. Algebraic germ.

1.1.1. *Weierstrass' theorems.* Let $f: B_\varepsilon(0) \rightarrow \mathbb{C}$ be a holomorphic function defined on polydisc $B_\varepsilon(0)$. For any $w = (z_2, \dots, z_n)$ we denote $f_w(z_1)$ the function $f(z_1, \dots, z_n)$. Now we're going to show that all zeros of f are caused by a factor of f which has the form of a Weierstrass polynomial.

Definition 1.1.1 (Weierstrass polynomial). A Weierstrass polynomial is a polynomial in z_1 of the form

$$z_1^d + \alpha_1(w)z_1^{d-1} + \dots + \alpha_d(w)$$

where coefficients $\alpha_i(w)$ are holomorphic functions on some small disc in \mathbb{C}^{n-1} vanishing at the origin.

Remark 1.1.1. Recall the one variable case, any holomorphic function $f(z)$ with a zero of order d at the origin can be written as $z^d h(z)$, where $h(0) \neq 0$. In fact, z^d is a Weierstrass polynomial since in this case, α_i are constants which vanish at origin, that's exactly zero.

Theorem 1.1.1 (Weierstrass preparation theorem). Let $f: B_\varepsilon(0) \rightarrow \mathbb{C}$ be a holomorphic function on the polydisc $B_\varepsilon(0)$. Assume $f(0) = 0$ and $f_0(z_1) \neq 0$. Then there exists a unique Weierstrass polynomial $g_w(z_1)$ and a holomorphic function h on some smaller polydisc $B_{\varepsilon'}(0) \subseteq B_\varepsilon(0)$ such that $f = gh$ and $h(0) \neq 0$.

Proof. See Proposition 1.1.6 in Page8 of [Huy05]. □

Theorem 1.1.2 (Weierstrass division theorem). Let $f \in \mathcal{O}_{\mathbb{C}^n, 0}$ and let $g \in \mathcal{O}_{\mathbb{C}^{n-1}, 0}[z_1]$ be a Weierstrass polynomial of degree d . Then there exist $r \in \mathcal{O}_{\mathbb{C}^{n-1}, 0}[z_1]$ of degree $< d$ and $h \in \mathcal{O}_{\mathbb{C}^n, 0}$ such that $f = gh + r$. The functions h and r are uniquely determined.

Proof. See Proposition 1.1.17 in Page15 of [Huy05]. □

1.1.2. *Stalk of sheaf of holomorphic functions.* Let's use $\mathcal{O}_{\mathbb{C}^n}$ to denote the sheaf² of holomorphic functions on \mathbb{C}^n , and use $\mathcal{O}_{\mathbb{C}^n, 0}$ to denote its stalk at origin. The elements in $\mathcal{O}_{\mathbb{C}^n, 0}$ are called germs. It's clear $\mathcal{O}_{\mathbb{C}^n, 0}$ is a local ring with maximal ideal \mathfrak{m} consisting of all functions that vanish at origin, which implies units in $\mathcal{O}_{\mathbb{C}^n, 0}$ are functions that don't vanish at origin.

By using Weierstrass preparation theorem, one can derive more about algebraic properties of $\mathcal{O}_{\mathbb{C}^n, 0}$. For example, Weierstrass preparation theorem can be rephrased by saying that after an appropriate coordinate choice any function $f \in \mathcal{O}_{\mathbb{C}^n, 0}$ can be uniquely written as $f = gh$, where $h \in \mathcal{O}_{\mathbb{C}^n, 0}$ is a unit and $g \in \mathcal{O}_{\mathbb{C}^{n-1}, 0}[z_1]$ is a Weierstrass polynomial. Furthermore, it also shows the following important property.

²Sheaf and its cohomology are important tools we will use once and again, if you're not familiar with it, see Appendix 20.

Theorem 1.1.3. The local ring $\mathcal{O}_{\mathbb{C}^n,0}$ is a UFD.

Proof. We prove the assumption by induction on n . For $n = 0$, the ring $\mathcal{O}_{\mathbb{C}^n,0} = \mathbb{C}$ is a field, and thus a UFD. Suppose that $\mathcal{O}_{\mathbb{C}^{n-1},0}$ is a UFD, for $f \in \mathcal{O}_{\mathbb{C}^n,0}$ we choose coordinates such that Weierstrass preparation theorem is applied, that is $f = gh$, where $g \in \mathcal{O}_{\mathbb{C}^{n-1},0}[z_1]$ is a Weierstrass polynomial and h is a unit in $\mathcal{O}_{\mathbb{C}^n,0}$. By induction we have $\mathcal{O}_{\mathbb{C}^{n-1},0}[z_1]$ is UFD, then g can be written as a product of irreducible elements of $\mathcal{O}_{\mathbb{C}^{n-1},0}[z_1]$. All that is left to show is that any irreducible element in $\mathcal{O}_{\mathbb{C}^{n-1},0}[z_1]$ is also irreducible as an element in $\mathcal{O}_{\mathbb{C}^n,0}$.

Assume $g \in \mathcal{O}_{\mathbb{C}^{n-1},0}[z_1]$ is a Weierstrass polynomial which is written as the product of non-units $g_i \in \mathcal{O}_{\mathbb{C}^{n-1},0}[z_1]$. There are two cases:

- (1) $g_i \in \mathcal{O}_{\mathbb{C}^{n-1},0}$. By induction hypothesis, g_i can be written as the product of irreducible elements of $\mathcal{O}_{\mathbb{C}^{n-1},0}$, which are also irreducible in $\mathcal{O}_{\mathbb{C}^n,0}$.
- (2) $g_i \notin \mathcal{O}_{\mathbb{C}^{n-1},0}$. In this case, g_i satisfies the hypothesis of Weierstrass preparation theorem since g_i is a Weierstrass polynomial, then g_i is non-trivial on the z_1 -line. So we can write $g_i = \tilde{g}_i h_i$, where \tilde{g}_i is also Weierstrass polynomial.

Note that degree of g as a polynomial in z_1 is finite, then repeating above process leads to a decomposition, with factors are either irreducible Weierstrass polynomials or elements in $\mathcal{O}_{\mathbb{C}^{n-1},0}$.

Now it suffices to show any irreducible Weierstrass polynomial g is actually irreducible as an element of $\mathcal{O}_{\mathbb{C}^n,0}$. Suppose $g = f_1 f_2$, where $f_1, f_2 \in \mathcal{O}_{\mathbb{C}^n,0}$ are non-units. We apply Weierstrass preparation theorem to obtain $f_i = g_i h_i, i = 1, 2$, and thus $g = (g_1 g_2)(f_1 f_2)$. By uniqueness one has $g = g_1 g_2$, which contradicts to the irreducibility of g as an element of $\mathcal{O}_{\mathbb{C}^{n-1},0}[z_1]$. \square

Another important fact is that $\mathcal{O}_{\mathbb{C}^n,0}$ is noetherian, which follows from Weierstrass division theorem.

Theorem 1.1.4. The local UFD $\mathcal{O}_{\mathbb{C}^n,0}$ is noetherian.

Proof. We prove the assumption by induction on n . For $n = 0$, it's clear since any field is noetherian. Suppose that $\mathcal{O}_{\mathbb{C}^{n-1},0}$ is noetherian, then Hilbert's basis theorem implies $\mathcal{O}_{\mathbb{C}^{n-1},0}[z_1]$ is also noetherian. Let $I \subseteq \mathcal{O}_{\mathbb{C}^n,0}$ be a non-trivial idea and choose $0 \neq f \in I$. Changing coordinates if necessary, we may assume Weierstrass preparation theorem is applied, that is $f = gh$, where $g \in \mathcal{O}_{\mathbb{C}^{n-1},0}[z_1]$ is a Weierstrass polynomial and h is a unit in $\mathcal{O}_{\mathbb{C}^n,0}$, hence $g \in I$. Furthermore, we assume $I \cap \mathcal{O}_{\mathbb{C}^{n-1},0}[z_1]$ is generated by g_1, \dots, g_k .

For any other $\tilde{f} \in I$, the Weierstrass division theorem implies $\tilde{f} = g\tilde{h} + r$ for some $r \in \mathcal{O}_{\mathbb{C}^{n-1},0}[z_1]$. Since $\tilde{f}, g\tilde{h} \in I$, we have $r \in I$ and therefore $r \in I \cap \mathcal{O}_{\mathbb{C}^{n-1},0}[z_1]$. Thus $\tilde{f} = g\tilde{h} + \sum_{i=1}^k a_i g_i$. This shows I is finitely generated by elements g, g_1, \dots, g_k . \square

Corollary 1.1.1. Let $g \in \mathcal{O}_{\mathbb{C}^n,0}$ be an irreducible element. If $f \in \mathcal{O}_{\mathbb{C}^n,0}$ vanishes on $Z(g) = \{z \mid g(z) = 0\}$, then g divides f .

Proof. By Weierstrass preparation theorem we may assume $g \in \mathcal{O}_{\mathbb{C}^{n-1},0}[z_1]$ is a Weierstrass polynomial with degree d . By the Weierstrass division theorem one finds $h \in \mathcal{O}_{\mathbb{C}^n,0}$ and $r \in \mathcal{O}_{\mathbb{C}^{n-1},0}[z_1]$ of degree $< d$ such that $f = gh + r$. For $w \in \mathbb{C}^{n-1}$, by assumption r_w vanishes on the zero set g_w . If all of zeros of g_w have multiplicity one, then $r_w \equiv 0$ since r_w is of degree $< d$. Now it suffices to show the set $w \in \mathbb{C}^{n-1}$ such that g_w has zeros with multiplicity > 1 is quite “small”.

Since g is irreducible and $\frac{\partial g}{\partial z_1}$ is of degree $d-1$, there exist elements $h_1, h_2 \in \mathcal{O}_{\mathbb{C}^{n-1},0}[z_1]$ and $0 \neq \gamma \in \mathcal{O}_{\mathbb{C}^{n-1},0}$ such that $h_1 g + h_2 \frac{\partial g}{\partial z_1} = \gamma$. So if g_w has a zero ξ of multiplicity > 1 , then $\gamma(w) = h_1(\xi, w)g_w(\xi) + h_2(\xi, w)\frac{\partial g_w}{\partial z_1}(\xi) = 0$. This shows such w is contained in the zero set of a non-trivial holomorphic function $\gamma \in \mathcal{O}_{\mathbb{C}^{n-1},0}$. Then the following exercise completes the proof. \square

Exercise 1.1.1. Let $U \subseteq \mathbb{C}^n$ be open and connected. Show that for any non-trivial holomorphic function $f: U \rightarrow \mathbb{C}$ the complement $U \setminus Z(f)$ of the zero set of f is connected and dense in U .

1.2. Analytic germ. For any $f \in \mathcal{O}_{\mathbb{C}^n,0}$, $Z(f)$ is not well-defined in fact since for another $g \in \mathcal{O}_{\mathbb{C}^n,0}$, which represents the same element with f , $Z(f)$ may not equal to $Z(g)$. However, there always exists an open neighborhood $0 \in U \subseteq \mathbb{C}^n$ such that $Z(f) \cap U = Z(g) \cap U$.

Definition 1.2.1 (germ of a set). The germ of a set in the origin $0 \in \mathbb{C}^n$ is given by a subset $X \subseteq \mathbb{C}^n$. Two germs of a set in the origin $X, Y \subseteq \mathbb{C}^n$ are same if there exists an open neighborhood $0 \in U \subseteq \mathbb{C}^n$ such that $X \cap U = Y \cap U$.

Unless otherwise specified, in this section we only consider germ of a set in the origin, and for convenience we just call it a germ.

Example 1.2.1. For $f \in \mathcal{O}_{\mathbb{C}^n,0}$, $Z(f)$ is a germ.

Definition 1.2.2 (analytic germ). A germ $X \subseteq \mathbb{C}^n$ is called analytic if there exist elements $f_1, \dots, f_k \in \mathcal{O}_{\mathbb{C}^n,0}$ such that $X = Z(f_1, \dots, f_k) := \bigcap_{i=1}^k Z(f_i)$.

Example 1.2.2. Let A be a subset of $\mathcal{O}_{\mathbb{C}^n,0}$. If we use (A) to denote the idea generated by A , then (A) is finitely generated since $\mathcal{O}_{\mathbb{C}^n,0}$ is noetherian. Thus $Z((A))$ is an analytic germ.

Definition 1.2.3. Let $X \subseteq \mathbb{C}^n$ be a germ. Then $I(X)$ denotes the set of all elements $f \in \mathcal{O}_{\mathbb{C}^n,0}$ with $X \subseteq Z(f)$.

Remark 1.2.1. It's clear $I(X)$ is an idea of $\mathcal{O}_{\mathbb{C}^n,0}$.

Lemma 1.2.1.

- (1) If $X_1 \subseteq X_2$ are germs, then $I(X_2) \subseteq I(X_1)$.
- (2) If $\mathfrak{a}_1 \subseteq \mathfrak{a}_2$ are two ideas of $\mathcal{O}_{\mathbb{C}^n,0}$, then $Z(\mathfrak{a}_2) \subseteq Z(\mathfrak{a}_1)$.

- (3) For any analytic germ one has $Z(I(X)) = X$.
(4) For any idea \mathfrak{a} of $\mathcal{O}_{\mathbb{C}^n,0}$ one has $\mathfrak{a} \subseteq I(Z(\mathfrak{a}))$

Proof. (1), (2) and (4) are clear. For (3). It's clear $X \subseteq Z(I(X))$. On the other hand since X is analytic germ there exist elements $f_1, \dots, f_k \in \mathcal{O}_{\mathbb{C}^n,0}$ such that $X = Z(f_1, \dots, f_k)$ as germs, thus $f_1, \dots, f_k \in I(X)$, so by (2) we have $Z(I(X)) \subseteq X = Z(f_1, \dots, f_k)$. This completes the proof of (3). \square

Definition 1.2.4 (irreducible germ). An analytic germ is irreducible if the following condition is satisfied: If $X = X_1 \cup X_2$, where X_1, X_2 are analytic germs, then $X = X_1$ or $X = X_2$.

Lemma 1.2.2. An analytic germ X is irreducible if and only if $I(X) \subseteq \mathcal{O}_{\mathbb{C}^n,0}$ is a prime ideal.

Proof. If X is irreducible and $f_1 f_2 \in I(X)$, then $X \subseteq Z(f_1 f_2) = Z(f_1) \cup Z(f_2)$, so we have $X = (X \cap Z(f_1)) \cup (X \cap Z(f_2))$ is a union of analytic germs. Then by irreducibility one has $X = X \cap Z(f_i)$ for some $i = 1$ or $i = 2$, and thus at least one of functions f_1 or f_2 vanishes on X . This shows $I(X)$ is prime.

Conversely, if $I(X)$ is a prime ideal and let $X = X_1 \cup X_2$ with X_1 and X_2 are analytic. If $f_i \in I(X_i), i = 1, 2$, then $f_1 f_2 \in I(X)$ since

$$X = X_1 \cup X_2 \subseteq Z(f_1) \cup Z(f_2) = Z(f_1 f_2)$$

Hence $f_1 \in I(X)$ or $f_2 \in I(X)$. Thus it suffices to shows that if $X \neq X_1$ and $X \neq X_2$, there exist elements $f_1 \in I(X_1) \setminus I(X)$ and $f_2 \in I(X_2) \setminus I(X)$. This follows immediately from (1) of Lemma 1.2.1. \square

Corollary 1.2.1. For $f \in \mathcal{O}_{\mathbb{C}^n,0}$, $Z(f)$ is irreducible if and only if there exists an irreducible $g \in \mathcal{O}_{\mathbb{C}^n,0}$ such that $f = g^k$ for some $k \in \mathbb{Z}_{>0}$.

Proof. If $f = g^k$ with g irreducible, then $Z(f) = Z(g)$ and if $h \in I(Z(g))$, then g divides h by Corollary 1.1.1, this shows $I(Z(g)) = (g)$ and thus $Z(f)$ is irreducible since $I(Z(f))$ also equals to (g) , which is prime. Conversely, if $f = \prod g_i^{n_i}$, then $Z(f) = \bigcup Z(g_i)$, which cannot be irreducible except for the case $f = g^k$ for some irreducible g . \square

Lemma 1.2.3. Every decreasing sequences of germs $\{X_i\}$ is stationary.

Proof. Consider its corresponding sequence $\{I(X_i)\}$, it's an increasing sequence, thus it's stationary since $\mathcal{O}_{\mathbb{C}^n,0}$ is noetherian, this completes the proof since for each i , $Z(I(X_i)) = X_i$. \square

Theorem 1.2.1. Every germ X admits a finite decomposition $X = \bigcup_{i=1}^N X_i$, where X_i is irreducible for each i and $X_i \not\subseteq X_j$ for $i \neq j$. The decomposition is unique apart from the ordering.

Proof. It suffices to show uniqueness since existence follows from above lemma. Assume $X = \bigcup_{l=1}^{N'} X'_l$ is another decomposition, note that $X_i = \bigcup_{l=1}^{N'} X_i \cap X'_l$, we must have $X_i = X_i \cap X'_{l(i)}$ since X_i is irreducible. Likewise

one has $X'_{l(i)} \cap X_j$, so we $i = j$ since $X_i \not\subseteq X_j$ for $i \neq j$, and this shows $X_i = X'_{l(j)}$ \square

1.3. Hilbert's Nullstellensatz.

Theorem 1.3.1. Let $X \subseteq \mathbb{C}^n$ be an irreducible analytic germ defined by a prime ideal $\mathfrak{p} \subseteq \mathcal{O}_{\mathbb{C}^n,0}$. Then one can find a coordinate system

$$(z_1, \dots, z_{n-d}, z_{n-d+1}, \dots, z_n)$$

such that the projection $(z_1, \dots, z_n) \rightarrow (z_{n-d+1}, \dots, z_n)$ induces a surjective map of germs $X \rightarrow \mathbb{C}^d$ and such that the induced ring homomorphism $\mathcal{O}_{\mathbb{C}^d,0} \rightarrow \mathcal{O}_{\mathbb{C}^n,0}/\mathfrak{p}$ is a finite integral ring extension.

Proof. See (4.19) of [Dem12]. \square

Theorem 1.3.2 (Hilbert's Nullstellensatz). If $I \subseteq \mathcal{O}_{\mathbb{C}^n,0}$ is any ideal, then $\sqrt{I} = I(Z(I))$.

Proof. It easy to see $\sqrt{I} \subseteq I(Z(I))$. Conversely, it suffices to show $I(Z(I)) \subseteq \mathfrak{p}$ for all prime ideals containing I since \sqrt{I} is the intersection of all prime ideals \mathfrak{p} containing I . If one has

$$\mathfrak{p} = \sqrt{\mathfrak{p}} = I(Z(\mathfrak{p}))$$

then the results follows from $Z(\mathfrak{p}) \subseteq Z(I)$. Thus we reduce the problem to the case to $I = \mathfrak{p}$ is a prime ideal. For $f \in I(Z(\mathfrak{p}))$, by Theorem 1.3.1 there exists an appropriate coordinate system (z_1, \dots, z_n) such that the induced element $\bar{f} \in \mathcal{O}_{\mathbb{C}^n,0}/\mathfrak{p}$ satisfies an irreducible algebraic equation $\bar{f}^k + a_1 \bar{f}^{k-1} + \dots + a_k = 0$ with $a_i \in \mathcal{O}_{\mathbb{C}^d,0}$. Since f vanishes along $Z(\mathfrak{p})$, the 0-th coefficient a_k does as well. As $Z(\mathfrak{p}) \rightarrow \mathbb{C}^d$ is surjective, this shows $a_k = 0$. Hence above algebraic equation cannot be irreducible unless $k = 1$. Therefore $\bar{f} = 0$ and thus $f \in \mathfrak{p}$. \square

Corollary 1.3.1. There is a one to one correspondence between prime ideals of $\mathcal{O}_{\mathbb{C}^n,0}$ and irreducible analytic germ given by $X \mapsto I(X)$ and $\mathfrak{p} \rightarrow Z(\mathfrak{p})$.

1.4. Dimension.

Definition 1.4.1 (dimension). Let X be an irreducible analytic germ defined by a prime ideal $\mathfrak{p} \subseteq \mathcal{O}_{\mathbb{C}^n,0}$. Then the dimension of X is defined by $n - \text{htp}$, where htp is the height of \mathfrak{p} .

Remark 1.4.1. For arbitrary analytic germ is of dimension d if all its irreducible components are of the same dimension d .

Remark 1.4.2. If $X \subseteq \mathbb{C}^n$ is an irreducible analytic germ of codimensional 1, then the prime ideal \mathfrak{p} defining X is of height 1. A basic result in commutative algebra says any prime ideal of height 1 in a UFD is principle. Therefore, $\mathfrak{p} = (f)$ for some irreducible $f \in \mathcal{O}_{\mathbb{C}^n,0}$.

1.5. Meromorphic functions and relatively prime.

Definition 1.5.1. Let $U \subseteq \mathbb{C}^n$ be an open subset. A meromorphic function f on U is a function on the complement of a nowhere dense subset $S \subseteq U$ with the following property: There exist an open covering $\{U_i\}$ of U and holomorphic functions $g_i, h_i: U \rightarrow \mathbb{C}$ with $h_i|_{U_i \setminus S} \cdot f|_{U_i \setminus S} = g_i|_{U_i \setminus S}$.

Remark 1.5.1. For any $z \in U$, the meromorphic function f in a neighborhood of z is given by g/h , where $g, h \in \mathcal{O}_{\mathbb{C}^n, z}$. If we assume g, h are chosen to be relatively prime, then they're unique up to units.

Proposition 1.5.1. Let $f \in \mathcal{O}_{\mathbb{C}^n, 0}$ be irreducible, then for sufficiently small ε and $z \in B_\varepsilon(0)$ the induced element $f \in \mathcal{O}_{\mathbb{C}^n, z}$ is irreducible.

Proof. Suppose $f \in \mathcal{O}_{\mathbb{C}^n, z}$ is reducible, that is $f = f_1 f_2$ where $f_i \in \mathcal{O}_{\mathbb{C}^n, z}$ non-units, i.e. $f_1(z) = f_2(z) = 0$. Thus $\frac{\partial f}{\partial z_1}(z) = \frac{\partial f_1}{\partial z_1}(z) f_2(z) + f_1(z) \frac{\partial f_2}{\partial z_1}(z) = 0$.

Thus the set of points $z \in B_\varepsilon(0)$ where f as an element of $\mathcal{O}_{\mathbb{C}^n, z}$ is reducible is contained in the analytic set $Z(f, \frac{\partial f}{\partial z_1})$. Now it suffices to show it's a proper subset of $Z(f)$ since f is irreducible, so is $Z(f)$. If not, then $\frac{\partial f}{\partial z_1}$ would vanish on $Z(f)$. Since f is irreducible, we can apply Corollary 1.1.1 to obtain $\frac{\partial f}{\partial z_1}$ divides f , a contradiction. \square

Proposition 1.5.2. If $f, g \in \mathcal{O}_{\mathbb{C}^n, 0}$ are relatively prime, then they're relatively prime in $\mathcal{O}_{\mathbb{C}^n, z}$, for z in a sufficiently small neighborhood of 0.

Proof. Without loss of generality, we may assume $f, g \in \mathcal{O}_{\mathbb{C}^{n-1}, 0}[z_1]$ are Weierstrass polynomials, then f and g are relatively prime if and only if their resultant $R \in \mathcal{O}_{\mathbb{C}^{n-1}}$ has non-zero germ at 0, therefore the germ of R is also non-zero in a sufficiently small neighborhood of 0. \square

2. COMPLEX MANIFOLD

2.1. Basic definitions and properties.

Definition 2.1.1 (holomorphic atlas). A holomorphic atlas on a smooth manifold is an atlas $\{(U_\alpha, \varphi_\alpha)\}$ of the form $\varphi_\alpha : U_\alpha \cong \varphi_\alpha(U_\alpha) \subseteq \mathbb{C}^n$ such that transition functions $\varphi_{\alpha\beta} := \varphi_\alpha \circ \varphi_\beta^{-1} : \varphi_\beta(U_{\alpha\beta}) \rightarrow \varphi_\alpha(U_{\alpha\beta})$ are holomorphic functions. Furthermore,

- (1) the pair $(U_\alpha, \varphi_\alpha)$ is called a holomorphic chart.
- (2) two holomorphic atlases are called equivalent, if the union of them is still a holomorphic atlas.

Definition 2.1.2 (complex manifold). A complex n -manifold X is a smooth $2n$ -manifold admitting an equivalence class of holomorphic atlases.

Remark 2.1.1. A complex manifold is called connected, compact, simply-connected and so on, if its underlying real manifold has this property.

Definition 2.1.3 (submanifold). Let X be a complex n -manifold and $Y \subseteq X$ be a smooth manifold of (real) dimension $2k$. Then Y is a complex submanifold if there exists a holomorphic atlas $\{(U_i, \varphi_i)\}$ of X such that $\varphi_i : U_i \cap Y \cong \varphi_i(U_i) \cap \mathbb{C}^k$.

Definition 2.1.4 (holomorphic map). Let X, Y be complex manifolds. A continuous map $f : X \rightarrow Y$ is a holomorphic map if for any holomorphic charts (U, φ) and (U', φ') of X and Y respectively, the map $\varphi' \circ f \circ \varphi^{-1} : \varphi(f^{-1}(U') \cap U) \rightarrow \varphi'(U')$ is holomorphic.

Definition 2.1.5 (biholomorphic). Let X, Y be two complex manifolds. X, Y are called biholomorphic, if there exists a holomorphic homeomorphism $f : X \rightarrow Y$.

Definition 2.1.6 (holomorphic function). A holomorphic function on complex manifold X is a holomorphic map $f : X \rightarrow \mathbb{C}$.

Notation 2.1.1. We always use \mathcal{O}_X to denote the sheaf of holomorphic functions on complex manifold X , and use $\Gamma(U, \mathcal{O}_X)$ to denote sections over open subset $U \subseteq X$.

Proposition 2.1.1. Let X be a compact connected complex manifold. Then $\Gamma(X, \mathcal{O}_X) = \mathbb{C}$.

Proof. It's clear from maximum principle. □

Definition 2.1.7 (meromorphic function). A meromorphic function on a complex manifold X is a map $f : X \rightarrow \coprod_{x \in X} \mathbb{C} \setminus \{0\}$ which associates to any $x \in X$ an element $f_x \in \mathbb{C} \setminus \{0\}$ such that for any $x_0 \in X$ there exists a neighborhood $U \subseteq X$ and two holomorphic functions $g, h : U \rightarrow \mathbb{C}$ with $f_x = g/h$ for all $x \in U$.

Notation 2.1.2. We always use \mathcal{K}_X to denote the sheaf of meromorphic functions on complex manifold X , and use $K(X)$ to denote $\Gamma(X, \mathcal{K}_X)$.

Definition 2.1.8 (algebraic dimension). The algebraic dimension of a compact connected complex manifold X is $a(X) := \text{trdeg}_{\mathbb{C}} K(X)$.

Proposition 2.1.2 (Siegel). Let X be a compact connected complex n -manifold. Then

$$\text{trdeg}_{\mathbb{C}} K(X) \leq n$$

Proof. See Proposition 2.1.9 in Page 54 of [Huy05]. □

2.2. Analytic subvariety.

Definition 2.2.1 (analytic subvariety). Let X be a complex manifold. An analytic subvariety of X is a closed subset $Y \subseteq X$ such that for any $x \in Y$ there exists an open neighborhood $x \in U \subseteq X$ such that $Y \cap U$ is a zero set of finitely many holomorphic functions $f_1, \dots, f_k \in \mathcal{O}(U)$.

Remark 2.2.1. Obviously, any analytic subvariety X defines an analytic germ in any point $z \in X$.

Definition 2.2.2 (irreducible analytic subvariety). An analytic subvariety Y is called irreducible, if it cannot be written as the union $Y = Y_1 \cup Y_2$ of two proper analytic subvarieties $Y_i \subseteq Y, i = 1, 2$.

Given an analytic subvariety Y of a complex manifold X .

Definition 2.2.3 (regular). A point $x \in Y$ is called regular point, if the functions f_1, \dots, f_k can be chosen such that $\varphi(x) \in \varphi(U)$ is a regular point of holomorphic map $f := (f_1 \circ \varphi^{-1}, \dots, f_k \circ \varphi^{-1}) : \varphi(U) \rightarrow \mathbb{C}^k$, where (U, φ) is a local chart of x .

Definition 2.2.4 (singular). A point $x \in Y$ is singular, if it's not regular.

Proposition 2.2.1. The set of regular points $Y_{\text{reg}} = Y \setminus Y_{\text{sing}}$ is a non-empty submanifold of X . Furthermore, if Y is irreducible, then Y_{reg} is connected.

Definition 2.2.5 (dimension). The dimension of an irreducible analytic subvariety Y is defined by $\dim Y = \dim Y_{\text{reg}}$.

2.3. Examples.

Example 2.3.1 (affine space). The n -dimensional complex plane \mathbb{C}^n is a complex manifold.

Example 2.3.2 (complex tori). If V is a complex vector space of dimension n and $\Gamma \subseteq V$ is a free abelian, discrete subgroup of order $2n$, then $X = V/\Gamma$ is a complex manifold, which is called complex tori.

Remark 2.3.1. The underlying manifolds of complex tori with different Γ are not very interesting since they are all diffeomorphic to $(S^1)^{2n}$. However, if you pick two lattices Γ_1, Γ_2 randomly, then \mathbb{C}^n/Γ_1 and \mathbb{C}^n/Γ_2 will not be biholomorphic to each other.

Example 2.3.3 (projective space). The projective space $\mathbb{C}\mathbb{P}^n$ is a complex manifold. Indeed, atlas are given by $U_i = \{[z] \in \mathbb{C}\mathbb{P}^n \mid z^j \neq 0\}, 0 \leq i \leq n$, and φ_i is defined as

$$\begin{aligned} \varphi_i: U_i &\rightarrow \mathbb{C}^n \\ [z] &\mapsto \left(\frac{z_0}{z^i}, \dots, \widehat{\frac{z^i}{z^i}}, \dots, \frac{z_n}{z^i}\right) \end{aligned}$$

The transition functions are calculated as follows: For $i < j$

$$\varphi_i \circ \varphi_j^{-1}: (u_1, \dots, u_n) \mapsto \left(\frac{u_1}{u_i}, \dots, \widehat{\frac{u_i}{u_i}}, \dots, \frac{u_{j-1}}{u_i}, \frac{1}{u_i}, \frac{u_{j+1}}{u_i}, \dots, \frac{u_n}{u_i}\right)$$

It's holomorphic on $U_i \cap U_j$.

Remark 2.3.2. $\mathbb{C}\mathbb{P}^n$ is compact since $\mathbb{C}\mathbb{P}^n$ is diffeomorphic to S^{2n+1}/S^1 , which is called Hopf fibration.

Definition 2.3.1 (projective manifold). A complex manifold X is called projective if X is biholomorphic to a closed complex submanifold of some projective space $\mathbb{C}\mathbb{P}^N$.

Example 2.3.4 (Grassmannian manifold). The Grassmannian manifold

$$Gr(k, n+1) = \{k\text{-dimensional subspace of } \mathbb{C}^{n+1}\}$$

Now we're going to show $Gr(k, n+1)$ is a manifold of dimension $k(n+1-k)$. Any $W \in Gr(k, n+1)$ is generated by the rows of a $k \times (n+1)$ matrix A of rank k . Let us denote the set of these matrices by $M_{k,n+1}$, which is an open subset of the set of all $k \times (n+1)$ matrices. The latter space is a complex manifold which is canonically isomorphic to $\mathbb{C}^{k(n+1)}$. Thus we obtain a natural surjection $\pi: M_{k,n+1} \rightarrow Gr(k, n+1)$, which is the quotient by the natural action of $GL(k, \mathbb{C})$ on $M_{k,n+1}$.

Let's fix an ordering $\{B_1, \dots, B_m\}$ of all $k \times k$ -minors of matrices $A \in M_{k,n+1}$. Define an open covering $Gr(k, n+1) = \bigcup_{i=1}^m U_i$, where U_i is the open subset $\{\pi(A) \mid \det(B_i) \neq 0\}$. Note that U_i is well-defined since if $\pi(A) = \pi(A')$, then A and A' differs an action of $GL(k, \mathbb{C})$, so $\det(B_i) \neq 0$ if and only if $\det(B'_i) \neq 0$. So without lose of generality, we may assume A is of form (B_i, C_i) , where C_i is a $k \times (n+1-k)$ matrix. Then the map $\varphi_i: U_i \rightarrow \mathbb{C}^{k(n+1-k)}$, given by $\pi(A) \rightarrow B_i^{-1}C_i$ is well-defined, and $\{(U_i, \varphi_i)\}$ will give atlas of $Gr(k, n+1)$, sicne all operations are matrix operation, thus they're holomorphic. This shows $Gr(k, n+1)$ is a complex manifold with dimension $k(n+1-k)$.

Remark 2.3.3. If V is a complex vector space of dimension $n+1$, then $Gr(k, V)$ is defined as the set consisting of all k -dimensional subspaces of V , which is biholomorphic to $Gr(k, n+1)$.

Example 2.3.5 (Plücker embedding). Let V be a complex vector space of dimension $n + 1$, then

$$\Phi: Gr(k, V) \hookrightarrow \mathbb{C}\mathbb{P}(\bigwedge^k V)$$

defined by $W \subseteq V$ with basis w_1, \dots, w_k is mapped to $[w_1 \wedge \dots \wedge w_k]$, is called Plücker embedding. It's well-defined, thanks to the following lemma.

Lemma 2.3.1. Let W be a complex vector space of dimension k , and $\mathcal{B}_1 = \{w_1, \dots, w_k\}$ and $\mathcal{B}_2 = \{v_1, \dots, v_k\}$ are two basis for W . Then $v_1 \wedge \dots \wedge v_k = \lambda w_1 \wedge \dots \wedge w_k$ for some $\lambda \in \mathbb{C}^*$.

Proof. If we express $w_j = a_{1j}v_1 + \dots + a_{kj}v_k$, then direct computation shows that

$$\begin{aligned} w_1 \wedge \dots \wedge w_k &= (a_{11}v_1 + \dots + a_{k1}v_k) \wedge \dots \wedge (a_{1k}v_1 + \dots + a_{kk}v_k) \\ &= \sum_{\sigma \in S_k} \text{sign}(\sigma) a_{1\sigma(1)} \dots a_{k\sigma(k)} v_1 \wedge \dots \wedge v_k \\ &= \lambda v_1 \wedge \dots \wedge v_k \end{aligned}$$

Note that λ is exactly the determinant of the change of basis matrix from \mathcal{B}_1 to \mathcal{B}_2 . \square

Remark 2.3.4. It's a little bit complicated to check it's injective.

2.4. Vector bundle.

2.4.1. In viewpoint of transition functions.

Definition 2.4.1 (complex vector bundle). Let X be a smooth manifold. A complex vector bundle E of rank r on X consists of the following data:

- (1) E is a smooth manifold with surjective map $\pi: E \rightarrow X$, such that
 - (1) For all $x \in X$, fibre E_x is a \mathbb{C} -vector space of dimension r .
 - (2) For all $x \in X$, there exists $u \in U \subseteq X$ and there is a homeomorphism $\varphi: \pi^{-1}(U) \rightarrow U \times \mathbb{C}^r$ such that

$$\begin{array}{ccc}
 \pi^{-1}(U) & \xrightarrow{\pi} & U \\
 \searrow \varphi & \curvearrowright & \nearrow p_1 \\
 & & U \times \mathbb{C}^r \\
 & & \xrightarrow{p_2} \mathbb{C}^r
 \end{array}$$

and for all $y \in U$, $E_y \xrightarrow{p_2 \circ \varphi} \mathbb{C}^r$ is a \mathbb{C} -vector space isomorphism. (U, φ) is called a trivialization of E over U .

Remark 2.4.1 (transition functions). Consider two local trivialization $(U_\alpha, \varphi_\alpha), (U_\beta, \varphi_\beta)$. Then $\varphi_\alpha \circ \varphi_\beta^{-1}: (U_\alpha \cap U_\beta) \times \mathbb{C}^r \rightarrow (U_\alpha \cap U_\beta) \times \mathbb{C}^r$ induces

$$g_{\alpha\beta}: U_\alpha \cap U_\beta \rightarrow \text{GL}(r, \mathbb{C})$$

where $g_{\alpha\beta}$ is called transition function. Furthermore, it satisfies

$$\begin{aligned}
 g_{\alpha\beta}g_{\beta\gamma}g_{\gamma\alpha} &= \text{id} && \text{on } U_\alpha \cap U_\beta \cap U_\gamma \\
 g_{\alpha\alpha} &= \text{id} && \text{on } U_\alpha
 \end{aligned}$$

In fact, transition functions contain all information about this vector bundle since a vector bundle is locally trivial, so how are these trivial pieces glued together really matters.

Definition 2.4.2 (complex vector bundle). Let X be a smooth manifold. A complex vector bundle E of rank r on X consists of the following data:

- (1) open covering $\{U_\alpha\}$ of X .
- (2) smooth functions $\{g_{\alpha\beta}: U_\alpha \cap U_\beta \rightarrow \text{GL}(r, \mathbb{C})\}$ satisfies

$$\begin{aligned}
 g_{\alpha\beta}g_{\beta\gamma}g_{\gamma\alpha} &= \text{id} && \text{on } U_\alpha \cap U_\beta \cap U_\gamma \\
 g_{\alpha\alpha} &= \text{id} && \text{on } U_\alpha
 \end{aligned}$$

Remark 2.4.2. The two definitions above are equivalent. The first definition implies the second clearly. The converse is a standard constructive method: If we already have an open covering and a set of transition functions, the vector bundle E is defined to be the quotient of the disjoint union $\coprod_{U_\alpha} (U \times \mathbb{C}^r)$ by the equivalence relation that puts $(p', v') \in U_\beta \times \mathbb{C}^r$ equivalent to $(p, v) \in U_\alpha \times \mathbb{C}^r$ if and only if $p = p'$ and $v' = g_{\alpha\beta}(p)v$. To connect this definition with the previous one, define the map π to send the equivalence class of any given (p, v) to p .

Definition 2.4.3 (holomorphic vector bundle). A holomorphic vector bundle $\pi: E \rightarrow X$ over a complex manifold X is a complex vector bundle with holomorphic transition functions.

Exercise 2.4.1. Show that the total space of a holomorphic vector bundle E is a complex manifold.

Proof. Since we already have a complex structure on X , we need to pull it back to E using π and use the holomorphic transition functions to show it do gives a complex structure on E . \square

Example 2.4.1 (trivial bundle). Let X be a smooth/complex manifold. Then $X \times \mathbb{C}^r$ is called trivial complex (holomorphic) vector bundle of rank r on X .

Definition 2.4.4 (subbundle). Let $\pi: E \rightarrow X$ be a complex (holomorphic) vector bundle. $F \subseteq E$ is called a subbundle of rank s , if

- (1) For all $x \in X$, $F \cap E_x$ is a subspace of E_x with dimension s .
- (2) $\pi|_F: F \rightarrow X$ induces a complex (holomorphic) vector bundle.

2.4.2. *In viewpoint of sheaf.* One may refer to Appendix 20 for more details about sheaf.

Definition 2.4.5 (section). Let X be a complex manifold and $\pi: E \rightarrow X$ be a complex (holomorphic) vector bundle. For any open subset $U \subseteq X$, a section of E over U is a smooth/holomorphic map $s: U \rightarrow E$ such that $\pi \circ s = \text{id}_U$.

Notation 2.4.1. The set of all smooth (or holomorphic) sections over open subset U is denoted by $C^\infty(U, E)$ (or $\Gamma(U, E)$).

One reason why sheaf plays an important role of study of complex geometry is that you can regard a vector bundle as a special sheaf.

Definition 2.4.6 (sheaf of sections). Let X be a complex manifold and $\pi: E \rightarrow X$ a holomorphic vector bundle. Then its sheaf of sections is defined as

$$\mathcal{O}_X(E)(U) = \Gamma(U, E)$$

Example 2.4.2. Let $E \rightarrow X$ be trivial holomorphic vector bundle. Then $\mathcal{O}_X(E)$ is exactly sheaf of holomorphic functions.

Example 2.4.3 (locally free sheaf). A sheaf \mathcal{F} over a topological space X is called locally free, if there exists an open covering $\{U_\alpha\}$ of X such that $\mathcal{F}|_{U_\alpha} \cong \mathcal{O}_{U_\alpha}^{\oplus r}$ of rank r .

Exercise 2.4.2. Let X be a complex manifold. There is one to one correspondence over X :

$$\{\text{holomorphic vector bundles}\} \xleftrightarrow{1-1} \{\text{locally free sheaves}\}$$

Proof. If $\pi: E \rightarrow X$ is a holomorphic vector bundle, then $\mathcal{O}_X(E)$ is a locally free sheaf. Indeed since we have local trivialization of holomorphic vector bundle $\{U_\alpha\}$, that is $E|_{U_\alpha} \cong U_\alpha \times \mathbb{C}^r$, and it's known to all that sections of a trivial holomorphic function is exactly holomorphic functions, thus $\mathcal{O}(E)|_{U_\alpha} \cong \mathcal{O}_{U_\alpha}^{\oplus r}$, that is $\mathcal{O}_X(E)$ is a locally free sheaf.

Conversely, if \mathcal{E} is locally free over an open covering $\{U_\alpha\}$ of X , then we just need to glue $U_\alpha \times \mathbb{C}^r \rightarrow U_\alpha$ together to get a vector bundle. Therefore we need a family of gluing data $g_{\alpha\beta}: (U_\alpha \cap U_\beta) \times \mathbb{C}^r \rightarrow (U_\alpha \cap U_\beta) \times \mathbb{C}^r$. Since \mathcal{E} is locally free, we have local isomorphism $f_\alpha: \mathcal{E}|_{U_\alpha} \rightarrow \mathcal{O}_{U_\alpha}^{\oplus r}$. Restricting to intersection $U_\alpha \cap U_\beta$, we get

$$f_{\alpha\beta} = f_\alpha|_{U_\alpha \cap U_\beta} \circ f_\beta^{-1}|_{U_\alpha \cap U_\beta}: \mathcal{O}_{U_\beta}^{\oplus r}|_{U_\alpha \cap U_\beta} \rightarrow \mathcal{O}_{U_\alpha}^{\oplus r}|_{U_\alpha \cap U_\beta}$$

Every such map is induced by a map

$$g_{\alpha\beta}: (U_\alpha \cap U_\beta) \times \mathbb{C}^r \rightarrow (U_\alpha \cap U_\beta) \times \mathbb{C}^r$$

that's gluing data we desire. \square

Definition 2.4.7 (cohomology of vector bundle). Let E be a holomorphic vector bundle on a complex manifold X . Then its q -th cohomology $H^q(X, E)$ is defined to be q -th sheaf cohomology of $\mathcal{O}_X(E)$

2.4.3. *Algebraic construction.* Let E, F be complex (holomorphic) vector bundles on X with transition functions $\{g_{\alpha\beta}\}, \{h_{\alpha\beta}\}$ respectively. Then by algebraic construction we have

- (1) $E \oplus F$, given by transition functions $\{\text{diag}(g_{\alpha\beta}, h_{\alpha\beta})\}$
- (2) $E \otimes F$, given by transition functions $\{g_{\alpha\beta} \otimes h_{\alpha\beta}\}$.
- (3) E^* , given by transition functions $\{(g_{\alpha\beta}^{-1})^T\}$.
- (4) $\text{Hom}(E, F) := E^* \otimes F$.
- (5) $\bigwedge^k E$, given by transition functions $\{\bigwedge^k g_{\alpha\beta}\}$.
- (6) Let $f: X \rightarrow Y$ be a smooth/holomorphic map, $\pi: E \rightarrow Y$ is a vector bundle with transition functions $\{g_{\alpha\beta}\}$, then transition functions of pullback bundle f^*E is given by $\{g_{\alpha\beta} \circ f\}$.

Remark 2.4.3. Here is an explicit construction of pullback bundle defined by

$$f^*E = \{(x, e) \in X \times E \mid f(x) = \pi(e)\} \subseteq X \times E$$

In fact, you can regard it as a push out as

$$\begin{array}{ccc} f^*E & \longrightarrow & E \\ \downarrow & & \downarrow \pi \\ X & \xrightarrow{f} & Y \end{array}$$

In particular, pullback bundle has universal property.

2.4.4. *Hermitian structure.* Let X be a smooth manifold and $\pi: E \rightarrow X$ be a complex vector bundle.

Definition 2.4.8 (Hermitian metric). A Hermitian metric h on E is a global section of $E^* \otimes \bar{E}$.

Remark 2.4.4 (local form). Let $\{e_\alpha\}$ be a local frame of E . Then Hermitian metric h is given by

$$h = h_{\alpha\bar{\beta}} e^\alpha \otimes \bar{e}^\beta$$

where $h_{\alpha\bar{\beta}}$ is a Hermitian matrix.

Proposition 2.4.1. Every complex vector bundle admits a Hermitian metric.

Proof. Use partition of unity. \square

2.4.5. *Line bundle.*

Definition 2.4.9 (line bundle). A complex (or holomorphic) line bundle L is a complex (or holomorphic) vector bundle of rank one.

Proposition 2.4.2. Let L be a complex line bundle over X . Then $L \otimes L^*$ is the trivial bundle.

Proof. Suppose $\{g_{\alpha\beta}\}$ is the transition functions of L , by Section 2.4.3 it's clear to see the transition functions of $L^* \otimes L$ is

$$(g_{\alpha\beta}^{-1})^T g_{\alpha\beta} = g_{\alpha\beta}^{-1} g_{\alpha\beta} = \text{id}$$

This completes the proof. \square

Proposition 2.4.3. Let L be a holomorphic over a compact complex manifold X . Then L is trivial if and only if both L and its dual L^* admit non-trivial global section.

Proposition 2.4.4. Let $\pi: E \rightarrow X$ be a complex line bundle. Then E is a trivial line bundle if and only if there exists a nowhere vanishing global section s .

Proof. It's clear there exists a nowhere vanishing global section if E is trivial. Conversely, if there exists a nowhere vanishing global section s . Consider the following map

$$\begin{aligned} \varphi: X \times \mathbb{C} &\rightarrow E \\ (x, \lambda) &\mapsto \lambda s(x) \end{aligned}$$

It's an isomorphism since fiberwisely one has $\varphi_x(\lambda) = \lambda s(x)$, and it's injective thus isomorphism since $s(x) \neq 0$. \square

Definition 2.4.10 (picard group). The picard group $\text{Pic}(X)$ denotes set of all holomorphic line bundles on X up to isomorphism, whose group structure is given by tensor product.

Proposition 2.4.5. There is a natural isomorphism $\text{Pic}(X) \cong H^1(X, \mathcal{O}_X^*)$.

Proof. For a line bundle L , it's completely determined by its transition functions $g_{\alpha\beta}: U_{\alpha\beta} \rightarrow \mathbb{C}^*$, which is holomorphic functions. It gives rise to an element in $\check{H}^1(X, \mathcal{O}_X^*)$ since $g_{\alpha\beta}$ satisfies cocycle conditions. Furthermore, Čech cohomology³ computes the sheaf cohomology for reasonable topological space, e.g. for manifolds. \square

Remark 2.4.5. This proposition gives us a method to compute Picard group of a complex manifold since there is exponential sequence as follows

$$0 \rightarrow \mathbb{Z} \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_X^* \rightarrow 0$$

which is a exact sequence of sheaves, then it gives a long exact sequence of cohomology groups as follows

$$\dots \rightarrow H^1(X, \mathbb{Z}) \rightarrow H^1(X, \mathcal{O}_X) \rightarrow H^1(X, \mathcal{O}_X^*) \rightarrow H^2(X, \mathbb{Z}) \rightarrow \dots$$

Thus $\text{Pic}(X)$ can in principle be computed by above exact sequence. Roughly speaking, $\text{Pic}(X)$ has two parts:

- (1) A discrete part, measured by its image in $H^2(X, \mathbb{Z})$.
- (2) A continuous part coming from the $H^1(X, \mathcal{O}_X)$, which is possibly trivial.

Proposition 2.4.6. The set $\mathcal{O}_{\mathbb{C}\mathbb{P}^n}(-1) \subseteq \mathbb{C}\mathbb{P}^n \times \mathbb{C}^{n+1}$ that consists of all pairs $(l, z) \in \mathbb{C}\mathbb{P}^n \times \mathbb{C}^{n+1}$ with $z \in l$ forms a holomorphic line bundle, called tautological line bundle

Proof. Let $\pi: \mathcal{O}_{\mathbb{C}\mathbb{P}^n}(-1) \rightarrow \mathbb{C}\mathbb{P}^n$ be the projection to the first factor. Consider open covering $\{U_i\}_{i=0}^n$ of $\mathbb{C}\mathbb{P}^n$, where

$$U_i = \{[l] = [l_0 : \dots : l_n] \in \mathbb{C}\mathbb{P}^n \mid l_i \neq 0\}$$

A canonical trivialization of $\mathcal{O}_{\mathbb{C}\mathbb{P}^n}(-1)$ over U_i is given by

$$\begin{aligned} \varphi_i: \pi^{-1}(U_i) &\rightarrow U_i \times \mathbb{C} \\ (l, z) &\mapsto (l, z_i) \end{aligned}$$

Its transition function is computed as follows

$$\begin{aligned} \varphi_i \circ \varphi_j^{-1}: (U_i \cap U_j) \times \mathbb{C} &\longrightarrow (U_i \cap U_j) \times \mathbb{C} \\ (l, w) &\mapsto (l, w \frac{l_i}{l_j}) \end{aligned}$$

where $l = (l_0 : \dots : l_n)$. This shows its transition function $g_{ij}(z) = z_i/z_j \in \mathbb{C}^*$ is holomorphic. \square

Definition 2.4.11 (line bundles on $\mathbb{C}\mathbb{P}^n$).

$$\begin{cases} \mathcal{O}_{\mathbb{C}\mathbb{P}^n}(-k) = \mathcal{O}_{\mathbb{C}\mathbb{P}^n}(-1)^{\otimes k} & k \in \mathbb{Z}_{>0} \\ \mathcal{O}_{\mathbb{C}\mathbb{P}^n}(k) = (\mathcal{O}_{\mathbb{C}\mathbb{P}^n}(-k))^* & k \in \mathbb{Z}_{>0} \\ \mathcal{O}_{\mathbb{C}\mathbb{P}^n}(0) = \mathbb{C}\mathbb{P}^n \times \mathbb{C} \end{cases}$$

³For more details, see Appendix 20.

Proposition 2.4.7. For $k \geq 0$. the space $\Gamma(\mathbb{CP}^n, \mathcal{O}_{\mathbb{CP}^n}(k))$ is canonically isomorphic to the space $\mathbb{C}[z_0, \dots, z_n]_k$ of all homogenous polynomials of degree k .

Corollary 2.4.1. For $k < 0$ the line bundle $\mathcal{O}_{\mathbb{CP}^n}(k)$ admits no global holomorphic section.

Proof. It follows from above result and Proposition 2.4.3. □

2.5. Euler sequence and adjunction formula.

2.5.1. Euler sequence.

Proposition 2.5.1. On \mathbb{CP}^n there exists a natural short exact sequence of holomorphic vector bundles

$$0 \rightarrow \mathcal{O}_{\mathbb{CP}^n} \xrightarrow{\phi} \mathcal{O}_{\mathbb{CP}^n}(1)^{\oplus n+1} \xrightarrow{\psi} T\mathbb{CP}^n \rightarrow 0$$

Exercise 2.5.1. For Grassmannian manifold $Gr(k, n)$, we have

$$0 \rightarrow E \rightarrow Gr(k, n) \otimes \mathbb{C}^n \rightarrow Q \rightarrow 0$$

Show that

$$T_{Gr(k,n)} \cong \text{Hom}(E, Q)$$

2.5.2. Adjunction formula.

Proposition 2.5.2 (adjunction formula). Let $\pi: L \rightarrow X$ is a holomorphic line bundle and s be a holomorphic section of L . Suppose that $D = \{x \in X \mid s(x) = 0\}$ is a smooth submanifold of codimensional 1. Show that the following sequence is exact

$$0 \rightarrow TD \rightarrow TX|_D \rightarrow L|_D \rightarrow 0$$

As a consequence

$$K_D^* \cong K_D^* \otimes L|_D = (K_X^* \otimes L)|_D$$

Or equivalently

$$K_D \cong (K_X \otimes L)|_D$$

This is called adjunction formula.

Proof. Firstly we have the following exact sequence

$$0 \rightarrow TD \rightarrow TX|_D \rightarrow N_D \rightarrow 0$$

where N_D is the normal bundle. Now it suffices to show $L|_D$ is isomorphic to the normal bundle of D . Note that $s|_D = 0$ and s is not identically zero on X , so ds gives an isomorphism between $L|_D$ and N_D in fact. By taking determinant we obtain the adjunction formula since for a exact sequence of vector bundle

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$$

we have

$$\det B = \det A \otimes \det C$$

and determinant of a line bundle is itself. □

Example 2.5.1. Let $X = \mathbb{CP}^n$ and $L = \mathcal{O}_{\mathbb{CP}^n}(-d)$. Proposition 2.4.7 shows that $D \subseteq \mathbb{CP}^n$ is a smooth hypersurface defined by zero set of a homogenous polynomial with degree d . Then we have

$$\begin{aligned} K_D^* &\cong (K_X^* \otimes L)|_D \\ &= (\mathcal{O}_{\mathbb{CP}^n}(n+1) \otimes \mathcal{O}_{\mathbb{CP}^n}(-d))|_D \\ &\cong \mathcal{O}_{\mathbb{CP}^n}(n+1-d)|_D \end{aligned}$$

Remark 2.5.1. As a consequence, D is called

$$\begin{cases} \text{Fano} & d < n+1 \\ \text{Calabi-Yau} & d = n+1 \\ \text{general type} & d > n+1 \end{cases}$$

These concepts we will define later.

3. DIVISOR AND LINE BUNDLE

In this section, unless otherwise specified, we assume X is a complex manifold.

3.1. Divisor.

Definition 3.1.1 (analytic hypersurface). An analytic hypersurface of X is an analytic subvariety $Y \subseteq X$ of codimensional one.

Remark 3.1.1. By Remark 1.4.2 one has a hypersurface is locally given as the zero set of a non-trivial holomorphic function.

Definition 3.1.2 (divisor). A divisor D on X is a locally finite⁴ formal linear combination $D = \sum a_i[Y_i]$ with $Y_i \subseteq X$ are irreducible hypersurfaces and $a_i \in \mathbb{Z}$.

Definition 3.1.3 (divisor group). The divisor group $\text{Div}(X)$ is the set of all divisors endowed with the natural group structure.

Definition 3.1.4 (effective). A divisor $D = \sum a_i[Y_i]$ is called effective, if $a_i \geq 0$ for all i . In this case, we write $D \geq 0$.

Proposition 3.1.1. Every hypersurfaces Y defines an effective divisor $\sum[Y_i] \in \text{Div}(X)$, where Y_i are irreducible components of Y .

Proof. It suffices to show the irreducible components of a hypersurface Y is locally finite. \square

Let $Y \subseteq X$ be a hypersurface and $x \in Y$. Suppose that Y defines an irreducible germ in x , that is this germ is the zero set of an irreducible $g \in \mathcal{O}_{X,x}$.

Definition 3.1.5 (order). Let f be a meromorphic function in a neighborhood of $x \in Y$. Then the order $\text{ord}_{Y,x}(f)$ of f in x with respect to Y is given by

$$f = g^{\text{ord}_{Y,x}(f)} h$$

where $h \in \mathcal{O}_{X,x}^*$.

Remark 3.1.2.

- (1) The order of f in x with respect to Y is independent of the choice of g since any two irreducible $g, g' \in \mathcal{O}_{X,x}$ with $Z(g) = Z(g')$ only differs by an element in $\mathcal{O}_{X,x}^*$.
- (2) More globally, one can define order $\text{ord}_Y(f)$ as $\text{ord}_Y(f) = \text{ord}_{Y,x}(f)$ for $x \in Y$ such that Y defines an irreducible germ in x . Such a point $x \in Y$ always exists, for example, one can choose a regular point $x \in Y_{\text{reg}}$. Moreover, it's independent of the choice of x since

⁴The sum is called locally finite, if for any $x \in X$, there exists an open neighborhood $U \subseteq X$ such that only finite many coefficients $a_i \neq 0$ with $Y_i \cap U \neq \emptyset$.

Definition 3.1.6 (zeros and poles). Let f be a meromorphic function on X . Then

- (1) f has zeros of order $d \geq 0$ along Y if $\text{ord}_Y(f) = d$.
- (2) f has poles of order $d \geq 0$ along Y if $\text{ord}_Y(f) = -d$.

Definition 3.1.7 (principal divisor). For $f \in K(X)$, the divisor associated to f is

$$(f) := \sum \text{ord}_Y(f)[Y]$$

where the sum is taken over all irreducible hypersurfaces $Y \subseteq X$. A divisor of this form is called principal.

Remark 3.1.3. The divisor (f) can be written as the difference of two effective divisors $(f) = Z(f) - P(f)$, where

$$Z(f) = \sum_{\text{ord}_Y(f) > 0} \text{ord}_Y(f)[Y], \quad P(f) = \sum_{\text{ord}_Y(f) < 0} \text{ord}_Y(f)[Y]$$

Proposition 3.1.2. There exists a natural isomorphism

$$H^0(X, \mathcal{K}_X^*/\mathcal{O}_X^*) \cong \text{Div}(X)$$

Proof. An element $f \in H^0(X, \mathcal{K}_X^*/\mathcal{O}_X^*)$ is given by non-trivial meromorphic functions $f_i \in K_X^*(U_i)$ such that $f_i f_j^{-1}$ is a holomorphic function without zeros on $U_i \cap U_j$, where $\{U_i\}$ is an open covering of X . Thus for any irreducible hypersurface $Y \subseteq X$ with $Y \cap U_i \cap U_j \neq \emptyset$, one has $\text{ord}_Y(f_i) = \text{ord}_Y(f_j)$. Hence $\text{ord}_Y(f)$ is well-defined for any irreducible hypersurface Y . Then one associates to f the divisor $(f) = \sum \text{ord}_Y(f)[Y] \in \text{Div}(X)$.

It's clear this map is a group homomorphism. To see it's bijective, we define the inverse as follows. If $D = \sum a_i[Y_i] \in \text{Div}(X)$ is given, then there exists an open covering $\{U_i\}$ of X such that $Y_i \cap U_j$ is defined by $g_{ij} \in \mathcal{O}(U_j)$ which is unique up to elements in $\mathcal{O}^*(U_j)$. Let $f_j := \prod_i g_{ij}^{a_i} \in \mathcal{K}_X^*(U_j)$ since g_{ij} and g_{ik} defines the same irreducible hypersurface, they only differ by an element in $\mathcal{O}^*(U_j \cap U_k)$. Thus f glue to an element $f \in H^0(X, \mathcal{K}_X^*/\mathcal{O}_X^*)$. It's clear these two maps are inverse to each other. \square

Remark 3.1.4. In algebraic geometry, elements in $H^0(X, \mathcal{K}_X^*/\mathcal{O}_X^*)$ are called Cartier divisors and elements in $\text{Div}(X)$ are called Weil divisors. Above isomorphism still holds in the algebraic setting under a weak smoothness assumption on X .

Corollary 3.1.1. There exists a natural group homomorphism

$$\begin{aligned} \text{Div}(X) &\rightarrow \text{Pic}(X) \\ D &\mapsto \mathcal{O}(D) \end{aligned}$$

where $\mathcal{O}(D)$ is defined in the proof.

Proof. If $D = \sum a_i[Y_i] \in \text{Div}(X)$ corresponds to $f \in H^0(X, \mathcal{K}_X^*/\mathcal{O}_X^*)$, which in turn is given by functions $f_i \in \mathcal{K}_X^*(U_i)$ for an open covering $\{U_i\}$. Then we define $\mathcal{O}(D) \in \text{Div}(X)$ with transition functions $\psi_{ij} := f_i f_j^{-1} \in \mathcal{O}_X^*(U_{ij})$.

If D, D' are two divisors, without loss of generality we may assume they're given by $\{f_i\}$ and $\{f'_i\}$ respectively on the same open covering, then $D + D'$, then $D + D'$ corresponds to $\{f_i + f'_i\}$. By definition $\mathcal{O}(D + D')$ is described by $\{\psi_{ij}\psi'_{ij}\}$, hence $\mathcal{O}(D + D') = \mathcal{O}(D) \otimes \mathcal{O}(D')$. \square

Remark 3.1.5. In fact, above corollary can be derived from the following exact sequence of sheaves

$$0 \rightarrow \mathcal{O}_X^* \rightarrow \mathcal{K}_X^* \rightarrow \mathcal{K}_X^*/\mathcal{O}_X^* \rightarrow 0$$

Then above group homomorphism is exactly the boundary map, the kernel of which coincides with the image of $H^0(X, \mathcal{K}_X^*) \rightarrow H^0(X, \mathcal{K}_X^*/\mathcal{O}_X^*)$, and the latter by definition is the set of principal divisors.

Definition 3.1.8 (linearly equivalent). Two divisors D, D' are called linearly equivalent, denoted by $D \sim D'$, if $D - D'$ is a principal divisor.

Corollary 3.1.2. The group homomorphism $\text{Div}(X) \rightarrow \text{Pic}(X)$ factorizes over an injection

$$\text{Div}(X)/\sim \hookrightarrow \text{Pic}(X)$$

3.2. Relations between divisor and line bundle. In general, $\text{Div}(X)/\sim \hookrightarrow \text{Pic}(X)$ is a strict inclusion, but we will see if a line bundle admits a non-trivial global section, then it's contained in the image. In order to show this, we need to construct a canonical map

$$\begin{aligned} H^0(X, L) \setminus \{0\} &\rightarrow \text{Div}(X) \\ s &\mapsto Z(s) \end{aligned}$$

The map is constructed as follows: Let $L \in \text{Pic}(X)$ on open covering $\{U_i\}$ be trivialized by $\psi_i: L|_{U_i} \rightarrow \mathcal{O}_{U_i}$. Then divisor $Z(s)$ is given by $f := \{f_i := \psi_i(s|_{U_i}) \in H^0(X, \mathcal{K}_X^*/\mathcal{O}_X^*)\}$.

Proposition 3.2.1. For $0 \neq s \in H^0(X, L)$, the line bundle $\mathcal{O}(Z(s))$ is isomorphic to L .

Proposition 3.2.2. For any effective divisor $D \in \text{Div}(X)$, there exists a section $0 \neq s \in H^0(X, \mathcal{O}(D))$ with $Z(s) = D$.

Corollary 3.2.1. Non-trivial sections $s_1 \in H^0(X, L_1)$ and $s_2 \in H^0(X, L_2)$ define linearly equivalent divisors $Z(s_1) \sim Z(s_2)$ if and only if $L_1 \cong L_2$.

Proof. If $L_1 \cong L_2$, then

If $Z(s_1) \sim Z(s_2)$, then by Corollary 3.1.2 one has $\mathcal{O}(Z(s_1)) \cong \mathcal{O}(Z(s_2))$, then this shows $L_1 \cong L_2$ since $\mathcal{O}(Z(s_i)) = L_i, i = 1, 2$. \square

Corollary 3.2.2. The image of the natural map $\text{Div}(X) \rightarrow \text{Pic}(X)$ is generated by those line bundles $L \in \text{Pic}(X)$ with $H^0(X, L) \neq 0$.

Proof. We have already seen if $H^0(X, L) \neq 0$, then L is contained in the image. Conversely, any divisor $D = \sum a_i[Y_i]$ can be written as $D = \sum a_i^+[Y_i] - \sum a_j^-[Y_j]$ with $a_k^\pm \geq 0$, and thus $\mathcal{O}(D) \cong \mathcal{O}(\sum a_i^+[Y_i]) \otimes \mathcal{O}(\sum a_j^-[Y_j])^*$. Both

$\mathcal{O}(\sum a_i^+[Y_i])$ and $\mathcal{O}(\sum a_j^-[Y_j])$ are associated to effective divisors, and therefore admit non-trivial global sections. \square

Remark 3.2.1. For projective manifolds, the map $\text{Div}(X) \rightarrow \text{Pic}(X)$ is surjective, but note that even for very easy manifolds, such as complex tori, this is no longer the case.

3.3. Ample line bundle.

Definition 3.3.1 (base point). Let L be a holomorphic line bundle on a complex manifold X . A point $x \in X$ is a base point of L if $s(x) = 0$ for all $s \in H^0(X, L)$.

Notation 3.3.1. The base locus $\text{Bs}(L)$ is the set of all base points of L .

Remark 3.3.1. If $\dim H^0(X, L) < \infty$, we can choose a basis of global sections s_1, \dots, s_N of it, then $\text{Bs}(L) = Z(s_1) \cap \dots \cap Z(s_N)$ is an analytic subvariety. Later we will see if X is compact, then $\dim H^0(X, L) < \infty$.

Proposition 3.3.1. Let L be a holomorphic line bundle on a complex manifold X and suppose $s_1, \dots, s_N \in H^0(X, L)$ is a basis. Then

$$\begin{aligned} \varphi_L: X \setminus \text{Bs}(L) &\rightarrow \mathbb{C}\mathbb{P}^N \\ x &\mapsto (s_0(x) : \dots : s_N(x)) \end{aligned}$$

defines a holomorphic map such that $\varphi_L^* \mathcal{O}_{\mathbb{C}\mathbb{P}^N}(-1) \cong L|_{X \setminus \text{Bs}(L)}$.

Definition 3.3.2 (ample line bundle). A holomorphic line bundle L on a complex manifold X is called ample if for some $k > 0$ and some linear system in $H^0(X, L^k)$ the associated map φ is an embedding.

Remark 3.3.2. By definition, a compact complex manifold is projective if and only if it admits an ample line bundle.

4. TANGENT AND COTANGENT BUNDLE

4.1. Complex and holomorphic tangent bundle.

Definition 4.1.1 (complex tangent bundle). Let X be a smooth n -manifold with an atlas $\{U_\alpha, \varphi_\alpha: U_\alpha \rightarrow V_\alpha \subseteq \mathbb{R}^n\}$. Then (real) tangent bundle $T_{\mathbb{R}}X$ is a vector bundle given by smooth transition functions

$$g_{\alpha\beta}: U_\alpha \cap U_\beta \rightarrow \mathrm{GL}(n, \mathbb{R})$$

$$x \mapsto J_{\mathbb{R}}(\varphi_\alpha \circ \varphi_\beta^{-1})(\varphi_\beta(x))$$

The complex tangent bundle $T_{\mathbb{C}}X$ is defined as the complexification of $T_{\mathbb{R}}X$, that is, $T_{\mathbb{R}}X \otimes_{\mathbb{R}} \mathbb{C}$.

Definition 4.1.2 (holomorphic tangent bundle). Let X be a complex n -manifold, with an atlas $\{U_\alpha, \varphi_\alpha: U_\alpha \rightarrow V_\alpha \subseteq \mathbb{C}^n\}$. Then holomorphic tangent bundle TX is given by holomorphic transition functions

$$g_{\alpha\beta}: U_\alpha \cap U_\beta \rightarrow \mathrm{GL}(n, \mathbb{C})$$

$$z \mapsto J_{\mathbb{C}}(\varphi_\alpha \circ \varphi_\beta^{-1})(\varphi_\beta(z))$$

Remark 4.1.1 (relations between complex tangent bundle and holomorphic tangent bundle). Let X be a complex n -manifold and $\{z^i = x^i + \sqrt{-1}y^i\}_{1 \leq i \leq n}$ be a local coordinate of X . Then $\{x^1, \dots, x^n, y^1, \dots, y^n\}$ gives a local coordinate of its underlying real $2n$ -manifold, and there is an almost complex structure J on $T_{\mathbb{R}}X$ given by

$$J\left(\frac{\partial}{\partial x^i}\right) = \frac{\partial}{\partial y^i}$$

$$J\left(\frac{\partial}{\partial y^i}\right) = -\frac{\partial}{\partial x^i}$$

Thus complex tangent bundle $T_{\mathbb{C}}X$ can be decomposed as $T_{\mathbb{C}}X = T^{1,0}X \oplus T^{0,1}X$ with respect to J , with local frames as follows:

- (1) $\{\frac{\partial}{\partial z^i} := \frac{1}{2}(\frac{\partial}{\partial x^i} - \sqrt{-1}\frac{\partial}{\partial y^i})\}$ is a local frame of $T^{1,0}X$.
- (2) $\{\frac{\partial}{\partial \bar{z}^i} := \frac{1}{2}(\frac{\partial}{\partial x^i} + \sqrt{-1}\frac{\partial}{\partial y^i})\}$ is a local frame of $T^{0,1}X$.

Indeed, direct computation shows

$$J\left(\frac{\partial}{\partial z^i}\right) = \frac{1}{2}\left(\frac{\partial}{\partial y^i} + \sqrt{-1}\frac{\partial}{\partial x^i}\right) = \frac{\sqrt{-1}}{2}\left(\frac{\partial}{\partial x^i} - \sqrt{-1}\frac{\partial}{\partial y^i}\right) = \sqrt{-1}\frac{\partial}{\partial z^i}$$

$$J\left(\frac{\partial}{\partial \bar{z}^i}\right) = \frac{1}{2}\left(\frac{\partial}{\partial y^i} - \sqrt{-1}\frac{\partial}{\partial x^i}\right) = -\frac{\sqrt{-1}}{2}\left(\frac{\partial}{\partial x^i} + \sqrt{-1}\frac{\partial}{\partial y^i}\right) = -\sqrt{-1}\frac{\partial}{\partial \bar{z}^i}$$

and for any section s of $T_{\mathbb{C}}X$, one has the following decomposition

$$s = \frac{1}{2}(s - \sqrt{-1}J(s)) + \frac{1}{2}(s + \sqrt{-1}J(s))$$

Note that TX is isomorphic to $(T_{\mathbb{R}}X, J)$ as complex vector bundle, and we claim $(T_{\mathbb{R}}X, J)$ is isomorphic to $T^{1,0}X$ as a complex vector bundle. Indeed, there is a natural inclusion $T_{\mathbb{R}}X \hookrightarrow T_{\mathbb{C}}X$, if we compose this inclusion with

projection $T_{\mathbb{C}}X = T^{1,0}X \oplus T_X^{0,1} \rightarrow T^{1,0}X$ onto the first summand, we obtain an \mathbb{C} -isomorphism $(T_{\mathbb{R}}X, J) \rightarrow T^{1,0}X$ with inverse map $2\text{Re}(-)$.

In particular, $T^{1,0}X$ is isomorphic to TX as complex vector bundles, so we can endow $T^{1,0}X$ with holomorphic structure such that $T^{1,0}X$ is isomorphic to TX as holomorphic vector bundles, and thus we can use $\{dz^i\}$ as local frame of holomorphic tangent bundle.

4.2. Bidegree forms. For complex manifold X , there is also an almost complex structure on $\Omega_{X,\mathbb{R}}^1$, that is dual bundle of $T_{\mathbb{R}}X$, and complexified dual space of $T_{\mathbb{R}}X$ admits an analogous decomposition:

$$\Omega_{X,\mathbb{C}}^1 = \Omega_{X,\mathbb{R}}^1 \otimes \mathbb{C} = \Omega_X^{1,0} \oplus \Omega_X^{0,1}$$

There is also a decomposition on its k -th wedge product as follows:

$$\Omega_{X,\mathbb{C}}^k = \bigwedge^k \Omega_{X,\mathbb{C}}^1 = \bigoplus_{p+q=k} \Omega_X^{p,q}$$

where $\Omega_X^{p,q} = \bigwedge^p \Omega_X^{1,0} \otimes \bigwedge^q \Omega_X^{0,1}$.

Definition 4.2.1 ((p, q) -form). A k -form ω of type (p, q) is a smooth section of $\Omega_X^{p,q}$, that is

$$\omega \in C^\infty(X, \Omega_X^{p,q}) \subseteq C^\infty(X, \Omega_{X,\mathbb{C}}^k)$$

Remark 4.2.1 (local form). Suppose $\{z^1, \dots, z^n\}$ is a local coordinate of X , and denote $z^i = x^i + \sqrt{-1}y^i$. Then $\{dx^1, \dots, dx^n, dy^1, \dots, dy^n\}$ gives a local frame of $\Omega_{X,\mathbb{R}}^1$, and induced almost complex structure is given by

$$\begin{aligned} J^*(dx^i)\left(\frac{\partial}{\partial x^i}\right) &= dx^i\left(J\left(\frac{\partial}{\partial x^i}\right)\right) = dx^i\left(\frac{\partial}{\partial y^i}\right) = 0 \\ J^*(dx^i)\left(\frac{\partial}{\partial y^i}\right) &= dx^i\left(J\left(\frac{\partial}{\partial y^i}\right)\right) = dx^i\left(-\frac{\partial}{\partial x^i}\right) = -1 \end{aligned}$$

that is

$$\begin{aligned} J^*(dx^i) &= -dy^i \\ J^*(dy^i) &= dx^i \end{aligned}$$

and similarly we have

- (1) $\{dz^i := dx^i + \sqrt{-1}dy^i\}$ is a local frame of $\Omega_X^{1,0}$.
- (2) $\{d\bar{z}^i := dx^i - \sqrt{-1}dy^i\}$ is a local frame of $\Omega_X^{0,1}$.

For a k -form, it locally looks like

$$\sum_{\substack{|I|=p, |J|=q \\ p+q=k}} f_{IJ} dz^I \wedge d\bar{z}^J$$

where f_{IJ} are smooth functions, and a k -form is a (p, q) -form if and only if locally it looks like

$$\sum_{|I|=p, |J|=q} f_{IJ} dz^I \wedge d\bar{z}^J$$

Exercise 4.2.1. For $\mathbb{C}^n \cong \mathbb{R}^{2n}$, one has

$$dx^1 \wedge dy^1 \wedge \cdots \wedge dx^n \wedge dy^n = \left(\frac{\sqrt{-1}}{2}\right)^n dz^1 \wedge d\bar{z}^1 \wedge \cdots \wedge dz^n \wedge d\bar{z}^n$$

Proof. It suffices to show the case $n = 1$, and we can compute directly as follows

$$\begin{aligned} \left(\frac{\sqrt{-1}}{2}\right) dz \wedge d\bar{z} &= \left(\frac{\sqrt{-1}}{2}\right) (dx + \sqrt{-1}dy) \wedge (dx - \sqrt{-1}dy) \\ &= \left(\frac{\sqrt{-1}}{2}\right) (-2\sqrt{-1}dx \wedge dy) \\ &= dx \wedge dy \end{aligned}$$

□

4.3. Dolbeault operators. For complex manifold X , naturally there is a differential operator

$$d: C^\infty(X, \Omega_{X,\mathbb{C}}^k) \rightarrow C^\infty(X, \Omega_{X,\mathbb{C}}^{k+1})$$

Since there is a decomposition for $\alpha \in C^\infty(X, \Omega_{X,\mathbb{C}}^k)$, it's natural to ask how to decompose $d\alpha \in C^\infty(X, \Omega_{X,\mathbb{C}}^{k+1})$.

Example 4.3.1. For smooth function α on X , locally a direct computation shows

$$\begin{aligned} d\alpha &= \frac{\partial\alpha}{\partial x^i} dx^i + \frac{\partial\alpha}{\partial y^i} dy^i \\ &= \frac{1}{2} \left(\frac{\partial\alpha}{\partial x^i} - \sqrt{-1} \frac{\partial\alpha}{\partial y^i} \right) dz^i + \frac{1}{2} \left(\frac{\partial\alpha}{\partial x^i} + \sqrt{-1} \frac{\partial\alpha}{\partial y^i} \right) d\bar{z}^i \\ &= \frac{\partial\alpha}{\partial z^i} dz^i + \frac{\partial\alpha}{\partial \bar{z}^i} d\bar{z}^i \end{aligned}$$

If we denote

$$\begin{aligned} \partial\alpha &= \frac{\partial\alpha}{\partial z^i} dz^i \\ \bar{\partial}\alpha &= \frac{\partial\alpha}{\partial \bar{z}^i} d\bar{z}^i \end{aligned}$$

then $d\alpha = \partial\alpha + \bar{\partial}\alpha$, where $\partial\alpha \in C^\infty(X, \Omega_X^{1,0})$ and $\bar{\partial}\alpha \in C^\infty(X, \Omega_X^{0,1})$. In general case, for $\alpha \in C^\infty(X, \Omega_X^{p,q})$, locally looks like

$$\alpha = \sum_{|I|=p, |J|=q} \alpha_{IJ} dz^J \wedge d\bar{z}^K$$

then

$$d\alpha = \sum_{|I|=p, |J|=q} \frac{\partial\alpha_{IJ}}{\partial z^l} dz^l \wedge dz^I \wedge d\bar{z}^J + \sum_{|I|=p, |J|=q} \frac{\partial\alpha_{IJ}}{\partial \bar{z}^l} d\bar{z}^l \wedge z^I \wedge \bar{z}^J$$

Thus one can define

$$\begin{aligned} \partial: C^\infty(X, \Omega_X^{p,q}) &\rightarrow C^\infty(X, \Omega_X^{p+1,q}) \\ \bar{\partial}: C^\infty(X, \Omega_X^{p,q}) &\rightarrow C^\infty(X, \Omega_X^{p,q+1}) \end{aligned}$$

such that $d = \partial + \bar{\partial}$.

Proposition 4.3.1.

(1)

$$\partial(\alpha \wedge \beta) = \partial\alpha \wedge \beta + (-1)^{\deg \alpha} \alpha \wedge \partial\beta$$

(2)

$$\partial^2 = \bar{\partial}^2 = 0, \quad \partial\bar{\partial} + \bar{\partial}\partial = 0$$

According to (2) of Proposition 4.3.1 there is following cochain complex

$$(4.1) \quad 0 \rightarrow C^\infty(X, \Omega_X^{p,0}) \xrightarrow{\bar{\partial}} C^\infty(X, \Omega_X^{p,1}) \xrightarrow{\bar{\partial}} \dots \xrightarrow{\bar{\partial}} C^\infty(X, \Omega_X^{p,n}) \rightarrow 0$$

Definition 4.3.1 (Dolbeault cohomology).

$$H^{p,q}(X) := H_{\bar{\partial}}^q(C^\infty(X, \Omega_X^{p,\bullet}))$$

Remark 4.3.1. Note that we have decomposition $C^\infty(X, \Omega_{X,\mathbb{C}}^k) = \bigoplus_{p+q=k} C^\infty(X, \Omega_X^{p,q})$, could we have the following decomposition

$$H^k(X, \mathbb{C}) = \bigoplus_{p+q=k} H^{p,q}(X)$$

In fact, for compact Kähler manifold, such decomposition do holds, which is called Hodge decomposition.

Example 4.3.2. Note that

$$H^{p,0}(X) = \{\alpha \in C^\infty(X, \Omega_X^{p,0}) \mid \bar{\partial}\alpha = 0\}$$

For $\alpha \in C^\infty(X, \Omega_X^{p,0})$ locally written as $\alpha = \sum_{|I|=p} \alpha_I dz^I$, one has

$$\bar{\partial}\alpha = \sum_{|I|=p} \frac{\partial \alpha_I}{\partial \bar{z}^l} d\bar{z}^l \wedge dz^I = 0 \iff \frac{\partial \alpha_I}{\partial \bar{z}^l} = 0$$

which implies α_I is a holomorphic function. This shows $H^{p,0}(X) = \Gamma(X, \Omega_X^p)$.

A natural question arises: what does this cohomology compute? In the context of smooth manifolds, de Rham cohomology calculates the cohomology of a constant sheaf, which heavily relies on the Poincaré lemma. In the complex setting, Dolbeault cohomology $H^{p,q}(X)$ determines the q -th sheaf cohomology of Ω_X^p , which is based on the following lemma.

Proposition 4.3.2 ($\bar{\partial}$ -Poincaré lemma). Let B be an sufficiently small open disc in \mathbb{C}^n . If $\alpha \in C^\infty(B, \Omega_X^{p,q})$ is $\bar{\partial}$ -closed and $q > 0$, then there exists $\beta \in C^\infty(B, \Omega_X^{p,q-1})$ such that $\alpha = \bar{\partial}\beta$.

Proof. See Corollary 1.3.9 of Page47 of [Huy05]. \square

Proposition 4.3.3 (functorial). Let $f: X \rightarrow Y$ be a holomorphic map between complex manifolds with pullback

$$f^*: C^\infty(Y, \Omega_{Y,\mathbb{C}}^k) \rightarrow C^\infty(X, \Omega_{X,\mathbb{C}}^k)$$

Then

$$f^*: C^\infty(Y, \Omega_{Y, \mathbb{C}}^{p,q}) \rightarrow C^\infty(X, \Omega_{X, \mathbb{C}}^{p,q})$$

and it induces

$$f^*: H^{p,q}(Y) \rightarrow H^{p,q}(X)$$

Example 4.3.3 (Dolbeault cohomology of a holomorphic vector bundle⁵). For a holomorphic vector bundle $E \rightarrow X$, we can also define

$$\bar{\partial}_E: C^\infty(X, \Omega_X^{0,q} \otimes E) \rightarrow C^\infty(X, \Omega_X^{0,q+1} \otimes E)$$

satisfies $\bar{\partial}_E^2 = 0$. Let's elaborate this construction: Since any global section is glued together by local sections, we just need to define $\bar{\partial}_E$ for local sections and check it is well-defined under the change of local chart. We can choose a local holomorphic frame $\{e^1, \dots, e^n\}$ for E on U , so any section $s \in C^\infty(U, \Omega_X^{0,q} \otimes E)$ locally can be written as $s = s^i \otimes e_i$ with $s^i \in C^\infty(U, \Omega_X^{0,q})$. Then we can define

$$\bar{\partial}_E(s) = \bar{\partial}s^i \otimes e^i$$

It's clear that this definition is independent of the choice of local chart since the transition functions are holomorphic and $\bar{\partial}$ kills them. Furthermore, $\bar{\partial}_E^2 = 0$ holds since $\bar{\partial}^2 = 0$. Thus we can construct a cochain complex and define its cohomology, denoted by

$$H^q(X, E) = H_{\bar{\partial}_E}^q(C^\infty(X, \Omega_X^{0,\bullet} \otimes E))$$

and similarly $H^q(X, E)$ computes the q -th sheaf cohomology of E .

⁵In previous case, $E = \Omega_X^{p,0}$

Part 2. Complex Differential Geometry

5. CONNECTIONS AND ITS CURVATURE

5.1. Connections on complex vector bundle. Let X be a complex manifold and $\pi: E \rightarrow X$ be a complex vector bundle.

5.1.1. *Basic definitions.*

Definition 5.1.1 (connection). A connection on E is a \mathbb{C} -linear operator

$$\nabla: C^\infty(X, E) \rightarrow C^\infty(X, \Omega_{X, \mathbb{C}}^1 \otimes E)$$

satisfying the Leibniz rule

$$\nabla(fs) = df \otimes s + f\nabla s$$

for $f \in C^\infty(X)$ and $s \in C^\infty(X, E)$.

Remark 5.1.1 (connection form). Let $\{e_\alpha\}$ be a local frame of E . Then any section s of E can be written as $s = s^\alpha e_\alpha$, and

$$\begin{aligned} \nabla(s^\alpha e_\alpha) &= ds^\alpha e_\alpha + s^\alpha \nabla s_\alpha \\ &= ds^\alpha e_\alpha + s^\alpha \omega_\alpha^\beta e_\beta \end{aligned}$$

where ω_α^β are 1-forms, which is called connection 1-form. In terms of Christoffel symbol, one has

$$\omega_\alpha^\beta = \Gamma_{i\alpha}^\beta dz^i + \Gamma_{i\alpha}^\beta d\bar{z}^i$$

5.1.2. *Curvature form.* Now we're going to extend connection to something called exterior derivative defined on sections of vector bundle valued k -forms as follows

$$\begin{aligned} d^\nabla: C^\infty(X, \Omega_{X, \mathbb{C}}^k \otimes E) &\rightarrow C^\infty(X, \Omega_{X, \mathbb{C}}^{k+1} \otimes E) \\ \omega \otimes s &\mapsto d\omega \otimes s + (-1)^k \omega \wedge \nabla s \end{aligned}$$

Definition 5.1.2 (curvature form). Let E be a complex vector bundle over a complex manifold X equipped with connection ∇ . There exists a section $\Theta \in C^\infty(X, \Omega_{X, \mathbb{C}}^2 \otimes \text{End } E)$, called curvature form, such that

$$(d^\nabla)^2 s = \Theta \wedge s$$

for all $s \in C^\infty(X, \Omega_{X, \mathbb{C}}^k \otimes E)$.

Remark 5.1.2 (local form). Let $\{e_\alpha\}$ be a local frame of E . The curvature form Θ can be written as

$$\Theta = \Theta_{i\bar{j}\alpha}^\beta dz^i \wedge d\bar{z}^j \otimes e^\alpha \otimes e_\beta$$

where $\Theta_{i\bar{j}\alpha}^\beta$ can also be expressed in terms of Christoffel symbols just like what we have seen in [Liu23].

5.2. Chern connection. In this section, we will introduce additional structures to enhance the complexity of a vector bundle E over a complex manifold X . These structures include Hermitian metrics and complex structures. We will explore connections that align harmoniously with these structures, similar to our approach in Riemannian geometry. By doing so, we will obtain the Chern connection, which runs parallel to the Levi-Civita connection.

5.2.1. *Compatibility with Hermitian metric.*

Definition 5.2.1 (Hermitian metric). Let E be a complex vector bundle. A Hermitian metric h on E is a smooth section of $E^* \otimes \overline{E}^*$.

Remark 5.2.1 (local form). Let $\{e_\alpha\}$ be a local frame of E . Then a Hermitian metric is determined by a positive definite Hermitian matrix $(h_{\alpha\bar{\beta}})$, that is

$$h = h_{\alpha\bar{\beta}} e^\alpha \otimes \bar{e}^\beta$$

where $h_{\alpha\bar{\beta}} = h(e_\alpha, \bar{e}_\beta)$.

Definition 5.2.2 (Hermitian vector bundle). A complex vector bundle E together with a Hermitian metric h is called a Hermitian vector bundle (E, h) .

Remark 5.2.2 (metric weight). Let L be a Hermitian line bundle. A Hermitian metric h is locally given by $e^{-2\varphi}$, where φ is a smooth function, which is called metric weight. Suppose $\{g_{\alpha\beta}\}$ is transition function of L with respect to open covering $\{U_\alpha\}$. Then h is given by a collection $\{h_\alpha \in C^\infty(U_\alpha)\}$ such that $h_\alpha = |g_{\alpha\beta}|^{-2} h_\beta$. In other words, a Hermitian metric is a collection of metric weights $\{\varphi_\alpha \in C^\infty(U_\alpha)\}$ such that

$$\varphi_\alpha = \varphi_\beta + \log |g_{\alpha\beta}|$$

Definition 5.2.3 (sesquilinear map). For a Hermitian vector bundle (E, h) over complex manifold X , there is a sesquilinear map

$$\begin{aligned} C^\infty(X, \Omega_{X,\mathbb{C}}^p \otimes E) \times C^\infty(X, \Omega_{X,\mathbb{C}}^q \otimes E) &\rightarrow C^\infty(X, \Omega_{X,\mathbb{C}}^{p+q}) \\ (s, t) &\mapsto \{s, t\} \end{aligned}$$

locally given by

$$\{s^\alpha e_\alpha, t^\beta e_\beta\} = h_{\alpha\bar{\beta}} s^\alpha \wedge \bar{t}^\beta$$

Definition 5.2.4 (metric connection). A connection ∇ on a Hermitian vector bundle (E, h) is called a metric connection, if

$$d\langle s, t \rangle = \{\nabla s, t\} + \{s, \nabla t\}$$

where s, t are sections of E .

Remark 5.2.3 (local form). If $\{e_\alpha\}$ is a local frame of E , then

$$\begin{aligned} dh_{\alpha\bar{\beta}} &= d\langle e_\alpha, \bar{e}_\beta \rangle \\ &= \{\nabla e_\alpha, \bar{e}_\beta\} + \{e_\alpha, \nabla \bar{e}_\beta\} \\ &= \omega_\alpha^\gamma h_{\gamma\bar{\beta}} + \overline{\omega_\beta^\gamma} h_{\alpha\bar{\gamma}} \end{aligned}$$

So in matrix notation, we have

$$dh = \omega h + h \bar{\omega}^T$$

In particular, if we take $\{e_\alpha\}$ to be orthogonal local frame of E with respect to h , we will find $\omega + \bar{\omega}^T = 0$, that is ω is skew-Hermitian matrix.

Proposition 5.2.1. Let (E, h) be a Hermitian vector bundle over a complex manifold X . A connection ∇ is a metric connection if and only if

$$d\{s, t\} = \{\nabla s, t\} + (-1)^p \{s, \nabla t\}$$

where $s \in C^\infty(X, \Omega_{X, \mathbb{C}}^p \otimes E)$ and $t \in C^\infty(X, \Omega_{X, \mathbb{C}}^q \otimes E)$.

Proposition 5.2.2. Let (E, h) be a Hermitian vector bundle equipped with connection ∇^E . Then ∇^E is a metric connection if and only if $\nabla^{E^* \otimes \bar{E}^*} h = 0$.

Proof. Direct computation shows

$$\begin{aligned} \nabla^{E^* \otimes \bar{E}^*} (h_{\alpha\bar{\beta}} e^\alpha \otimes \bar{e}^\beta) &= dh_{\alpha\bar{\beta}} \otimes e^\alpha \otimes \bar{e}^\beta + h_{\alpha\bar{\beta}} \nabla^{E^*} e^\alpha \otimes \bar{e}^\beta + h_{\alpha\bar{\beta}} e^\alpha \otimes \nabla^{\bar{E}^*} \bar{e}^\beta \\ &= dh_{\alpha\bar{\beta}} \otimes e^\alpha \otimes \bar{e}^\beta - h_{\alpha\bar{\beta}} \omega_\gamma^\alpha e^\gamma \otimes \bar{e}^\beta - h_{\alpha\bar{\beta}} \bar{\omega}_\gamma^\beta e^\alpha \otimes \bar{e}^\gamma \\ &= (dh_{\alpha\bar{\beta}} - \omega_\alpha^\gamma h_{\gamma\bar{\beta}} - \bar{\omega}_\beta^\gamma h_{\alpha\bar{\gamma}}) e^\alpha \otimes \bar{e}^\beta \end{aligned}$$

This shows desired result. \square

5.2.2. *Compatibility with complex structure.* For a complex manifold X , we have decomposition

$$\Omega_{X, \mathbb{C}}^1 = \Omega_X^{1,0} \oplus \Omega_X^{0,1}$$

Let $E \rightarrow X$ be a complex vector bundle with connection ∇ . Then we can decompose $\nabla = \nabla^{1,0} + \nabla^{0,1}$ by composing the projection as follows

$$\begin{array}{ccc} & & C^\infty(X, \Omega_X^{1,0} \otimes E) \\ & \nearrow & \\ C^\infty(X, E) & \xrightarrow{\nabla} & C^\infty(X, \Omega_{X, \mathbb{C}}^1 \otimes E) \\ & \searrow & \\ & & C^\infty(X, \Omega_X^{0,1} \otimes E) \end{array}$$

If we write $\nabla = d + \omega$ locally, then

$$\begin{aligned} \nabla^{1,0} &= \partial + \omega^{1,0} \\ \nabla^{0,1} &= \bar{\partial} + \omega^{0,1} \end{aligned}$$

Definition 5.2.5 (complex connection). A connection ∇ on a holomorphic vector bundle E over a complex manifold X is said to be compatible with complex structure if $\nabla^{0,1} = \bar{\partial}_E$.

Remark 5.2.4 (local form). Let $\{e_\alpha\}$ be a holomorphic local form of E , and denote

$$\nabla e_\alpha = (\Gamma_{i\alpha}^\beta dz^i + \Gamma_{\bar{i}\alpha}^\beta d\bar{z}^i) e_\beta$$

that is

$$\nabla^{0,1} e_\alpha = \Gamma_{\bar{i}\alpha}^\beta e_\beta d\bar{z}^i$$

But since $\{e_\alpha\}$ is holomorphic, that is $\bar{\partial}_E e_\alpha = 0$, which implies ∇ is complex if and only if $\Gamma_{\bar{i}\alpha}^\beta = 0$.

5.2.3. *Chern connection.*

Theorem 5.2.1 (Chern connection). Let X be a complex manifold, (E, h) a Hermitian holomorphic vector bundle. Then there exists a unique metric connection called Chern connection such that it's compatible with complex structure.

Proof. If metric connection ∇ is compatible with complex structure, then the following three equations are equivalent

$$\begin{aligned} dh &= \omega h + h \bar{\omega}^t \\ \partial h &= \omega h \\ \bar{\partial} h &= h \bar{\omega}^t \end{aligned}$$

since ω is a $(1, 0)$ -valued matrix. This shows Chern connection is uniquely determined by $\omega = (\partial h)h^{-1}$. \square

Remark 5.2.5 (local form). Chern connection is locally determined by

$$\frac{\partial h_{\alpha\bar{\beta}}}{\partial z^i} = \Gamma_{i\alpha}^\gamma h_{\gamma\bar{\beta}}$$

Definition 5.2.6 (Chern curvature). Let X be a complex manifold and (E, h) be a Hermitian holomorphic vector bundle. The Chern curvature Θ_h of (E, h) is defined as the curvature of Chern connection with respect to h .

Corollary 5.2.1. Let X be a complex manifold and (E, h) a Hermitian holomorphic vector bundle equipped with Chern connection ∇ locally given by ω . Then

- (1) $\partial\omega = \omega \wedge \omega$.
- (2) $\Theta_h = \bar{\partial}\omega$.
- (3) $\bar{\partial}\Theta_h = 0$.

Proof. For (1). Since $\omega = (\partial h)h^{-1}$, then directly computation shows

$$\begin{aligned} \partial\omega &= -\partial h \wedge \partial(h^{-1}) \\ &= -\partial h \wedge (-h^{-1}\partial h h^{-1}) \\ &= (\partial h)h^{-1} \wedge (\partial h)h^{-1} \\ &= \omega \wedge \omega \end{aligned}$$

For (2). Θ_h locally looks like

$$\Theta_h = d\omega - \omega \wedge \omega = d\omega - \partial\omega = \bar{\partial}\omega$$

For (3). It's clear from (2). \square

Remark 5.2.6 (local form). The Chern curvature can be expressed in terms of Christoffel symbol as follows

$$\Theta_h = \Theta_{i\bar{j}\alpha}^\gamma dz^i \wedge d\bar{z}^j \otimes e^\alpha \otimes e_\gamma$$

where $\Theta_{i\bar{j}\alpha}^\gamma = -\frac{\partial\Gamma_{i\alpha}^\gamma}{\partial\bar{z}^j}$. In other type one has

$$\begin{aligned} \Theta_{i\bar{j}\alpha\bar{\beta}} &= h_{\gamma\bar{\beta}} \Theta_{i\bar{j}\alpha}^\gamma \\ &= -h_{\gamma\bar{\beta}} \partial_{\bar{j}} (h^{\gamma\bar{\delta}} \frac{\partial h_{\alpha\bar{\delta}}}{\partial z^i}) \\ &= -\frac{\partial^2 h_{\alpha\bar{\beta}}}{\partial z^i \partial \bar{z}^j} + h^{\gamma\bar{\delta}} \frac{\partial h_{\alpha\bar{\delta}}}{\partial z^i} \frac{\partial h_{\gamma\bar{\beta}}}{\partial \bar{z}^j} \end{aligned}$$

5.2.4. Useful formulas of Chern connection. Let X be a complex manifold and (E, h) be a Hermitian holomorphic vector bundle over X . Let ∇ be Chern connection determined by Christoffel symbol $\Gamma_{i\alpha}^\beta$ on (E, h) with curvature Θ_h . Suppose $\{z^i\}$ is local coordinate of X , $\{e_\alpha\}$ is the local frame of E , and $\{e^\alpha\}$ and $\{\bar{e}_\alpha\}$ denote local frames of E^* and \bar{E} respectively.

Proposition 5.2.3.

$$\Gamma_{i\alpha}^\beta = h^{\beta\bar{\gamma}} \frac{\partial h_{\alpha\bar{\gamma}}}{\partial z^i}$$

Proof. See Remark 5.2.5. \square

Proposition 5.2.4.

$$\begin{aligned} \Theta_{i\bar{j}\alpha}^\gamma &= -\frac{\partial\Gamma_{i\alpha}^\gamma}{\partial\bar{z}^j} \\ \Theta_{i\bar{j}\alpha\bar{\beta}} &= -\frac{\partial^2 h_{\alpha\bar{\beta}}}{\partial z^i \partial \bar{z}^j} + h^{\gamma\bar{\delta}} \frac{\partial h_{\alpha\bar{\delta}}}{\partial z^i} \frac{\partial h_{\gamma\bar{\beta}}}{\partial \bar{z}^j} \end{aligned}$$

Proof. See Remark 5.2.6. \square

Proposition 5.2.5.

$$\begin{aligned} \nabla_{\frac{\partial}{\partial z^i}} e_\alpha &= \Gamma_{i\alpha}^\beta e_\beta, & \nabla_{\frac{\partial}{\partial \bar{z}^i}} e_\alpha &= 0, & \nabla_{\frac{\partial}{\partial z^i}} \bar{e}_\alpha &= 0, & \nabla_{\frac{\partial}{\partial \bar{z}^i}} \bar{e}_\alpha &= \Gamma_{i\alpha}^{\bar{\beta}} \bar{e}_\beta \\ \nabla_{\frac{\partial}{\partial z^i}} e^\alpha &= -\Gamma_{i\beta}^\alpha e^\beta, & \nabla_{\frac{\partial}{\partial \bar{z}^i}} e^\alpha &= 0, & \nabla_{\frac{\partial}{\partial z^i}} \bar{e}^\alpha &= 0, & \nabla_{\frac{\partial}{\partial \bar{z}^i}} \bar{e}^\alpha &= -\Gamma_{i\beta}^{\bar{\alpha}} \bar{e}^\beta \end{aligned}$$

Proof. It suffices to show the first two equalities, and others can be obtained from taking conjugates and dualities. The first one holds from definition of Christoffel symbol, and $\Gamma_{i\alpha}^\beta = 0$ holds from the Remark 5.2.4. \square

Corollary 5.2.2.

(1) For $s \in C^\infty(X, E)$, locally written as $s = s^\alpha e_\alpha$, one has

$$\begin{aligned}\nabla_{\frac{\partial}{\partial z^i}} s &= \left(\frac{\partial s^\beta}{\partial z^i} + s^\alpha \Gamma_{i\alpha}^\beta \right) e_\beta \\ \nabla_{\frac{\partial}{\partial \bar{z}^i}} s &= \frac{\partial s^\beta}{\partial \bar{z}^i} e_\beta\end{aligned}$$

(2) For $s \in C^\infty(X, \bar{E})$, locally written as $s = s^\alpha \bar{e}_\alpha$, one has

$$\begin{aligned}\nabla_{\frac{\partial}{\partial z^i}} s &= \frac{\partial s^\beta}{\partial z^i} \bar{e}_\beta \\ \nabla_{\frac{\partial}{\partial \bar{z}^i}} s &= \left(\frac{\partial s^\beta}{\partial \bar{z}^i} + s^\alpha \bar{\Gamma}_{i\alpha}^\beta \right) \bar{e}_\beta\end{aligned}$$

(3) For $s \in C^\infty(X, E^*)$, locally written as $s = s_\alpha e^\alpha$, one has

$$\begin{aligned}\nabla_{\frac{\partial}{\partial z^i}} s &= \left(\frac{\partial s_\beta}{\partial z^i} - s_\alpha \Gamma_{i\beta}^\alpha \right) e^\beta \\ \nabla_{\frac{\partial}{\partial \bar{z}^i}} s &= \frac{\partial s_\beta}{\partial \bar{z}^i} e^\beta\end{aligned}$$

(4) For $s \in C^\infty(X, \bar{E}^*)$, locally written as $s = s_\alpha \bar{e}^\alpha$, one has

$$\begin{aligned}\nabla_{\frac{\partial}{\partial z^i}} s &= \frac{\partial s_\beta}{\partial z^i} \bar{e}^\beta \\ \nabla_{\frac{\partial}{\partial \bar{z}^i}} s &= \left(\frac{\partial s_\beta}{\partial \bar{z}^i} - s_\alpha \bar{\Gamma}_{i\beta}^\alpha \right) \bar{e}^\beta\end{aligned}$$

Proposition 5.2.6 (Ricci identity). For $s \in C^\infty(X, E)$, locally written as $s = s^\alpha e_\alpha$, one has

$$\nabla_{\frac{\partial}{\partial z^i}} \nabla_{\frac{\partial}{\partial \bar{z}^j}} s^\beta - \nabla_{\frac{\partial}{\partial \bar{z}^j}} \nabla_{\frac{\partial}{\partial z^i}} s^\beta = \Theta_{ij\alpha}^\beta s^\alpha$$

Proof. Direct computation shows

$$\begin{aligned}\nabla_{\frac{\partial}{\partial z^i}} \nabla_{\frac{\partial}{\partial \bar{z}^j}} s &= \nabla_{\frac{\partial}{\partial z^i}} \left(\frac{\partial s^\alpha}{\partial \bar{z}^j} e_\alpha \right) \\ &= \frac{\partial^2 s^\beta}{\partial z^i \partial \bar{z}^j} e_\beta + \Gamma_{i\alpha}^\beta \frac{\partial s^\alpha}{\partial \bar{z}^j} e_\beta\end{aligned}$$

that is

$$\nabla_{\frac{\partial}{\partial z^i}} \nabla_{\frac{\partial}{\partial \bar{z}^j}} s^\beta = \frac{\partial^2 s^\beta}{\partial z^i \partial \bar{z}^j} + \Gamma_{i\alpha}^\beta \frac{\partial s^\alpha}{\partial \bar{z}^j}$$

Direct computation also shows

$$\nabla_{\frac{\partial}{\partial \bar{z}^j}} \nabla_{\frac{\partial}{\partial z^i}} s^\beta = \frac{\partial^2 s^\beta}{\partial \bar{z}^j \partial z^i} + \frac{\partial s^\alpha}{\partial \bar{z}^j} \Gamma_{i\alpha}^\beta + s^\alpha \frac{\partial \Gamma_{i\alpha}^\beta}{\partial \bar{z}^j}$$

Thus

$$\begin{aligned}\nabla_{\frac{\partial}{\partial z^i}} \nabla_{\frac{\partial}{\partial \bar{z}^j}} s^\beta - \nabla_{\frac{\partial}{\partial \bar{z}^j}} \nabla_{\frac{\partial}{\partial z^i}} s^\beta &= -s^\alpha \frac{\partial \Gamma_{i\alpha}^\beta}{\partial \bar{z}^j} \\ &= \Theta_{ij\alpha}^\beta s^\alpha\end{aligned}$$

□

5.3. First Chern class.

5.3.1. *First Chern class of complex line bundle.* Let $\pi: X \rightarrow L$ be a complex line bundle with connection ∇ over a complex manifold X . Then curvature Θ is a global section of $\Omega_{X,\mathbb{C}}^2$ since $\text{End } L$ is trivial bundle. Furthermore, Θ locally looks like $d\omega$ since for line bundle $\omega \wedge \omega = 0$. An immediate consequence is $d\Theta = 0$, that is Θ gives a cohomology class

$$[\Theta] \in H^2(X, \mathbb{C})$$

Definition 5.3.1 (first Chern class of line bundle). Let L be a complex line bundle over complex manifold X equipped with connection ∇ . The first Chern class of L is defined as

$$c_1(L) := \left[\frac{\sqrt{-1}}{2\pi} \Theta \right] \in H^2(X, \mathbb{C})$$

where Θ is the curvature of ∇ .

Proposition 5.3.1 (topological invariance). $c_1(L) \in H^2(X, \mathbb{C})$ is independent of the choice of connection.

Proof. Let $\tilde{\nabla}$ be another connection which is locally given by $\tilde{\omega}$. Then for section s of $\Omega_{X,\mathbb{C}}^k \otimes L$, one has

$$\begin{aligned} (\nabla - \tilde{\nabla})s &= (ds + \omega \wedge s) - (ds + \tilde{\omega} \wedge s) \\ &= (\omega - \tilde{\omega}) \wedge s \end{aligned}$$

Note that $\omega - \tilde{\omega}$ is a global section of $\Omega_{X,\mathbb{C}}^1$, so $\Theta - \tilde{\Theta}$ is exact. □

Proposition 5.3.2. Let $\pi: L \rightarrow X$ be a complex line bundle over a complex manifold X . Then $c_1(L) \in H^2(X, \mathbb{R})$.

Proof. Equip L with a Hermitian metric h , then for a metric connection ∇ , locally we have

$$\bar{\omega} = -\omega$$

Thus

$$\overline{\frac{\sqrt{-1}}{2\pi} \Theta} = -\frac{\sqrt{-1}}{2\pi} \bar{\Theta} = -\frac{\sqrt{-1}}{2\pi} d\bar{\omega} = \frac{\sqrt{-1}}{2\pi} d\omega = \frac{\sqrt{-1}}{2\pi} \Theta$$

□

Remark 5.3.1. Here are two facts here we don't prove:

- (1) $c_1(L) \in H^2(X, \mathbb{Z})$.
- (2) L is determined by $c_1(L)$.

Definition 5.3.2 (first Chern class of vector bundle). Let E be a complex vector bundle over complex manifold X . The first Chern class of E is defined to be the first Chern class of $\det E$.

5.3.2. *First Chern class of Hermitian holomorphic line bundle.* Let X be a complex manifold, (L, h) a Hermitian holomorphic line bundle, and ∇ is the Chern connection of (L, h) with Chern curvature Θ_h . Then by Proposition 5.3.2 and Corollary 5.2.1, we have

$$\left[\frac{\sqrt{-1}}{2\pi}\Theta_h\right] \in H^2(X, \mathbb{R}) \cap H^{1,1}(X)$$

Remark 5.3.2 (local form). Suppose Hermitian metric h is given by metric weight $\{\varphi_\alpha\}$, that is locally $h = e^{-2\varphi_\alpha}$. Then direct computation shows first Chern class is locally given by

$$\frac{\sqrt{-1}}{2\pi}\Theta_h = \frac{\sqrt{-1}}{\pi}\partial\bar{\partial}\varphi_\alpha$$

Proposition 5.3.3. $\left[\frac{\sqrt{-1}}{2\pi}\Theta_h\right] \in H^{1,1}(X)$ is independent of h .

Proof. Note that any two metric on a line bundle differ a smooth function which is positive everywhere, so if h and h' are two different metrics, we can write $\|e(z)\|_{h'} = e^f\|e(z)\|_h$ for some globally defined smooth function f . So by Remark 5.3.2, we have the difference of first Chern classes coming from different metrics is $\frac{\sqrt{-1}}{\pi}\partial\bar{\partial}f$, and it's trivial in $H^{1,1}(X)$ since f is globally defined. \square

5.4. **Lefschetz (1, 1)-theorem.** Now we know that given a Hermitian holomorphic line bundle (L, h) , its Chern curvature we will get a real (1, 1)-form. So we may wonder the converse of this statement. Is there any real (1, 1)-form comes from such a Hermitian holomorphic line bundle? That's main theorem for this section.

Theorem 5.4.1 (Lefschetz (1, 1)-theorem). Let X be a complex manifold and $[\omega] \in H^2(X, \mathbb{R}) \cap H^{1,1}(X)$. If

$$[\omega] \in \text{im}\{H^2(X, \mathbb{Z}) \rightarrow H^2(X, \mathbb{R})\},$$

then there exists a Hermitian holomorphic line bundle (L, h) such that

$$\frac{\sqrt{-1}}{2\pi}\Theta_h = \omega$$

Before proving this theorem, let's explain these notations. Here $H^2(X, \mathbb{Z})$ and $H^2(X, \mathbb{R})$ are sheaf cohomology of constant sheaves $\underline{\mathbb{Z}}$ and $\underline{\mathbb{R}}$. By de Rham theorem⁶, there is no difference between sheaf cohomology of $\underline{\mathbb{R}}$ and de Rham cohomology, but it's meaningless to consider de Rham cohomology with \mathbb{Z} -coefficient.

Here we use the isomorphisms

$$\begin{aligned} H^2(X, \mathbb{Z}) &\cong \check{H}^2(X, \underline{\mathbb{Z}}) \\ H^2(X, \mathbb{R}) &\cong \check{H}^2(X, \underline{\mathbb{R}}) \end{aligned}$$

⁶See appendix 20.

and consider the map in terms of Čech cohomology

$$\check{H}^2(X, \mathbb{Z}) \rightarrow \check{H}^2(X, \mathbb{R}).$$

Above isomorphism is called comparison theorem, and can be proved by technique of spectral sequences in general. Here we give an explicit construction in dimension two, that is, construct a Čech 2-cocycle from a closed 2-form.

In sketch, the philosophy of this construction is that we can descend the degree of differential forms, but the price we pay is to consider functions defined on intersections of many open subsets.

Proof of comparison theorem in dimension two. Let X be a smooth manifold and $Z^1(X) \subset \Omega_{X, \mathbb{R}}^1$ be the sheaf of closed 1-form. Then we have the following sequence of sheaves

$$0 \rightarrow \underline{\mathbb{R}} \rightarrow C^\infty(X) \xrightarrow{d} Z^1 \rightarrow 0.$$

By Poincaré lemma it's an exact sequence. Similarly, there is also an exact sequence

$$0 \rightarrow Z^1 \rightarrow \Omega_{X, \mathbb{R}}^1 \xrightarrow{d} Z^2 \rightarrow 0,$$

where Z^2 is the sheaf of closed 2-forms. By the definition of de Rham cohomology, we have

$$H^2(X, \mathbb{R}) = \frac{C^\infty(X, Z^2)}{dC^\infty(X, \Omega_{X, \mathbb{R}}^1)}$$

In order to avoid taking limit in Čech cohomology, we choose a good enough open covering⁷ $\mathfrak{U} = \{U_\alpha\}_{\alpha \in I}$ such that

(1)

$$d: C^\infty(U_\alpha, \Omega_{U_\alpha, \mathbb{R}}^1) \rightarrow C^\infty(U_\alpha, Z^2)$$

is surjective for any $\alpha \in I$.

(2)

$$d: C^\infty(U_\alpha \cap U_\beta) \rightarrow C^\infty(U_\alpha \cap U_\beta, Z^1)$$

is surjective for any $\alpha, \beta \in I$.

Let ω be a closed 2-form. For any $\alpha \in I$, we choose $A_\alpha \in C^\infty(U_\alpha, \Omega_{U_\alpha, \mathbb{R}}^1)$ such that

$$\omega|_{U_\alpha} = dA_\alpha.$$

Then

$$\prod_{\alpha, \beta} (A_\alpha - A_\beta)$$

is a Čech 1-cocycle in $C^1(\mathfrak{U}, Z^1)$ since $d(A_\alpha - A_\beta)|_{U_\alpha \cap U_\beta} = \omega - \omega = 0$. For any $\alpha, \beta \in I$, we choose $f_{\alpha\beta} \in C^\infty(U_\alpha \cap U_\beta)$ such that

$$(A_\alpha - A_\beta)_{\alpha\beta} = df_{\alpha\beta}.$$

⁷In fact, it's called Leray covering, and Leray's theorem about Čech cohomology says that the Čech cohomology with respect to Leray covering is exactly the Čech cohomology.

Note that

$$f_{\beta\gamma} - f_{\alpha\gamma} + f_{\alpha\beta}|_{U_\alpha \cap U_\beta \cap U_\gamma}$$

is d-closed by the same reason, and thus it's locally constant. Then

$$\tilde{\omega} = \prod_{\alpha, \beta, \gamma} (f_{\beta\gamma} - f_{\alpha\gamma} + f_{\alpha\beta})$$

is a Čech 2-cocycle in $C^2(\mathfrak{U}, \mathbb{R})$. Then by Leray's theorem, we obtain a Čech 2-cocycle $\tilde{\omega} \in \check{H}^2(X, \mathbb{R})$ from a closed 2-form $\omega \in H^2(X, \mathbb{R})$. \square

Now let's prove Lefschetz (1, 1)-theorem.

Proof of theorem 5.4.1. Let $\mathfrak{U} = \{U_\alpha\}_{\alpha \in I}$ be an open covering consisting of open polydisk such that for all $\alpha, \beta \in I$, the intersection $U_\alpha \cap U_\beta$ is simply-connected.

For a d-closed real (1, 1)-form ω , after a refinement if necessary, Lemma 0.2.1 implies that there exist smooth functions $\varphi_\alpha: U_\alpha \rightarrow \mathbb{R}$ such that

$$\omega|_{U_\alpha} = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \varphi_\alpha.$$

Then on any two intersection $U_\alpha \cap U_\beta$, one has $\partial \bar{\partial}(\varphi_\alpha - \varphi_\beta) = 0$. Again after a refinement if necessary, Lemma 0.2.2 implies that there exist holomorphic functions $f_{\alpha\beta}$ such that

$$(\varphi_\alpha - \varphi_\beta)|_{U_\alpha \cap U_\beta} = 2\text{Re}(f_{\alpha\beta}) = f_{\alpha\beta} + \overline{f_{\alpha\beta}}.$$

For $\prod f_{\alpha\beta} \in C^1(\mathfrak{U}, \mathcal{O}_X)$, one has

$$(\delta f)_{\alpha\beta\gamma} = (f_{\beta\gamma} - f_{\alpha\gamma} + f_{\alpha\beta})|_{U_\alpha \cap U_\beta \cap U_\gamma}$$

Since $2\text{Re}(f_{\beta\gamma} - f_{\alpha\gamma} + f_{\alpha\beta})_{\alpha\beta\gamma} = 0$, it must be a locally constant pure imaginary number, that is, it lies in $2\pi\sqrt{-1}\mathbb{R}(U_\alpha \cap U_\beta \cap U_\gamma)$.

For real 1-form

$$A_\alpha = \frac{\sqrt{-1}}{4\pi} (\bar{\partial} \varphi_\alpha - \partial \varphi_\alpha),$$

a direct computation shows that $\omega|_{U_\alpha} = dA_\alpha$, and that's why we define A_α in this form.

Recall what we have done in the proof of comparison theorem: If we want to find Čech cocycle which corresponding to ω , we need to consider $A_\alpha - A_\beta$ on the intersection $U_\alpha \cap U_\beta$. A direct computation shows that

$$\begin{aligned} \partial(\varphi_\beta - \varphi_\alpha) &= \partial(f_{\alpha\beta} + \overline{f_{\alpha\beta}}) \\ &= \partial f_{\alpha\beta} \\ &= df_{\alpha\beta} \\ \bar{\partial}(\varphi_\beta - \varphi_\alpha) &= d\overline{f_{\alpha\beta}}. \end{aligned}$$

Thus

$$(A_\beta - A_\alpha)_{\alpha\beta} = \frac{\sqrt{-1}}{4\pi} d(\overline{f_{\alpha\beta}} - f_{\alpha\beta}) = \frac{1}{2\pi} d(\text{Im}(f_{\alpha\beta})).$$

Then the Čech 2-cocycle $\check{\omega}$ corresponding to ω is

$$\begin{aligned}\check{\omega} &= \prod \left(\frac{1}{2\pi} \operatorname{Im}(f_{\beta\gamma} - f_{\alpha\gamma} + f_{\alpha\beta}) \right)_{\alpha\beta\gamma} \\ &= \prod \left(\frac{1}{2\pi\sqrt{-1}} (f_{\beta\gamma} - f_{\alpha\gamma} + f_{\alpha\beta}) \right)_{\alpha\beta\gamma}.\end{aligned}$$

By hypothesis one has $[\check{\omega}]$ is an image of $[\prod n_{\alpha\beta\gamma}] \in \check{H}^2(X, \mathbb{Z})$. However, it doesn't mean that $f_{\alpha\beta}$ are exactly integers, but not too bad, we just need some correction terms, that is

$$\prod \left(\frac{1}{2\pi\sqrt{-1}} (f_{\beta\gamma} - f_{\alpha\gamma} + f_{\alpha\beta}) \right)_{\alpha\beta\gamma} = \prod n_{\alpha\beta\gamma} + \delta \left(\prod c_{\alpha\beta} \right)$$

where $\prod (c_{\alpha\beta}) \in C^1(\mathfrak{U}, \mathbb{R})$ is 1-cochain. If we define $f'_{\alpha\beta} = f_{\alpha\beta} - 2\pi\sqrt{-1}c_{\alpha\beta}$, then

$$(f'_{\beta\gamma} - f'_{\alpha\gamma} + f'_{\alpha\beta})_{\alpha\beta\gamma} = 2\pi\sqrt{-1}n_{\alpha\beta\gamma} \in 2\pi\sqrt{-1}\mathbb{Z}(U_\alpha \cap U_\beta \cap U_\gamma).$$

Now consider the holomorphic function from $U_\alpha \cap U_\beta$ to \mathbb{C}^* defined by $g_{\alpha\beta} = \exp(-f'_{\alpha\beta})$. A direct computation shows that it satisfies the cocycle condition

$$g_{\beta\gamma}g_{\alpha\gamma}^{-1}g_{\alpha\beta} = 1,$$

since $e^{2\pi\sqrt{-1}} = 1$. Then $\{g_{\alpha\beta}\}$ is a collection of transition functions, and gives a holomorphic line bundle L .

Now it suffices to construct a Hermitian metric on L , and calculate its curvature to complete the proof. Note that

$$(\varphi_\alpha - \varphi_\beta)_{U_\alpha \cap U_\beta} = 2\operatorname{Re}(f_{\alpha\beta}) = 2\operatorname{Re}(f_{\alpha\beta})' = -\log |g_{\alpha\beta}|^2.$$

Consider the Hermitian metric h , which is locally given by

$$h_\alpha = \exp(-\varphi_\alpha)$$

on U_α . It's well-defined since $h_\beta = |g_{\alpha\beta}|^2 h_\alpha = g_{\alpha\beta}^T h_\alpha \overline{g_{\alpha\beta}}$. Moreover,

$$\frac{\sqrt{-1}}{2\pi} \Theta_h = \frac{\sqrt{-1}}{2\pi} \partial\bar{\partial}\varphi_\alpha = \omega.$$

This completes the proof. \square

Remark 5.4.1. Consider the exponential sequence

$$0 \rightarrow \mathbb{Z} \xrightarrow{2\pi\sqrt{-1}} \mathcal{O}_X \xrightarrow{\exp} \mathcal{O}_X^* \rightarrow 0$$

and the induced long exact sequence

$$\cdots \rightarrow H^1(X, \mathcal{O}_X^*) \xrightarrow{\delta} H^2(X, \mathbb{Z}) \rightarrow H^2(X, \mathcal{O}_X) \rightarrow \cdots$$

Let L be a holomorphic line bundle determined by its transition functions $\{g_{\alpha\beta}\}$. The proof for Lefschetz (1, 1)-theorem shows that δ maps $\{g_{\alpha\beta}\}$ to $-c_1(L)$.

6. HERMITIAN GEOMETRY

6.1. Hermitian manifold and Riemannian manifold. A Hermitian manifold is a complex manifold X together with a Hermitian metric h on holomorphic tangent bundle TX . One way to construct a Hermitian metric on TX is to consider special Riemannian metric on the underlying real manifold.

Suppose $\{z^i = x^i + \sqrt{-1}x^I\}$ is a local coordinate of X , where $1 \leq i \leq n$ and $n+1 \leq I = i+n \leq 2n$. Then $\{x^i, x^I\}$ gives a local coordinate of underlying real manifold of X , and there is a natural almost complex structure J on $T_{\mathbb{R}}X$ which is given by

$$\begin{aligned} J\left(\frac{\partial}{\partial x^i}\right) &= \frac{\partial}{\partial x^I} \\ J\left(\frac{\partial}{\partial x^I}\right) &= -\frac{\partial}{\partial x^i} \end{aligned}$$

Definition 6.1.1 (compatibility). A Riemannian metric g on $T_{\mathbb{R}}X$ is called compatible with almost complex structure, if

$$g(V, W) = g(JV, JW)$$

for all $V, W \in C^\infty(X, T_{\mathbb{R}}X)$.

Proposition 6.1.1. Let g be a Riemannian metric on $T_{\mathbb{R}}X$ which is compatible with J and locally given by

$$g = g_{ij}dx^i \otimes dx^j + g_{iJ}dx^i \otimes dx^J + g_{IJ}dx^I \otimes dx^J + g_{IJ}dx^I \otimes dx^J$$

Then

$$\begin{aligned} g_{ij} &= g_{IJ} \\ g_{iJ} &= g_{Ji} = -g_{jI} = -g_{Ij} \end{aligned}$$

Proof. Direct computation. □

Notation 6.1.1.

$$\begin{pmatrix} g^{il} & g^{iL} \\ g^{Ii} & g^{IL} \end{pmatrix} \begin{pmatrix} g_{lj} & g_{lJ} \\ g_{Lj} & g_{LJ} \end{pmatrix} = \mathbf{I}_{2n}$$

In other words,

$$\begin{aligned} g^{il}g_{lj} + g^{iL}g_{Lj} &= \delta_j^i \\ g^{il}g_{lJ} + g^{iL}g_{LJ} &= 0 \end{aligned}$$

Let g be a Riemannian metric on underlying real manifold which is compatible with J . Then its \mathbb{C} -linear extension $g_{\mathbb{C}}$ gives a matrix

$$G = \begin{pmatrix} (g_{\mathbb{C}}(\frac{\partial}{\partial z^i}, \frac{\partial}{\partial z^j}))_{n \times n} & (g_{\mathbb{C}}(\frac{\partial}{\partial z^i}, \frac{\partial}{\partial \bar{z}^j}))_{n \times n} \\ (g_{\mathbb{C}}(\frac{\partial}{\partial \bar{z}^i}, \frac{\partial}{\partial z^j}))_{n \times n} & (g_{\mathbb{C}}(\frac{\partial}{\partial \bar{z}^i}, \frac{\partial}{\partial \bar{z}^j}))_{n \times n} \end{pmatrix}$$

Direct computation shows that

$$\begin{aligned} g_{\mathbb{C}}\left(\frac{\partial}{\partial z^i}, \frac{\partial}{\partial z^j}\right) &= 0 \\ g_{\mathbb{C}}\left(\frac{\partial}{\partial \bar{z}^i}, \frac{\partial}{\partial \bar{z}^j}\right) &= 0 \end{aligned}$$

and if we denote $H = (h_{i\bar{j}})_{n \times n}$, where

$$(6.1) \quad h_{i\bar{j}} := g_{\mathbb{C}}\left(\frac{\partial}{\partial z^i}, \frac{\partial}{\partial \bar{z}^j}\right) = \frac{1}{2}(g_{ij} + \sqrt{-1}g_{iJ})$$

Then

$$G = \begin{pmatrix} 0 & H \\ \bar{H} & 0 \end{pmatrix}$$

Moreover, H is a positive definite Hermitian matrix since G is a positive definite symmetric matrix. Thus h gives a Hermitian metric on $T^{1,0}X$, which makes X a Hermitian manifold. Conversely, if h is a Hermitian metric on $T^{1,0}X$, there is also a Riemannian metric on $T_{\mathbb{R}}X$ given by

$$\begin{aligned} g_{ij} &= 2\operatorname{Re}h_{i\bar{j}} \\ g_{iJ} &= 2\operatorname{Im}h_{i\bar{j}} \\ g_{Ij} &= -g_{iJ} \\ g_{IJ} &= g_{ij} \end{aligned}$$

From now on, g always denotes a Riemannian metric on the underlying real manifold which is compatible with J , and h be the Hermitian metric corresponding to g .

Proposition 6.1.2.

$$\sqrt{\det g} = 2^n \det h$$

Proof. Direct computation shows

$$\begin{aligned} \det g &= \det \begin{pmatrix} (g_{ij})_{n \times n} & (g_{iJ})_{n \times n} \\ (g_{Ij})_{n \times n} & (g_{IJ})_{n \times n} \end{pmatrix} \\ &\stackrel{(1)}{=} \det \begin{pmatrix} (g_{ij})_{n \times n} & (g_{iJ})_{n \times n} \\ (-g_{iJ})_{n \times n} & (g_{ij})_{n \times n} \end{pmatrix} \\ &= \det \begin{pmatrix} (g_{ij} + \sqrt{-1}g_{iJ})_{n \times n} & (g_{iJ})_{n \times n} \\ (-g_{iJ} + \sqrt{-1}g_{ij})_{n \times n} & (g_{ij})_{n \times n} \end{pmatrix} \\ &= \det \begin{pmatrix} (g_{ij} + \sqrt{-1}g_{iJ})_{n \times n} & (g_{iJ})_{n \times n} \\ \mathbf{O} & (g_{ij} + \sqrt{-1}g_{iJ})_{n \times n} \end{pmatrix} \\ &= (2^n \det h)^2 \end{aligned}$$

where (1) holds from Proposition 6.1.1. □

Notation 6.1.2. $h^{i\bar{j}}$ is defined by the (i, j) -entry of $(H^{-1})^T$, that is $h^{i\bar{l}}h_{j\bar{l}} = \delta_j^i$, and $h^{\bar{i}j} := \overline{h^{i\bar{j}}}$. Note that

$$G^{-1} = (G^{-1})^T = \begin{pmatrix} 0 & (h^{i\bar{j}})_{n \times n} \\ (h^{\bar{i}j})_{n \times n} & 0 \end{pmatrix}$$

Proposition 6.1.3.

$$h^{\bar{i}j} = 2(g^{ij} - \sqrt{-1}g^{iJ})$$

Proof. Direct computation shows

$$\begin{aligned} h^{i\bar{l}}h_{j\bar{l}} &= (g^{il} - \sqrt{-1}g^{iL})(g_{jl} + \sqrt{-1}g_{jL}) \\ &= g^{il}g_{jl} + g^{iL}g_{jL} + \sqrt{-1}(g^{il}g_{jL} - g^{iL}g_{jL}) \\ &= g^{il}g_{lj} + g^{iL}g_{Lj} - \sqrt{-1}(g^{il}g_{lJ} + g^{iL}g_{Lj}) \\ &= \delta_j^i \end{aligned}$$

□

Definition 6.1.2 (fundamental form). The fundamental form ω of g is defined as

$$\omega(V, W) := g_{\mathbb{C}}(JV, W)$$

where $V, W \in C^\infty(X, T_{\mathbb{C}}X)$.

Proposition 6.1.4.

$$\omega = \sqrt{-1}h_{i\bar{j}}dz^i \wedge d\bar{z}^j$$

In particular, ω is a real $(1, 1)$ -form.

Proof. Direct computation shows

$$\begin{aligned} \omega\left(\frac{\partial}{\partial z^i}, \frac{\partial}{\partial z^j}\right) &= g_{\mathbb{C}}\left(J\left(\frac{\partial}{\partial z^i}\right), \frac{\partial}{\partial z^j}\right) = \sqrt{-1}g_{\mathbb{C}}\left(\frac{\partial}{\partial z^i}, \frac{\partial}{\partial z^j}\right) = 0 \\ \omega\left(\frac{\partial}{\partial z^i}, \frac{\partial}{\partial \bar{z}^j}\right) &= g_{\mathbb{C}}\left(J\left(\frac{\partial}{\partial z^i}\right), \frac{\partial}{\partial \bar{z}^j}\right) = \sqrt{-1}g_{\mathbb{C}}\left(\frac{\partial}{\partial z^i}, \frac{\partial}{\partial \bar{z}^j}\right) = \sqrt{-1}h_{i\bar{j}} \\ \omega\left(\frac{\partial}{\partial \bar{z}^j}, \frac{\partial}{\partial z^i}\right) &= g_{\mathbb{C}}\left(J\left(\frac{\partial}{\partial \bar{z}^j}\right), \frac{\partial}{\partial z^i}\right) = -\sqrt{-1}g_{\mathbb{C}}\left(\frac{\partial}{\partial \bar{z}^j}, \frac{\partial}{\partial z^i}\right) = -\sqrt{-1}h_{\bar{j}i} \\ \omega\left(\frac{\partial}{\partial \bar{z}^i}, \frac{\partial}{\partial \bar{z}^j}\right) &= g_{\mathbb{C}}\left(J\left(\frac{\partial}{\partial \bar{z}^i}\right), \frac{\partial}{\partial \bar{z}^j}\right) = -\sqrt{-1}g_{\mathbb{C}}\left(\frac{\partial}{\partial \bar{z}^i}, \frac{\partial}{\partial \bar{z}^j}\right) = 0 \end{aligned}$$

This shows

$$\begin{aligned} \omega &= \sqrt{-1}h_{i\bar{j}}dz^i \otimes d\bar{z}^j - \sqrt{-1}h_{\bar{j}i}d\bar{z}^j \otimes dz^i \\ &= \sqrt{-1}h_{i\bar{j}}dz^i \otimes d\bar{z}^j - \sqrt{-1}h_{\bar{j}i}d\bar{z}^j \otimes dz^i \\ &= \sqrt{-1}h_{i\bar{j}}dz^i \wedge d\bar{z}^j \end{aligned}$$

where the last step holds since h is Hermitian, that is $h_{\bar{j}i} = h_{i\bar{j}}$. □

Proposition 6.1.5.


$$2h + \sqrt{-1}\omega = g_{\mathbb{C}}$$

Proof. Direct computation shows

$$\begin{aligned}
h - \frac{1}{2}g_{\mathbb{C}} &= h_{i\bar{j}}dz^i \otimes d\bar{z}^j - \frac{1}{2}(h_{i\bar{j}}dz^i \otimes d\bar{z}^j + h_{\bar{i}j}d\bar{z}^i \otimes dz^j) \\
&= \frac{1}{2}h_{i\bar{j}}dz^i \otimes d\bar{z}^j - \frac{1}{2}h_{\bar{i}j}d\bar{z}^i \otimes dz^j \\
&= \frac{1}{2}h_{i\bar{j}}dz^i \wedge d\bar{z}^j \\
&= -\frac{\sqrt{-1}}{2}\omega
\end{aligned}$$

□

Remark 6.1.1. In fact, Hermitian metric h , Riemannian metric g on underlying real manifold and fundamental form ω are the same things on a complex manifold X , and any of them gives a Hermitian structure on X .



The explicit correspondences are listed as follows:

- 1 $2\omega(-, -) = -\text{Im}h(-, -)$
- 2 $2h(-, -) = \omega(-, J-) - \sqrt{-1}\omega(-, -)$
- 3 $2g_{\mathbb{C}}(-, -) = \text{Re}h(-, -)$
- 4 $2h(-, -) = g_{\mathbb{C}}(-, -) - \sqrt{-1}g_{\mathbb{C}}(J-, -)$
- 5 $g_{\mathbb{C}}(-, -) = \omega(-, J-)$
- 6 $\omega(-, -) = g_{\mathbb{C}}(J-, -)$

In later discussion, we may say a Hermitian manifold (X, h) or (X, ω) when we're emphasizing its Hermitian metric or fundamental form.

Theorem 6.1.1 (normal coordinate). Let (X, h) be a Hermitian manifold. For any $p \in X$, there exists a local holomorphic coordinate $\{z^i\}$ centered at p such that

$$h_{i\bar{j}}(p) = \delta_{i\bar{j}} \quad \text{and} \quad \frac{\partial h_{i\bar{j}}}{\partial \bar{z}^k} + \frac{\partial h_{i\bar{k}}}{\partial \bar{z}^j} = 0$$

Proof. Without lose of generality, we may assume

$$\omega = \sqrt{-1}(\delta_{i\bar{j}} + a_{i\bar{j}l}w^l + a_{\bar{i}jl}\bar{w}^l + O(|w|^2))dw^i \wedge d\bar{w}^j$$

□

6.2. Curvatures of Hermitian manifold. Sometimes we need to consider Hermitian holomorphic vector bundles over a Hermitian manifold, so there are two Hermitian metrics. In this case, in order to distinguish them, we always say a Hermitian holomorphic vector bundle (E, h) over a Hermitian manifold (X, g) .

Definition 6.2.1 (curvatures of vector bundle). Let (E, h) be a Hermitian holomorphic vector bundle on Hermitian manifold (X, g) . Then

- (1) the first Chern-Ricci curvature of (E, h) is locally given by

$$\text{Ric}^{(1)}(h) = \sqrt{-1} h^{\alpha\bar{\beta}} \Theta_{i\bar{j}\alpha\bar{\beta}} dz^i \wedge d\bar{z}^j$$

- (2) the second Chern-Ricci curvature of (E, h) is locally given by

$$\text{Ric}^{(2)}(h) = \sqrt{-1} g^{i\bar{j}} \Theta_{i\bar{j}\alpha\bar{\beta}} e^\alpha \otimes \bar{e}^\beta$$

- (3) the Chern scalar curvature of (E, h) is locally given by

$$s = g^{i\bar{j}} h^{\alpha\bar{\beta}} \Theta_{i\bar{j}\alpha\bar{\beta}}$$

Remark 6.2.1. For convenience, we always use the following notations.

- (1) $\text{Ric}^{(1)}(h) = \sqrt{-1} \text{tr}_h \Theta_h$ and $\text{Ric}^{(2)}(h) = \sqrt{-1} \text{tr}_g \Theta_h$, where Θ_h is the Chern curvature of (E, h) .
(2) For a real $(1, 1)$ -form φ locally written as $\sqrt{-1} \varphi_{i\bar{j}} dz^i \wedge d\bar{z}^j$, $\text{tr}_g \varphi$ denotes the function $g^{i\bar{j}} \varphi_{i\bar{j}}$. In particular, the Chern scalar curvature is denoted by $\text{tr}_g \text{tr}_h \Theta_h$, where Θ_h is the Chern curvature of (E, h) .

Definition 6.2.2 (curvatures of Hermitian manifold). Let (X, h) be a Hermitian manifold. Then

- (1) the first (or second) Chern-Ricci curvature of (X, h) is defined to be the first (or second) Chern-Ricci curvature of its holomorphic tangent bundle.
(2) the Chern scalar curvature of (X, h) is defined to be the Chern scalar curvature of its holomorphic tangent bundle.

Proposition 6.2.1. Let (E, h) be a Hermitian holomorphic vector bundle. The first Chern-Ricci curvature of (E, h) gives the first Chern class of (E, h) up to a scalar.

Proof. A direct computation shows

$$\frac{\sqrt{-1}}{2\pi} h^{\alpha\bar{\beta}} \left(-\frac{\partial^2 h_{\alpha\bar{\beta}}}{\partial z^i \partial \bar{z}^j} + h^{\gamma\bar{\delta}} \frac{\partial h_{\alpha\bar{\delta}}}{\partial z^i} \frac{\partial h_{\gamma\bar{\beta}}}{\partial \bar{z}^j} \right) = -\frac{\sqrt{-1}}{2\pi} \frac{\partial^2 \log \det(h_{\alpha\bar{\beta}})}{\partial z^i \partial \bar{z}^j}$$

Thus the first Chern-Ricci curvature of (E, h) gives the Chern curvature of $(\det E, \det h)$ up a scalar $1/2\pi$, and by Definition 5.3.2 the first Chern class of (E, h) is defined to be first Chern class of $\det E$. \square

Definition 6.2.3 (holomorphic sectional curvature). Let (X, h) be a Hermitian manifold and $v = v^i \frac{\partial}{\partial z^i} \in T_p X$ be a unit vector. The holomorphic sectional curvature in the direction v is defined as

$$\text{HSC}_p(v) := \Theta_{i\bar{j}k\bar{l}} v^i \bar{v}^j v^k \bar{v}^l$$

Definition 6.2.4 (holomorphic bisectonal curvature). Let (X, h) be a Hermitian manifold and $v = v^i \frac{\partial}{\partial z^i}, w = w^i \frac{\partial}{\partial z^i} \in T_p X$ be unit vectors. The holomorphic sectional curvature in the direction v, w is defined as

$$\text{HBSC}_p(v, w) := \Theta_{i\bar{j}k\bar{l}} v^i \bar{v}^j w^k \bar{w}^l$$

6.3. Useful formulas of Hermitian geometry. In this section we collect some useful formulas in Hermitian geometry. Unless otherwise specified, we assume (X, h) is a Hermitian n -manifold with fundamental form ω , and g is the Riemannian metric on the underlying real manifold.

Proposition 6.3.1.

$$\omega^n = (\sqrt{-1})^n n! \det(h_{i\bar{j}}) dz^1 \wedge d\bar{z}^1 \wedge \cdots \wedge dz^n \wedge d\bar{z}^n$$

Proof. Direct computation shows

$$\begin{aligned} \omega^n &= (\sqrt{-1})^n (h_{i\bar{j}} dz^i \wedge d\bar{z}^j)^n \\ &= (\sqrt{-1})^n \sum_{\substack{\sigma \in S_n \\ \tau \in S_n}} h_{i_{\sigma(1)} \bar{j}_{\tau(1)}} \cdots h_{i_{\sigma(n)} \bar{j}_{\tau(n)}} dz^{i_{\sigma(1)}} \wedge d\bar{z}^{j_{\tau(1)}} \wedge \cdots \wedge dz^{i_{\sigma(n)}} \wedge d\bar{z}^{j_{\tau(n)}} \\ &= (\sqrt{-1})^n \sum_{\substack{\sigma \in S_n \\ \tau \in S_n}} (-1)^{|\sigma|} (-1)^{|\tau|} h_{i_{\sigma(1)} \bar{j}_{\tau(1)}} \cdots h_{i_{\sigma(n)} \bar{j}_{\tau(n)}} dz^1 \wedge d\bar{z}^1 \wedge \cdots \wedge dz^n \wedge d\bar{z}^n \\ &= (\sqrt{-1})^n \sum_{\sigma \in S_n} \sum_{\rho \in S_n} (-1)^{|\rho|} h_{i_{\rho(1)} \bar{j}_1} \cdots h_{i_{\rho(n)} \bar{j}_n} dz^1 \wedge d\bar{z}^1 \wedge \cdots \wedge dz^n \wedge d\bar{z}^n \\ &= (\sqrt{-1})^n n! \det(h_{i\bar{j}}) dz^1 \wedge d\bar{z}^1 \wedge \cdots \wedge dz^n \wedge d\bar{z}^n \end{aligned}$$

□

Corollary 6.3.1.

$$\omega^{n-1} = (\sqrt{-1})^{n-1} (n-1)! \sum_{\substack{i_1 < \cdots < i_{n-1}, i_l \neq p \\ j_1 < \cdots < j_{n-1}, j_l \neq q}} h(p, q) dz^{i_1} \wedge d\bar{z}^{j_1} \wedge \cdots \wedge dz^{i_{n-1}} \wedge d\bar{z}^{j_{n-1}}$$

where $h(p, q)$ the cofactor of h without row p and column q .

Proof. Direct computation shows

$$\begin{aligned}
\omega^{n-1} &= (\sqrt{-1})^{n-1} (h_{i\bar{j}} dz^i \wedge d\bar{z}^j)^{n-1} \\
&= (\sqrt{-1})^{n-1} \sum_{\substack{\sigma \in S_{n-1} \\ \tau \in S_{n-1}}} h_{i_{\sigma(1)} \bar{j}_{\tau(1)}} \cdots h_{i_{\sigma(n-1)} \bar{j}_{\tau(n-1)}} dz^{i_{\sigma(1)}} \wedge d\bar{z}^{j_{\tau(1)}} \wedge \cdots \wedge dz^{i_{\sigma(n-1)}} \wedge d\bar{z}^{j_{\tau(n-1)}} \\
&= (\sqrt{-1})^{n-1} \sum_{\substack{i_1 < \cdots < i_{n-1}, i_l \neq p \\ j_1 < \cdots < j_{n-1}, j_l \neq q}} \sum_{\substack{\sigma \in S_{n-1} \\ \rho \in S_{n-1}}} (-1)^{|\rho|} h_{i_{\rho(1)} \bar{j}_1} \cdots h_{i_{\rho(n-1)} \bar{j}_{n-1}} dz^{i_1} \wedge d\bar{z}^{j_1} \wedge \cdots \wedge dz^{i_{n-1}} \wedge d\bar{z}^{j_{n-1}} \\
&= (\sqrt{-1})^{n-1} (n-1)! \sum_{\substack{i_1 < \cdots < i_{n-1}, i_l \neq p \\ j_1 < \cdots < j_{n-1}, j_l \neq q}} h(p, q) dz^{i_1} \wedge d\bar{z}^{j_1} \wedge \cdots \wedge dz^{i_{n-1}} \wedge d\bar{z}^{j_{n-1}}
\end{aligned}$$

□

Corollary 6.3.2.

$$\omega^{n-2} = (\sqrt{-1})^{n-2} (n-2)! \sum_{\substack{i_1 < \cdots < i_{n-2}, i_l \neq p, s \\ j_1 < \cdots < j_{n-2}, j_l \neq q, t}} h \begin{pmatrix} p, q \\ s, t \end{pmatrix} dz^{i_1} \wedge d\bar{z}^{j_1} \wedge \cdots \wedge dz^{i_{n-2}} \wedge d\bar{z}^{j_{n-2}}$$

where $h \begin{pmatrix} p, q \\ s, t \end{pmatrix}$ the cofactor of h without rows p, s and columns q, t .

Proposition 6.3.2. $\omega^n/n!$ is the volume form of the underlying real manifold with respect to g .

Proof. Direct computation shows

$$\begin{aligned}
\omega^n &\stackrel{(1)}{=} (\sqrt{-1})^n n! \det(h_{i\bar{j}}) dz^1 \wedge d\bar{z}^1 \wedge \cdots \wedge dz^n \wedge d\bar{z}^n \\
&\stackrel{(2)}{=} n! 2^n \det(h_{i\bar{j}}) dx^1 \wedge dy^1 \wedge \cdots \wedge dx^n \wedge dy^n \\
&\stackrel{(3)}{=} n! \text{vol}
\end{aligned}$$

where

- (1) holds from Proposition 6.3.1.
- (2) holds from Exercise 4.2.1.
- (3) holds from Proposition 6.1.2.

□

Proposition 6.3.3.

(1) If φ is a real $(1, 1)$ -form, then

$$\varphi \wedge \omega^{n-1} = \frac{1}{n} \text{tr}_\omega \varphi \cdot \omega^n$$

(2) If φ is a $(1, 0)$ -form, then

$$\sqrt{-1} \varphi \wedge \bar{\varphi} \wedge \omega^{n-1} = \frac{1}{n} |\varphi|^2 \cdot \omega^n$$

(3) If φ is a $(0, 1)$ -form, then

$$\sqrt{-1}\varphi \wedge \bar{\varphi} \wedge \omega^{n-1} = -\frac{1}{n}|\varphi|^2 \cdot \omega^n$$

Proof. For (1). Suppose $\varphi = \sqrt{-1}\varphi_{i\bar{j}}dz^i \wedge d\bar{z}^j$. Then by Proposition 6.3.1 one has

$$\frac{1}{n} \operatorname{tr}_\omega \varphi \cdot \omega^n = (\sqrt{-1})^n (n-1)! h^{i\bar{j}} \varphi_{i\bar{j}} \det h dz^1 \wedge d\bar{z}^1 \wedge \cdots \wedge dz^n \wedge d\bar{z}^n$$

and by Corollary 6.3.1 one has

$$\begin{aligned} \varphi \wedge \omega^{n-1} &= (\sqrt{-1})^n (n-1)! \varphi_{i\bar{j}} dz^i \wedge d\bar{z}^j \wedge \sum_{\substack{i_1 < \cdots < i_{n-1}, i_l \neq p \\ j_1 < \cdots < j_{n-1}, j_l \neq q}} h(p, q) dz^{i_1} \wedge d\bar{z}^{j_1} \wedge \cdots \wedge dz^{i_{n-1}} \wedge d\bar{z}^{j_{n-1}} \\ &= (\sqrt{-1})^n (n-1)! \sum_{1 \leq p, q \leq n} \varphi_{p\bar{q}} (-1)^{p+q} h(p, q) dz^1 \wedge d\bar{z}^1 \wedge \cdots \wedge dz^n \wedge d\bar{z}^n \end{aligned}$$

Note that by expansion of determinant one has

$$h^{i\bar{j}} \varphi_{i\bar{j}} \det h = h^{i\bar{j}} \varphi_{i\bar{j}} \sum_{k=1}^n h_{k\bar{j}} (-1)^{k+j} h(k, j) = \sum_{1 \leq i, j \leq n} \varphi_{i\bar{j}} (-1)^{i+j} h(i, j)$$

For (2). Suppose $\varphi = \varphi_i dz^i$. Then

$$\sqrt{-1}\varphi \wedge \bar{\varphi} = \sqrt{-1}\varphi_i \bar{\varphi}_{\bar{j}} dz^i \wedge d\bar{z}^j$$

is a real $(1, 1)$ -form, and it's clear

$$\operatorname{tr}_\omega \sqrt{-1}\varphi \wedge \bar{\varphi} = |\varphi|^2$$

then by (1) we obtain desired result, and (3) follows from (2) directly. \square

Proposition 6.3.4.

$$\langle dz^i \wedge \alpha, \beta \rangle = \langle \alpha, h^{p\bar{i}} t_p \beta \rangle$$

holds for α, β with appropriate bidegrees.

Proposition 6.3.5. Let (E, h) be a Hermitian holomorphic vector bundle over a Hermitian manifold (X, g) and φ be a real $(1, 1)$ -form. Then

$$\{\varphi, \varphi\} \frac{\omega^{n-2}}{(n-2)!} = (|\operatorname{tr}_\omega \varphi|^2 - |\varphi|^2) \frac{\omega^n}{n!}$$

Proof. Suppose $\varphi = \sqrt{-1}\varphi_{i\bar{j}}^\alpha dz^i \wedge d\bar{z}^j \otimes e_\alpha$. Then by Corollary 6.3.2 one has

$$\text{LHS} = (\sqrt{-1})^n \varphi_{i\bar{j}}^\alpha \bar{\varphi}_{k\bar{l}}^\beta h_{\alpha\bar{\beta}} dz^i \wedge d\bar{z}^j \wedge dz^k \wedge d\bar{z}^l \wedge \sum_{\substack{i_1 < \cdots < i_{n-2}, i_l \neq p, s \\ j_1 < \cdots < j_{n-2}, j_l \neq q, t}} g \binom{p, q}{s, t} dz^{i_1} \wedge d\bar{z}^{j_1} \wedge \cdots \wedge dz^{i_{n-2}} \wedge d\bar{z}^{j_{n-2}}$$

If we want to insert $dz^i \wedge d\bar{z}^j \wedge dz^l \wedge d\bar{z}^k$ into $\sum_{\substack{i_1 < \dots < i_{n-2}, i_l \neq p, s \\ j_1 < \dots < j_{n-2}, j_l \neq q, t}} dz^{i_1} \wedge d\bar{z}^{j_1} \wedge \dots \wedge dz^{i_{n-2}} \wedge d\bar{z}^{j_{n-2}}$, there are the following four cases

$$\begin{cases} i = p, l = s, j = q, k = t \\ i = p, l = s, j = t, k = q \\ i = s, l = p, j = q, k = t \\ i = s, l = p, j = t, k = q \end{cases}$$

So case by case one has

$$\begin{aligned} & \varphi_{ij}^\alpha \overline{\varphi_{kl}^\beta} dz^i \wedge d\bar{z}^j \wedge dz^l \wedge d\bar{z}^k \wedge \sum_{\substack{i_1 < \dots < i_{n-2}, i_l \neq p, s \\ j_1 < \dots < j_{n-2}, j_l \neq q, t}} dz^{i_1} \wedge d\bar{z}^{j_1} \wedge \dots \wedge dz^{i_{n-2}} \wedge d\bar{z}^{j_{n-2}} \\ = & \sum_{\substack{1 \leq p, q \leq n \\ 1 \leq s, t \leq n}} (\varphi_{pq}^\alpha \overline{\varphi_{ts}^\beta} - \varphi_{pt}^\alpha \overline{\varphi_{qs}^\beta} - \varphi_{sq}^\alpha \overline{\varphi_{tp}^\beta} + \varphi_{st}^\alpha \overline{\varphi_{qp}^\beta}) (-1)^{p+q+t+s} dz^1 \wedge d\bar{z}^1 \wedge \dots \wedge dz^n \wedge d\bar{z}^n \end{aligned}$$

On the other hand, direct computation shows

$$\begin{aligned} |\mathrm{tr}_\omega \varphi|^2 &= \varphi_{ij}^\alpha \overline{\varphi_{kl}^\beta} g^{i\bar{j}} g^{l\bar{k}} h_{\alpha\bar{\beta}} \\ |\varphi^2| &= \varphi_{ij}^\alpha \overline{\varphi_{kl}^\beta} g^{i\bar{k}} g^{l\bar{j}} h_{\alpha\bar{\beta}} \end{aligned}$$

and thus

$$\mathrm{RHS} = (\sqrt{-1})^n \varphi_{ij}^\alpha \overline{\varphi_{kl}^\beta} h_{\alpha\bar{\beta}} (g^{i\bar{j}} g^{l\bar{k}} - g^{i\bar{k}} g^{l\bar{j}}) \det g$$

Note that Laplacian's theorem implies

$$\begin{aligned} (g^{i\bar{j}} g^{l\bar{k}} - g^{i\bar{k}} g^{l\bar{j}}) \det g &= (g^{i\bar{j}} g^{l\bar{k}} - g^{i\bar{k}} g^{l\bar{j}}) \sum_{\substack{1 \leq p, s \leq n \\ 1 \leq j, k \leq n}} (g_{p\bar{j}} g_{s\bar{k}} - g_{s\bar{j}} g_{p\bar{k}}) (-1)^{p+j+s+k} g \binom{p, j}{s, k} \\ &= (\delta_p^i \delta_s^l - \delta_s^i \delta_p^l - \delta_s^i \delta_p^l + \delta_p^i \delta_s^l) (-1)^{p+j+s+k} g \binom{p, j}{s, k} \end{aligned}$$

This shows the RHS equals to the LHS. \square

6.4. Gauduchon metric.

Theorem 6.4.1 (Gauduchon metric). If X be a complex manifold, then there exists a Hermitian metric ω such that

$$\partial\bar{\partial}\omega^{n-1} = 0,$$

which is called Gauduchon metric.

Proof. See [Gau77]. \square

Corollary 6.4.1. Let $f: X \rightarrow \mathbb{R}$ be a smooth function on a compact complex manifold X . If

$$\sqrt{-1}\partial\bar{\partial}f \geq 0,$$

then f is a constant.

Proof. Let ω be the Gauduchon metric on X . Then

$$0 \leq \int_X \sqrt{-1} \partial \bar{\partial} f \wedge \omega^{n-1} = \int_X \sqrt{-1} f \wedge \partial \bar{\partial} \omega^{n-1} = 0$$

which implies $\sqrt{-1} \partial \bar{\partial} f = 0$. Note that

$$(6.2) \quad \partial \bar{\partial} f^2 = \partial(2f \bar{\partial} f) = 2\partial f \wedge \bar{\partial} f + 2f \partial \bar{\partial} f = 2\partial f \wedge \bar{\partial} f$$

Then

$$\begin{aligned} 0 &\stackrel{(1)}{=} \int_X \sqrt{-1} \partial \bar{\partial} f^2 \wedge \omega^{n-1} \\ &\stackrel{(2)}{=} 2 \int_X \sqrt{-1} \partial f \wedge \bar{\partial} f \wedge \omega^{n-1} \\ &\stackrel{(3)}{=} \frac{2}{n} \int_X |\partial f|^2 \omega^n \end{aligned}$$

where

(1) holds from ω is a Gauduchon metric.

(2) holds from equation (6.2).

(3) holds from (2) of Proposition 6.3.3.

This shows $\partial f = 0$, and this also shows $df = 0$ since f is real-valued, that is f is a constant. \square

Corollary 6.4.2. Let $f: X \rightarrow \mathbb{R}$ be a smooth function on a compact complex manifold X . If there exists a Hermitian metric ω such that

$$\sqrt{-1} \operatorname{tr}_\omega \partial \bar{\partial} f \geq 0,$$

then f is a constant.

Proof. By (1) of Proposition 6.3.3 one has

$$\int_X \sqrt{-1} \operatorname{tr}_\omega \partial \bar{\partial} f \wedge \omega^n = \int_X n \sqrt{-1} \partial \bar{\partial} f \wedge \omega^{n-1}$$

The argument in above corollary still works. \square

6.5. Second fundamental form. Let (E, h) be a Hermitian holomorphic vector bundle over complex manifold X with rank r and S be a holomorphic subbundle of E with rank s . Then there is an exact sequence of holomorphic vector bundles

$$0 \rightarrow S \rightarrow E \rightarrow Q \rightarrow 0$$

where Q is the quotient bundle, which is isomorphic to S^\perp as complex vector bundle.

6.5.1. *Second fundamental form of subbundle.* Suppose ∇^E is the Chern connection on E and define $\nabla^S := \pi_S \circ \nabla^E$, where $\pi_S: E \rightarrow S$ is the orthogonal projection.

- (1) It's clear ∇^S is compatible with complex structure of S since ∇^E is Chern connection of E , and S is holomorphic subbundle of E .
- (2) For sections s, t of S , one has

$$\begin{aligned} dh(s, t) &= h(\nabla^E s, t) + h(s, \nabla^E t) \\ &\stackrel{(a)}{=} h(\pi_S \circ \nabla^E s, t) + h(s, \pi_S \circ \nabla^E t) \\ &= h(\nabla^S s, t) + h(s, \nabla^S t), \end{aligned}$$

where (a) holds from π_S is orthogonal projection.

This shows that ∇^S is the Chern connection of S with respect to Hermitian metric induced by the one on E .

Definition 6.5.1 (second fundamental form). The second fundamental form of the subbundle S of E is defined as

$$B = \nabla^E - \nabla^S: C^\infty(X, S) \rightarrow C^\infty(X, \Omega_X^{1,0} \otimes Q).$$

Remark 6.5.1 (local form). For $p \in X$, suppose $\{e_\alpha\}_{1 \leq \alpha \leq r}$ is a holomorphic local frame of E such that $\{e_\alpha\}_{1 \leq \alpha \leq s}$ is a holomorphic local frame of S , and for convenience we assume $h_{\alpha\bar{\beta}}(p) = \delta_{\alpha\bar{\beta}}$. By formula of Chern connection, for $1 \leq \alpha \leq s$, one has

$$\begin{aligned} \nabla^E e_\alpha(p) &= \sum_{\beta=1}^r \frac{h_{\alpha\bar{\beta}}}{\partial z^i}(p) dz^i \otimes e_\beta \\ \nabla^S e_\alpha(p) &= \sum_{\beta=1}^s \frac{h_{\alpha\bar{\beta}}}{\partial z^i}(p) dz^i \otimes e_\beta \end{aligned}$$

Then for $1 \leq \alpha \leq s$, one has

$$B e_\alpha(p) = \sum_{\beta=s+1}^r \frac{h_{\alpha\bar{\beta}}}{\partial z^i}(p) dz^i \otimes e_\beta$$

Thus with respect to local frame we choose, one has

$$B(p) = \sum_{\alpha=1}^s \sum_{\beta=s+1}^r \frac{h_{\alpha\bar{\beta}}}{\partial z^i}(p) dz^i \otimes e^\alpha \otimes e_\beta$$

One has its conjugate transpose is

$$B^*(p) = \sum_{\beta=s+1}^r \sum_{\alpha=1}^s \frac{h_{\beta\bar{\alpha}}}{\partial \bar{z}^j}(p) d\bar{z}^j \otimes e^\beta \otimes e_\alpha$$

then $B^* \wedge B$ gives a section of $\Omega_X^{1,1} \otimes S^* \otimes S$. To be explicit, for $1 \leq \alpha \leq s$, one has

$$\begin{aligned} B^* \wedge B e_\alpha(p) &= B^* \left(\sum_{\gamma=s+1}^r \frac{h_{\alpha\bar{\gamma}}}{\partial z^i}(p) dz^i \otimes e_\gamma \right) \\ &= - \sum_{\beta=1}^s \sum_{\gamma=s+1}^r \frac{h_{\alpha\bar{\gamma}}}{\partial z^i} \frac{h_{\gamma\bar{\beta}}}{\partial \bar{z}^j}(p) dz^i \wedge d\bar{z}^j e_\beta \end{aligned}$$

which implies

$$B^* \wedge B(p) = - \sum_{\alpha, \beta=1}^s \left\{ \sum_{\gamma=s+1}^r \frac{\partial h_{\alpha\bar{\gamma}}}{\partial z^i} \frac{\partial h_{\gamma\bar{\beta}}}{\partial \bar{z}^j}(p) \right\} dz^i \wedge d\bar{z}^j \otimes e^\alpha \otimes e_\beta$$

On the other hand, direct computation shows

$${}^E\Theta|_S(p) - {}^S\Theta(p) = \sum_{\alpha, \beta=1}^s \left\{ \sum_{\gamma=s+1}^r \frac{\partial h_{\alpha\bar{\gamma}}}{\partial z^i} \frac{\partial h_{\gamma\bar{\beta}}}{\partial \bar{z}^j}(p) \right\} dz^i \wedge d\bar{z}^j \otimes e^\alpha \otimes \bar{e}^\beta$$

Thus

$${}^S\Theta = {}^E\Theta|_S + B^* \wedge B$$

where B^* is conjugate transpose of B since $e_\beta = \delta_{\beta\bar{\beta}} \bar{e}^\beta$.

6.5.2. *Second fundamental form of quotient bundle.* Here we consider the following exact sequence

$$0 \rightarrow Q^* \rightarrow E^* \rightarrow S^* \rightarrow 0$$

The second fundamental form C of quotient bundle Q is defined as the second fundamental form of subbundle Q^* .

7. KÄHLER GEOMETRY

7.1. Kähler manifold.

Definition 7.1.1 (Kähler manifold). A Hermitian manifold (X, h) is called a Kähler manifold, if its fundamental form ω is d-closed⁸.

Remark 7.1.1. Note that $d\omega = 0$ is equivalent to $\partial\omega = 0$, and is also equivalent to $\bar{\partial}\omega = 0$ since ω is a real $(1, 1)$ -form.

Remark 7.1.2 (local form). By Proposition 6.1.4 one has

$$\omega = \sqrt{-1}h_{i\bar{j}}dz^i \wedge d\bar{z}^j$$

So Kähler condition $d\omega = 0$ can be computed explicitly as follows

$$\begin{aligned} d\omega &= \sqrt{-1}d(h_{i\bar{j}}dz^i \wedge d\bar{z}^j) \\ &= \sqrt{-1}\left(\frac{\partial h_{i\bar{j}}}{\partial z^k}dz^k \wedge dz^i \wedge d\bar{z}^j - \frac{\partial h_{i\bar{j}}}{\partial \bar{z}^k}dz^i \wedge dz^k \wedge d\bar{z}^j\right) \\ &= 0 \end{aligned}$$

So locally Kähler condition can be written as follows

$$\begin{aligned} \partial_k h_{i\bar{j}} &= \partial_i h_{k\bar{j}} \\ \partial_{\bar{k}} h_{i\bar{j}} &= \partial_{\bar{j}} h_{i\bar{k}} \end{aligned}$$

holds for all i, j, k .

Proposition 7.1.1. Let (X, h) be a Kähler manifold. Then the first Chern-Ricci curvature coincides with the second Chern-Ricci curvature.

Proof. Note that

$$\Theta_{i\bar{j}k\bar{l}} = -\frac{\partial^2 h_{i\bar{j}}}{\partial z^k \partial \bar{z}^l} + h^{p\bar{q}} \frac{\partial h_{k\bar{q}}}{\partial z^i} \frac{\partial h_{p\bar{l}}}{\partial \bar{z}^j}$$

Thus if (X, h) is Kähler, then

$$\Theta_{i\bar{j}k\bar{l}} = \Theta_{k\bar{j}i\bar{l}} = \Theta_{i\bar{l}k\bar{j}}$$

As a consequence, one has

$$\text{Ric}^{(1)}(h) = \sqrt{-1}h^{k\bar{l}}\Theta_{i\bar{j}k\bar{l}}dz^i \wedge d\bar{z}^j = \sqrt{-1}h^{k\bar{l}}\Theta_{k\bar{l}i\bar{j}}dz^i \wedge d\bar{z}^j = \text{Ric}^{(2)}(h)$$

This completes the proof. \square

Definition 7.1.2 (Kähler-Einstein metric). A Kähler metric ω is called a Kähler-Einstein metric, if there exists $\lambda \in \mathbb{R}$ such that

$$\text{Ric}(\omega) = \lambda\omega$$

Example 7.1.1. Any complex curve⁹ X is Kähler since $d\omega = 0$ automatically holds.

⁸ ω is called Kähler form and h is called Kähler metric.

⁹In other words, a Riemann surface.

Proposition 7.1.2. A submanifold of a Kähler manifold is still Kähler.

Proof. If (X, ω) is a Kähler manifold and Y is a submanifold, the restriction of ω to Y gives Kähler form of Y . \square

Proposition 7.1.3. Let (X, ω) be a compact Kähler n -manifold. Then $H^{2k}(X, \mathbb{R}) \neq 0$ for $0 \leq k \leq n$.

Proof. Note that $d(\omega^k) = 0$ holds for $0 \leq k \leq n$ since $d\omega = 0$, that is $[\omega^k] \in H^{2k}(X, \mathbb{R})$. By Proposition 6.3.2 one has $\omega^n = n! \text{vol}$, so the integral pairing

$$\int_X \omega^k \wedge \omega^{n-k} = n! \int_X \text{vol} \neq 0$$

implies $[\omega^k] \neq 0$ for $0 \leq k \leq n$. \square

Theorem 7.1.1. Let (X, h) be a Kähler manifold. Then there exists a holomorphic coordinate (z^1, \dots, z^n) such that

$$h_{i\bar{j}}(z) = \delta_{i\bar{j}} - \Theta_{i\bar{j}k\bar{l}}(p) z^k \bar{z}^l + O(|z|^2).$$

7.2. Levi-Civita connection encounters Chern connection. Let X be a complex n -manifold and $\{z^i = x^i + \sqrt{-1}x^I\}$ be a local coordinate of X , where $1 \leq i \leq n$ and $n+1 \leq I = i+n \leq 2n$. Then $\{x^i, x^I\}$ gives a local coordinate of underlying real manifold of X . Let g be a Riemannian metric on $T_{\mathbb{R}}X$ which is compatible with natural almost complex structure J on $T_{\mathbb{R}}X$ with Levi-Civita connection ∇ . Now consider \mathbb{C} -linear extension of ∇

$$\tilde{\nabla}: C^\infty(X, T_{\mathbb{C}}X) \rightarrow C^\infty(X, \Omega_{X, \mathbb{C}}^1 \otimes T_{\mathbb{C}}X)$$

It's clear $\tilde{\nabla}$ gives a connection on $T_{\mathbb{C}}X$.

Notation 7.2.1. For $A \in \{1, \dots, n, \bar{1}, \dots, \bar{n}\}$, we denote

$$z^A = \begin{cases} z^i & A = i \\ \bar{z}^i & A = \bar{i} \end{cases}$$

Proposition 7.2.1. Let $\{\frac{\partial}{\partial z^i}, \frac{\partial}{\partial \bar{z}^i}\}$ be a local frame of $T_{\mathbb{C}}X$. Then

$$\tilde{\nabla}_{\frac{\partial}{\partial z^A}} \frac{\partial}{\partial z^B} = \Gamma_{AB}^C \frac{\partial}{\partial z^C} = \frac{1}{2} h^{CE} \left(\frac{\partial h_{EB}}{\partial z^A} + \frac{\partial h_{AE}}{\partial z^B} - \frac{\partial h_{AB}}{\partial z^E} \right) \frac{\partial}{\partial z^C}$$

where $A, B, C, E \in \{1, \dots, n, \bar{1}, \dots, \bar{n}\}$.

Proof. By proof of Koszul formula, it suffices to check

- (1) $\tilde{\nabla}$ is compatible with $g_{\mathbb{C}}$.
- (2) $\tilde{\nabla}$ is torsion-free, that is for $X, Y \in C^\infty(X, T_{\mathbb{C}}X)$, one has

$$\tilde{\nabla}_X Y - \tilde{\nabla}_Y X = [X, Y]$$

Above two claims can be checked easily by using the fact ∇ is Levi-Civita connection and the \mathbb{C} -linearity of $\tilde{\nabla}$. \square

Corollary 7.2.1.

$$\begin{aligned}\overline{\Gamma_{ij}^k} &= \Gamma_{ij}^k = \frac{1}{2}h^{k\bar{l}}\left(\frac{\partial h_{j\bar{l}}}{\partial z^i} + \frac{\partial h_{i\bar{l}}}{\partial z^j}\right) \\ \overline{\Gamma_{i\bar{j}}^k} &= \Gamma_{i\bar{j}}^k = \frac{1}{2}h^{k\bar{l}}\left(\frac{\partial h_{j\bar{l}}}{\partial \bar{z}^i} - \frac{\partial h_{j\bar{i}}}{\partial \bar{z}^l}\right) \\ \overline{\Gamma_{ij}^k} &= \Gamma_{ij}^k = 0\end{aligned}$$

Proof. Note that

$$h_{ij} = h_{\bar{i}\bar{j}} = h^{ij} = h^{\bar{i}\bar{j}} = 0$$

□

Theorem 7.2.1. Let (X, h) be a Kähler manifold with induced Riemannian metric g on underlying real manifold, and suppose ∇ is Levi-Civita connection with respect to g . Then Chern connection with respect to h can be obtained from the restriction of \mathbb{C} -linear extension of ∇ to $T^{1,0}X$.

Proof. Let $\tilde{\nabla}$ be the \mathbb{C} -linear extension of ∇ and $\{\frac{\partial}{\partial z^i}\}$ be a local frame of $T^{1,0}X$. Then by definition one has

$$\tilde{\nabla} \frac{\partial}{\partial z^j} = \Gamma_{ij}^k dz^i \otimes \frac{\partial}{\partial z^k} + \Gamma_{i\bar{j}}^{\bar{k}} dz^i \otimes \frac{\partial}{\partial \bar{z}^k} + \Gamma_{i\bar{j}}^k d\bar{z}^i \otimes \frac{\partial}{\partial z^k} + \Gamma_{i\bar{j}}^{\bar{k}} d\bar{z}^i \otimes \frac{\partial}{\partial \bar{z}^k}$$

By Corollary 7.2.1 one has $\Gamma_{i\bar{j}}^{\bar{k}} = 0$ automatically, and if Kähler condition holds, then

$$\begin{aligned}\Gamma_{ij}^k &= h^{k\bar{l}} \frac{\partial h_{j\bar{l}}}{\partial z^i} \\ \Gamma_{i\bar{j}}^k &= \Gamma_{i\bar{j}}^{\bar{k}} = 0\end{aligned}$$

Thus $\tilde{\nabla}|_{T^{1,0}X}$ gives a connection on $T^{1,0}X$, and by formula of Chern connection, it's exactly the Chern connection with respect to h . □

7.3. Curvatures of Kähler metric. In this section, let (X, h) be a Kähler manifold with induced Riemannian metric g on underlying real manifold, and suppose $\{z^i = x^i + \sqrt{-1}x^I\}$ is a local coordinate of X , where $1 \leq i \leq n$ and $n+1 \leq I = i+n \leq 2n$.

Notation 7.3.1.

(1)

$$\begin{aligned}R_{ijkl} &= R\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}, \frac{\partial}{\partial x^k}, \frac{\partial}{\partial x^l}\right) \\ R_{ijKL} &= R\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}, \frac{\partial}{\partial x^K}, \frac{\partial}{\partial x^L}\right)\end{aligned}$$

where R is curvature tensor of Levi-Civita connection ∇ with respect to g .

(2)

$$\Theta_{i\bar{j}k\bar{l}} = \Theta\left(\frac{\partial}{\partial z^i}, \frac{\partial}{\partial \bar{z}^j}, \frac{\partial}{\partial z^k}, \frac{\partial}{\partial \bar{z}^l}\right)$$

where Θ is Chern curvature with respect to h .

7.3.1. Ricci curvature and scalar curvature.

Lemma 7.3.1.

$$\begin{aligned} R_{ijkl} &= R_{ijKL} \\ R_{ijKl} &= -R_{ijkL} \end{aligned}$$

Proof. It follows from Kähler condition. \square **Corollary 7.3.1.**

$$\begin{aligned} R_{ij} &= R_{IJ} \\ R_{iJ} &= R_{Ji} = -R_{jI} = -R_{Ij} \end{aligned}$$

Proof. It follows from Proposition 6.1.1 and Lemma 7.3.1. \square **Theorem 7.3.1.**

$$\Theta_{i\bar{j}} = \frac{1}{2}(R_{ij} + \sqrt{-1}R_{iJ})$$

where $\Theta_{i\bar{j}}$ is given by $\text{Ric}(h) = \sqrt{-1}\Theta_{i\bar{j}}dz^i \wedge d\bar{z}^j$, while R_{ij} and R_{iJ} are Riemannian Ricci curvatures.

Proof. Suppose $\{\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^I}\}$ is an orthonormal frame of real tangent bundle with respect to g , and thus $\{u_i := \frac{1}{\sqrt{2}}(\frac{\partial}{\partial x^i} - \sqrt{-1}\frac{\partial}{\partial x^I})\}$ gives an orthonormal frame of holomorphic tangent bundle with respect to h . Then

$$\begin{aligned} \Theta_{i\bar{j}} &= \sum_k \left(R\left(\frac{\partial}{\partial z^i}, \frac{\partial}{\partial \bar{z}^j}, u_k, \bar{u}_k\right) \right) \\ &= \sum_k \frac{1}{2} \left(\underbrace{R\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}, u_k, \bar{u}_k\right)}_{\text{part I}} + \underbrace{\sqrt{-1}R\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^J}, u_k, \bar{u}_k\right)}_{\text{part II}} \right) \end{aligned}$$

For part I, one has

$$\begin{aligned} \sum_k R\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}, u_k, \bar{u}_k\right) &= \sum_k \frac{1}{2} R\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}, \frac{\partial}{\partial x^k} - \sqrt{-1}\frac{\partial}{\partial x^K}, \frac{\partial}{\partial x^k} + \sqrt{-1}\frac{\partial}{\partial x^K}\right) \\ &= \sum_k \frac{\sqrt{-1}}{2} (R_{ijkK} - R_{ijKk}) \\ &= \sum_k \sqrt{-1}R_{ijkK} \\ &= \sum_k \sqrt{-1}(-R_{kijK} - R_{jkiK}) \\ &= \sum_k \sqrt{-1}(R_{kiJk} + R_{KijK}) \\ &= \sqrt{-1}R_{iJ} \end{aligned}$$

Here we used Lemma 7.3.1 and first Bianchi identity. Similarly for part II one has

$$\begin{aligned}
\sum_k \sqrt{-1} R\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^J}, u_k, \bar{u}_k\right) &= \sum_k \frac{\sqrt{-1}}{2} R\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^J}, \frac{\partial}{\partial x^k} - \sqrt{-1} \frac{\partial}{\partial x^K}, \frac{\partial}{\partial x^k} + \sqrt{-1} \frac{\partial}{\partial x^K}\right) \\
&= \sum_k -R_{iJkK} \\
&= \sum_k (R_{kiJK} + R_{JkiK}) \\
&= \sum_k (R_{kijK} + R_{KijK}) \\
&= R_{ij}
\end{aligned}$$

This shows the desired result. \square

Corollary 7.3.2. Let (X, h) be a Kähler manifold with induced Riemannian metric g on underlying real manifold. Then

- (1) h has positive Chern-Ricci curvature if and only if g has positive Ricci curvature.
- (2) h is Kähler-Einstein with Einstein constant λ if and only if g is an Einstein metric with Einstein constant λ .

Corollary 7.3.3. Let (X, h) be a Kähler manifold with induced Riemannian metric g on underlying real manifold. Let s_R be the Riemannian scalar curvature and s is the Chern scalar curvature. Then $s_R = 2s$.

Proof. On one hand, direct computation shows

$$\begin{aligned}
s &= h^{i\bar{j}} \Theta_{i\bar{j}} \\
&= 2(g^{ij} - \sqrt{-1}g^{iJ}) \cdot \frac{1}{2}(R_{ij} + \sqrt{-1}R_{iJ}) \\
&= g^{ij} R_{ij} + g^{iJ} R_{iJ}
\end{aligned}$$

On the other hand,

$$\begin{aligned}
s_R &= g^{ij} R_{ij} + g^{iJ} R_{iJ} + g^{Ij} R_{Ij} + g^{IJ} R_{IJ} \\
&\stackrel{(1)}{=} 2g^{ij} R_{ij} + 2g^{iJ} R_{iJ} \\
&= 2s
\end{aligned}$$

where (1) holds from Proposition 6.1.1 and Corollary 7.3.1. \square

7.3.2. Holomorphic sectional curvature and holomorphic bisectional curvature. Recall that for a Hermitian manifold (X, h) and unit vectors $v = v^i \frac{\partial}{\partial z^i}$, $w = w^i \frac{\partial}{\partial z^i} \in T_p X$, the holomorphic sectional curvature is defined by

$$\text{HSC}_p(v) = \Theta_{i\bar{j}k\bar{l}} v^i \bar{v}^j v^k \bar{v}^l$$

and the holomorphic bisectonal curvature is defined by

$$\text{HBSC}_p(v, w) = \Theta_{i\bar{j}k\bar{l}} v^i \bar{v}^j w^k \bar{w}^l$$

Now suppose (X, h) is a Kähler manifold, we will see the holomorphic sectional curvature and holomorphic bisectonal curvature are closely related to the Riemannian sectional curvature.

Suppose unit vectors $v, w \in T_p X$ are given by

$$\begin{aligned} v &= \frac{1}{\sqrt{2}}(x - \sqrt{-1}Jx) \\ w &= \frac{1}{\sqrt{2}}(y - \sqrt{-1}Jy) \end{aligned}$$

where x, y are real vectors with $g(x, x) = g(y, y) = 1$. Then the holomorphic bisectonal curvature is computed by

$$\begin{aligned} R(v, \bar{v}, w, \bar{w}) &= \frac{1}{2}R(x - \sqrt{-1}Jx, x + \sqrt{-1}Jx, w, \bar{w}) \\ &= \sqrt{-1}R(x, Jx, w, \bar{w}) \\ &= -R(x, Jx, y, Jy) \\ &= R(x, y, y, x) + R(x, Jy, Jy, x) \end{aligned}$$

In particular, the holomorphic sectional curvature of v is exactly the sectional curvature of the plane spanned by x and Jx .

Note that the holomorphic bisectonal curvature is a sum of two sectional curvatures. Hence the holomorphic bisectonal curvature carries less information than the sectional curvature. On the other hand, the holomorphic bisectonal curvature carries more information than the Ricci curvature, since for real vector x , one has

$$\text{Ric}_p(x) = \sum_{i=1}^n \left(R\left(\frac{\partial}{\partial x^i}, x, x, \frac{\partial}{\partial x^i}\right) + R\left(J\left(\frac{\partial}{\partial x^i}\right), x, x, J\left(\frac{\partial}{\partial x^i}\right)\right) \right)$$

As a consequence, positive sectional curvature implies positive holomorphic bisectonal curvature, and positive holomorphic bisectonal curvature implies positive Ricci curvature.

Definition 7.3.1 (constant holomorphic sectional curvature). Let (X, h) be a Kähler manifold. (X, h) has constant holomorphic sectional curvature c if

$$\text{HSC}_p(v) = c$$

for all unit vector $v \in T_p X$.

Remark 7.3.1. In other words, the sectional curvature of all J -invariant planes equal to c , that is, $R(x, Jx, Jx, x) = c$ for all unit real vector x .

The definition given above is the most natural way to define constant holomorphic sectional curvature, and now let's try to give another descriptions which is easy to use.

Proposition 7.3.1. Let (X, h) be a Kähler manifold with constant holomorphic sectional curvature c . Then $R = cR_0$, where

$$R_0(X, Y, Z, W) = \frac{1}{4} \{g(X, W)g(Y, Z) - g(Y, W)g(X, Z) - g(X, JZ)g(Y, JW) \\ + g(X, JW)g(Y, JZ) - 2g(X, JY)g(Z, JW)\}$$

Proof. See Proposition 7.3 of Chapter IX in [KN69]. \square

Proposition 7.3.2. Let (X, h) be a Kähler manifold. It has constant holomorphic sectional curvature c if and only if

$$\Theta_{i\bar{j}k\bar{l}} = \frac{c}{2}(h_{i\bar{j}}h_{k\bar{l}} + h_{i\bar{l}}h_{k\bar{j}})$$

Proof. A direct computation shows

$$\begin{aligned} \Theta_{i\bar{j}k\bar{l}} &= R\left(\frac{\partial}{\partial z^i}, \frac{\partial}{\partial \bar{z}^j}, \frac{\partial}{\partial z^k}, \frac{\partial}{\partial \bar{z}^l}\right) \\ &= cR_0\left(\frac{\partial}{\partial z^i}, \frac{\partial}{\partial \bar{z}^j}, \frac{\partial}{\partial z^k}, \frac{\partial}{\partial \bar{z}^l}\right) \\ &= \frac{c}{4} \left\{ g\left(\frac{\partial}{\partial z^i}, \frac{\partial}{\partial \bar{z}^l}\right)g\left(\frac{\partial}{\partial \bar{z}^j}, \frac{\partial}{\partial z^k}\right) - g\left(\frac{\partial}{\partial \bar{z}^j}, \frac{\partial}{\partial \bar{z}^l}\right)g\left(\frac{\partial}{\partial z^i}, \frac{\partial}{\partial z^k}\right) - g\left(\frac{\partial}{\partial z^i}, J\frac{\partial}{\partial z^k}\right)g\left(\frac{\partial}{\partial \bar{z}^j}, J\frac{\partial}{\partial \bar{z}^l}\right) \right. \\ &\quad \left. + g\left(\frac{\partial}{\partial z^i}, J\frac{\partial}{\partial \bar{z}^l}\right)g\left(\frac{\partial}{\partial \bar{z}^j}, J\frac{\partial}{\partial z^k}\right) - 2g\left(\frac{\partial}{\partial z^i}, J\frac{\partial}{\partial \bar{z}^j}\right)g\left(\frac{\partial}{\partial z^k}, J\frac{\partial}{\partial \bar{z}^l}\right) \right\} \\ &= \frac{c}{4}(h_{i\bar{l}}h_{k\bar{j}} + h_{i\bar{j}}h_{k\bar{l}} + 2h_{i\bar{j}}h_{k\bar{l}}) \\ &= \frac{c}{2}(h_{i\bar{l}}h_{k\bar{j}} + h_{i\bar{j}}h_{k\bar{l}}) \end{aligned}$$

On the other hand, it's clear (X, h) has constant holomorphic sectional curvature c if $\Theta_{i\bar{j}k\bar{l}} = \frac{c}{2}(h_{i\bar{j}}h_{k\bar{l}} + h_{i\bar{l}}h_{k\bar{j}})$. \square

Remark 7.3.2.

- (1) Above formula differs a sign from the one given in Proposition 7.6 of Chapter IX in [KN69], since the curvature notation we defined here differs a sign from the one defined by Kobayashi.
- (2) In [Tia00], he called this by constant holomorphic bisectional curvature, and it maybe a bit confusing for beginners. If (X, h) has constant holomorphic sectional curvature, in this case you have that the bisectional curvature of X, Y equals

$$R(X, JX, JY, Y) = \frac{c}{2}(1 + g(X, Y)^2 + g(X, JY))$$

As you can see, this expression is not constant (as you vary X, Y among unit vectors), but it varies between $c/2$ and c according to the relative position of the 2-planes spanned by (X, JX) and (Y, JY) . In particular you see that there is no such thing as a non-flat ‘‘Kähler manifold with constant holomorphic bisectional curvature’’, because if there was such a thing, in particular the holomorphic sectional curvature would be constant, but then the curvature tensor would be given by the above

formula, and the bisectional curvature would actually be NOT constant unless $c = 0$, that is, all the curvatures are constant because the metric is flat.

7.4. Fubini-Study metric.

Proposition 7.4.1 (Fubini-Study metric). Let $\mathbb{C}\mathbb{P}^n = \bigcup_{i=0}^n U_i$ be the canonical open covering, that is $U_i = \{(z^0 : \cdots : z^n) \mid z^i \neq 0\}$. Then there is a Kähler metric ω_{FS} on $\mathbb{C}\mathbb{P}^n$, called Fubini-Study metric, such that

$$\omega_{FS}|_{U_i} = \frac{\sqrt{-1}}{2} \partial \bar{\partial} \log \left(\frac{\sum_{l=0}^n |z^l|^2}{|z^i|^2} \right)$$

Proof. Note that $\omega_{FS}|_{U_i}$ can be written as

$$\omega_{FS}|_{U_i} = \frac{\sqrt{-1}}{2} \partial \bar{\partial} \log \left(\sum_{l=1}^n |w^l|^2 + 1 \right)$$

where $w^l = z^l/z^i$. The following steps show that $\{\omega_{FS}|_{U_i}\}_{i=0}^n$ gives a real $(1, 1)$ -form which is closed.

(1) It's globally defined since direct computation shows

$$\begin{aligned} \log \left(\frac{\sum_{k=0}^n |z^k|^2}{|z^i|^2} \right) &= \log \left(\frac{|z^j|^2}{|z^i|^2} \frac{\sum_{k=0}^n |z^k|^2}{|z^j|^2} \right) \\ &= \log \left(\frac{|z^j|^2}{|z^i|^2} \right) + \log \left(\frac{\sum_{k=0}^n |z^k|^2}{|z^j|^2} \right) \\ &= \log \left(\frac{\sum_{k=0}^n |z^k|^2}{|z^j|^2} \right) \end{aligned}$$

where the last equality holds since $|z^j|^2/|z^i|^2$ is a nowhere vanishing holomorphic function.

(2) It's real since $\bar{\partial} \bar{\partial} = \bar{\partial} \partial = -\partial \bar{\partial}$.

(3) It's ∂ -closed since each $\omega_{FS}|_{U_i}$ is ∂ -closed.

It remains to show ω is positive. A direct computation yields

$$\partial \bar{\partial} \log \left(1 + \sum_{l=1}^n |w^l|^2 \right) = \frac{1}{(1 + \sum_{l=1}^n |w^l|^2)^2} h_{i\bar{j}} dw^i \wedge d\bar{w}^j$$

where $h_{i\bar{j}} = (1 + \sum_{l=1}^n |w^l|^2) \delta_{ij} - w^i \bar{w}^j$. Now it suffices to show $h_{i\bar{j}}$ is positive definite, for $u \neq 0$, one has

$$\begin{aligned} u^T (h_{i\bar{j}}) \bar{u} &= (u, u) + (w, w)(u, u) - u^T \bar{w} w^T \bar{u} \\ &= (u, u) + (w, w)(u, u) - (u, w)(w, u) \\ &= (u, u) + (w, w)(u, u) - \overline{(w, u)}(w, u) \\ &= (u, u) + (w, w)(u, u) - |(w, u)|^2 > 0 \end{aligned}$$

□

Corollary 7.4.1. Any projective manifold is Kähler.

Proof. By Proposition 7.1.2, the submanifold of Kähler manifold is still Kähler. \square

Proposition 7.4.2. $\mathcal{O}_{\mathbb{C}\mathbb{P}^n}(1)$ is a positive holomorphic line bundle.

Proof. Note that line bundle $\mathcal{O}_{\mathbb{C}\mathbb{P}^n}(1)$ can be given transition functions $\{U_i, g_{ij}\}$, where $\{U_i\}$ is canonical open covering and $g_{ij} = z^j/z^i$. Consider $h_i: U_i \rightarrow \mathbb{R}_{>0}$ given by

$$h_i = \frac{|z^i|^2}{\sum_{l=0}^n |z^l|^2}$$

Then h_i can be glued together to obtain a Hermitian metric on $\mathcal{O}_{\mathbb{C}\mathbb{P}^n}(1)$ since $h_i = h_j |g_{ij}|^2$. And it's clear to see Hermitian metric corresponding to curvature of Chern connection with respect to this metric is Fubini-Study metric. \square

Remark 7.4.1. Note that $\mathcal{O}_{\mathbb{C}\mathbb{P}^n}(-1)$ is a subbundle of $\mathbb{C}\mathbb{P}^n \times \mathbb{C}^{n+1}$, so we can obtain a natural Hermitian metric of $\mathcal{O}_{\mathbb{C}\mathbb{P}^n}(-1)$ by restricting standard Hermitian metric of $\mathbb{C}\mathbb{P}^n \times \mathbb{C}^{n+1}$, and Hermitian metric on $\mathcal{O}_{\mathbb{C}\mathbb{P}^n}(1)$ we defined before is exactly the dual metric of this natural metric.

Theorem 7.4.1. The Fubini-Study metric ω_{FS} on $\mathbb{C}\mathbb{P}^n$ is a Kähler-Einstein metric with Einstein constant $n+1$ and has constant holomorphic sectional curvature.

Remark 7.4.2.

- (1) In our definition (or the one given by Kobayashi in [KN69]), Fubini-Study metric has constant holomorphic sectional curvature 2, and in the definition given by Gang Tian in [Tia00], Fubini-Study metric has constant holomorphic sectional curvature 1, but it doesn't matter since up to a rescaling they're all the same.
- (2) There are other models that have constant holomorphic sectional curvature, such as

Example 7.4.1. Let $X = \mathbb{C}^n$ equipped with $\omega = \sqrt{-1}/2 dz^i \wedge d\bar{z}^i$. Then (X, ω) is flat, and thus has constant holomorphic sectional curvature 0.

Example 7.4.2. Let $X = \mathbb{B}^n = \{z \in \mathbb{C}^n : |z| < 1\}$ equipped with

$$\omega = \frac{\sqrt{-1}}{2} \partial\bar{\partial} \log(1 - |z|^2)$$

Then (X, ω) has constant holomorphic sectional curvature -2 .

Also, there is a theorem parallel to the Hopf's theorem in Riemannian geometry.

Theorem 7.4.2 (uniformization theorem). If (X, h) is a complete Kähler manifold of constant holomorphic sectional curvature, then its universal covering is one of above examples. Moreover, up to rescaling, h pulls back to one of the metrics in the above examples.

Part 3. Hodge theory

8. HODGE THEOREM

8.1. Hodge star and adjoint operators. Let (X, ω) be a compact Hermitian manifold. A (p, q) -form α can be locally written as

$$\alpha = \frac{1}{p! \times q!} \alpha_{i_1 \dots i_p \bar{j}_1 \dots \bar{j}_q} dz^{i_1} \wedge \dots \wedge dz^{i_p} \wedge d\bar{z}^{j_1} \wedge \dots \wedge d\bar{z}^{j_q}$$

Then for $\alpha, \beta \in C^\infty(X, \Omega_X^{p,q})$, the local inner product is defined as

$$\langle \alpha, \beta \rangle = \frac{1}{p! \times q!} h^{i_1 \bar{k}_1} \dots h^{i_p \bar{k}_p} h^{l_1 \bar{j}_1} \dots h^{l_q \bar{j}_q} \alpha_{i_1 \dots i_p \bar{j}_1 \dots \bar{j}_q} \overline{\beta_{k_1 \dots k_p \bar{l}_1 \dots \bar{l}_q}}$$

which is a smooth function on X .

Definition 8.1.1 (inner product on (p, q) -form). An inner product on the space of (p, q) -form is defined as

$$(\alpha, \beta) := \int_X \langle \alpha, \beta \rangle \frac{\omega^n}{n!}$$

where $\alpha, \beta \in C^\infty(X, \Omega_X^{p,q})$. This also gives an inner product on $\Omega_{X, \mathbb{C}}^k = \bigoplus_{p+q=k} \Omega_X^{p,q}$.

Holding the inner product $(-, -)$, the formal adjoint operator of d is defined as an operator

$$d^*: C^\infty(X, \Omega_{X, \mathbb{C}}^k) \rightarrow C^\infty(X, \Omega_{X, \mathbb{C}}^{k-1})$$

satisfying $(\alpha, d\beta) = (d^*\alpha, \beta)$ for α, β with appropriate degrees, similarly one can define ∂^* and $\bar{\partial}^*$. In order to construct these adjoint operators, we need to introduce the well-known Hodge star operator.

Definition 8.1.2 (Hodge star operator). There exists an operator

$$\star : C^\infty(X, \Omega_X^{p,q}) \rightarrow C^\infty(X, \Omega_X^{n-q, n-p})$$

such that

$$(\alpha, \beta) = \int_X \alpha \wedge \star \bar{\beta}$$

Remark 8.1.1. It's well-defined since $\bar{\beta}$ is a (q, p) -form, and thus $\star \bar{\beta}$ is a $(n-p, n-q)$ -form.

Lemma 8.1.1.

- (1) $\star 1 = \omega^n / n!$
- (2) $\star \omega = \omega^{n-1} / (n-1)!$
- (3) $\star \bar{\psi} = \overline{\star \psi}$
- (4) $\star \star = (-1)^{p+q}$ on $C^\infty(X, \Omega_X^{p,q})$
- (5) $(\star \varphi, \star \psi) = (\varphi, \psi)$

Proposition 8.1.1. $d^* = -\star d\star$

Proof. For arbitrary $\alpha \in C^\infty(X, \Omega_{X, \mathbb{C}}^{p+q})$ and $\beta \in C^\infty(X, \Omega_{X, \mathbb{C}}^{p+q+1})$, then

$$\begin{aligned}
(d\alpha, \beta) &= \int_X d\alpha \wedge \star \beta \\
&= \int_X d(\alpha \wedge \star \beta) - (-1)^{p+q} \alpha \wedge d \star \beta \\
&= (-1)^{p+q+1} \int_X \alpha \wedge d \star \beta \\
&\stackrel{(1)}{=} (-1)^{p+q+1} (-1)^{2n-(p+q+1)+1} \int_X \alpha \wedge \star \star d \star \beta \\
&= -(\alpha, \star d \star \beta)
\end{aligned}$$

where (1) holds from (4) of Lemma 8.1.1. \square

Proposition 8.1.2.

$$\begin{aligned}
\partial^* &= -\star \bar{\partial} \star \\
\bar{\partial}^* &= -\star \partial \star
\end{aligned}$$

Proof. Direct computation. \square

Definition 8.1.3 (Lefschetz operator). Let (X, ω) be a compact Kähler manifold. The Lefschetz operator is defined as

$$\begin{aligned}
L: C^\infty(X, \Omega_X^{p,q}) &\rightarrow C^\infty(X, \Omega_X^{p+1,q+1}) \\
\alpha &\mapsto \omega \wedge \alpha
\end{aligned}$$

Lemma 8.1.2. $\Lambda := L^* = (-1)^{p+q} \star L \star$ on (p, q) -forms.

Proof. For $\alpha \in C^\infty(X, \Omega_X^{p,q}), \beta \in C^\infty(X, \Omega_X^{p+1,q+1})$, direct computation shows

$$\begin{aligned}
(L\alpha, \beta) &= \int_X L\alpha \wedge \star \beta \\
&= \int_X \omega \wedge \alpha \wedge \star \beta \\
&\stackrel{(1)}{=} \int_X \alpha \wedge \omega \wedge \star \beta \\
&\stackrel{(2)}{=} \int_X \alpha \wedge (-1)^{p+q} \star \star \omega \wedge \star \beta \\
&= (\alpha, (-1)^{p+q} \star L \star \beta)
\end{aligned}$$

where

- (1) holds from ω is a 2-form.
- (2) holds from (4) of Lemma 8.1.1. \square

8.1.1. *Useful formulas of adjoint operators.*

Proposition 8.1.3. Let (X, h) be a compact Kähler manifold. Then locally

$$\begin{cases} \partial = dz^i \wedge \nabla_i \\ \partial^* = -h^{i\bar{j}} \iota_i \circ \nabla_j = -h^{i\bar{j}} \nabla_j \circ \iota_i \end{cases} \quad \begin{cases} \bar{\partial} = d\bar{z}^i \wedge \nabla_{\bar{i}} \\ \bar{\partial}^* = -h^{i\bar{j}} \iota_{\bar{j}} \circ \nabla_i = -h^{i\bar{j}} \nabla_i \circ \iota_{\bar{j}} \end{cases}$$

Proof. Here we only give the proof of the case ∂ and ∂^* , the proof for the other two cases are same. It suffices to check pointwisely, and at each point we may also choose normal coordinate in Theorem 7.1.1. For (p, q) -form α , locally written as $\alpha = \alpha_{J\bar{K}} dz^J \wedge d\bar{z}^K$. Then

$$\partial\alpha = \frac{\partial\alpha_{J\bar{K}}}{\partial z^i} dz^i \wedge dz^J \wedge d\bar{z}^K$$

and

$$\begin{aligned} dz^i \wedge \nabla_i \alpha &= dz^i \wedge \nabla_i (\alpha_{J\bar{K}} dz^J \wedge d\bar{z}^K) \\ &= dz^i \wedge \frac{\partial\alpha_{J\bar{K}}}{\partial z^i} dz^J \wedge d\bar{z}^K + \alpha_{J\bar{K}} \nabla_i (dz^J \wedge d\bar{z}^K) \\ &\stackrel{(1)}{=} \frac{\partial\alpha_{J\bar{K}}}{\partial z^i} dz^i \wedge dz^J \wedge d\bar{z}^K \end{aligned}$$

where (1) holds from our choice of normal coordinate. To see formula of ∂^* , take arbitrary forms α, β with appropriate bidegrees, then

$$\begin{aligned} (\partial\alpha, \beta) &= (dz^i \wedge \nabla_i \alpha, \beta) \\ &\stackrel{(2)}{=} (\nabla_i \alpha, h^{p\bar{i}} \iota_p \beta) \\ &\stackrel{(3)}{=} -(\alpha, h^{p\bar{i}} \nabla_i \circ \iota_p \beta) \end{aligned}$$

where

(2) holds from Proposition 6.3.4.

(3) holds from Stokes' theorem and the fact Chern connection is compatible with metric.

This shows

$$\partial^* = -h^{i\bar{j}} \nabla_j \circ \iota_i \stackrel{(4)}{=} -h^{i\bar{j}} \iota_i \circ \nabla_j$$

where (4) holds from $\iota_i \circ \nabla_j = \nabla_j \circ \iota_i$. \square

Proposition 8.1.4. Let (X, ω) be a compact Kähler manifold. Then locally

$$\Lambda = \sqrt{-1} h^{i\bar{j}} \iota_i \circ \iota_{\bar{j}} = -\sqrt{-1} h^{i\bar{j}} \iota_{\bar{j}} \circ \iota_i$$

Proof. For arbitrary forms α, β with appropriate bidegrees, direct computation shows

$$\begin{aligned}
(\omega \wedge \alpha, \beta) &= (\sqrt{-1}h_{i\bar{j}}dz^i \wedge d\bar{z}^j \wedge \alpha, \beta) \\
&\stackrel{(1)}{=} (\sqrt{-1}h_{i\bar{j}}d\bar{z}^j \wedge \alpha, h^{p\bar{i}}\iota_p\beta) \\
&\stackrel{(2)}{=} (\sqrt{-1}h_{i\bar{j}}\alpha, h^{p\bar{i}}h^{j\bar{q}}\iota_{\bar{q}} \circ \iota_p\beta) \\
&\stackrel{(3)}{=} (\alpha, -\sqrt{-1}h_{j\bar{i}}h^{p\bar{i}}h^{j\bar{q}}\iota_{\bar{q}} \circ \iota_p\beta) \\
&= (\alpha, -\sqrt{-1}h^{p\bar{i}}\iota_{\bar{i}} \circ \iota_p\beta)
\end{aligned}$$

where

(1) and (2) hold from Proposition 6.3.4.

(3) holds from $h_{i\bar{j}}$ is Hermitian, that is $\bar{h}_{i\bar{j}} = h_{j\bar{i}}$.

This shows

$$\Lambda = -\sqrt{-1}h^{i\bar{j}}\iota_{\bar{j}} \circ \iota_i \stackrel{(4)}{=} \sqrt{-1}h^{i\bar{j}}\iota_i \circ \iota_{\bar{j}}$$

where (4) holds from $\iota_i \circ \iota_{\bar{j}} = -\iota_{\bar{j}} \circ \iota_i$. \square

8.2. Hodge theorem.

Definition 8.2.1 (Laplacian). Laplacian Δ_\bullet is an operator defined by $\Delta_\bullet := \bullet\bullet^* + \bullet^*\bullet$, where \bullet can be d, ∂ and $\bar{\partial}$.

Definition 8.2.2 (harmonic). A form α is called Δ_\bullet -harmonic if $\Delta_\bullet\alpha = 0$, where \bullet can be d, ∂ and $\bar{\partial}$.

Notation 8.2.1. \mathcal{H}^k denotes the space of Δ_d -harmonic k -forms, and $\mathcal{H}^{p,q}$ denotes the space of $\Delta_{\bar{\partial}}$ -harmonic forms of type (p, q) .

Lemma 8.2.1. α is Δ_\bullet -harmonic if and only if $\bullet\alpha = 0, \bullet^*\alpha = 0$, where \bullet can be d, ∂ and $\bar{\partial}$.

Proof. Direct computation shows

$$\begin{aligned}
(\alpha, \Delta_d\alpha) &= (\alpha, dd^*\alpha) + (\alpha, d^*d\alpha) \\
&= \|d^*\alpha\|^2 + \|d\alpha\|^2
\end{aligned}$$

This shows α is Δ_d -harmonic if and only if $d\alpha = d^*\alpha = 0$, the other cases are same. \square

Theorem 8.2.1 (Hodge theorem). Let (X, h) be a compact Hermitian n -manifold. Then

- (1) $\mathcal{H}^{p,q}$ is finite dimensional.
- (2) There is a decomposition $C^\infty(X, \Omega_X^{p,q}) = \mathcal{H}^{p,q} \oplus \Delta_{\bar{\partial}}(C^\infty(X, \Omega_X^{p,q}))$, which is orthogonal with respect to inner products in Definition 8.1.1.

Corollary 8.2.1. There is the following orthonormal decomposition

$$C^\infty(X, \Omega_X^{p,q}) = \mathcal{H}^{p,q} \oplus \bar{\partial}(C^\infty(X, \Omega_X^{p,q-1})) \oplus \bar{\partial}^*(C^\infty(X, \Omega_X^{p,q+1}))$$

Corollary 8.2.2.

$$\begin{aligned}\ker \bar{\partial} &= \mathcal{H}^{p,q} \oplus \bar{\partial}^*(C^\infty(X, \Omega_X^{p,q-1})) \\ \ker \bar{\partial}^* &= \mathcal{H}^{p,q} \oplus \bar{\partial}(C^\infty(X, \Omega_X^{p,q+1}))\end{aligned}$$

Corollary 8.2.3. The natural map $\mathcal{H}^{p,q} \rightarrow H^{p,q}(X)$ is an isomorphism. In particular, $H^{p,q}(X)$ is finite dimensional.

In order to give the following isomorphism

$$\star: \mathcal{H}^{p,q} \rightarrow \mathcal{H}^{n-q,n-p}$$

Parallel to the real case¹⁰, it suffices to have

$$\star \circ \Delta_{\bar{\partial}} = \Delta_{\bar{\partial}} \circ \star$$

But something bad happens since we only have $\bar{\partial}^* = -\star \partial \star$, and direct computation only yields $\Delta_{\bar{\partial}} \circ \star = \star \circ \Delta_{\partial}$. So it fails generally since $\Delta_{\bar{\partial}} \neq \Delta_{\partial}$. There are two ways to deal with this gap. The first way is that we will see later if X is compact Kähler manifold, then $\Delta_{\partial} = \Delta_{\bar{\partial}}$, that is Theorem 9.1.1. Then

Corollary 8.2.4. If (X, ω) is a compact Kähler n -manifold, then $\star: \mathcal{H}^{p,q} \rightarrow \mathcal{H}^{n-q,n-p}$ is an isomorphism.

Another way is to consider

$$\begin{aligned}\bar{\star}: C^\infty(X, \Omega_X^{p,q}) &\rightarrow C^\infty(X, \Omega_X^{n-p,n-q}) \\ \alpha &\mapsto \star \bar{\alpha}\end{aligned}$$

then direct computation shows

$$\bar{\star} \circ \Delta_{\bar{\partial}} = \Delta_{\bar{\partial}} \circ \bar{\star}$$

Corollary 8.2.5. If (X, h) is a compact Hermitian manifold, then $\bar{\star}: \mathcal{H}^{p,q} \rightarrow \mathcal{H}^{n-p,n-q}$ is an isomorphism.

Corollary 8.2.6. $H^{p,q}(X) \cong H^{n-p,n-q}(X)$.

Remark 8.2.1. This is a special case of Serre duality.

¹⁰See Hodge theory in [Liu23].

9. HODGE DECOMPOSITION

9.1. Kähler identities.

Definition 9.1.1 (commutator of differential operators). Let A, B be two differential operators. The commutator of A, B is defined as

$$[A, B] := AB - (-1)^{\deg A \deg B} BA$$

Lemma 9.1.1 (Jacobi identity). Let A, B, C be differential operators. Then $(-1)^{\deg A \deg C} [A, [B, C]] + (-1)^{\deg B \deg A} [B, [C, A]] + (-1)^{\deg C \deg B} [C, [A, B]] = 0$

Remark 9.1.1. In our case, the degree of $d, d^*, \partial, \partial^*, \bar{\partial}, \bar{\partial}^*$ is one, and the degree of L and Λ is zero¹¹.

Proposition 9.1.1 (Kähler identities). If (X, ω) is a compact Kähler manifold, then

$$\begin{aligned} [\bar{\partial}^*, L] &= \sqrt{-1} \partial \\ [\partial^*, L] &= -\sqrt{-1} \cdot \bar{\partial} \\ [\Lambda, \bar{\partial}] &= -\sqrt{-1} \partial^* \\ [\Lambda, \partial] &= \sqrt{-1} \cdot \bar{\partial}^* \end{aligned}$$

Proof. By taking conjugates and adjoints, it suffices to prove the first identity, which is a first order identity of differential equation. But by Theorem 7.1.1, locally we have $h_{i\bar{j}} = \delta_{i\bar{j}} + O(|\xi^2|)$. Thus it suffices to check Kähler identity for the case $U \subseteq \mathbb{C}^n$ equipped with standard Hermitian metric.

Suppose (p, q) -form α is locally given by $\alpha = \alpha_{JK} dz^J \wedge d\bar{z}^K$, then by Proposition 8.1.3 one has $\bar{\partial}^* \alpha = -\sum_l \iota_{\frac{\partial}{\partial \bar{z}^l}} \frac{\partial \alpha}{\partial z^l}$. Thus

$$\begin{aligned} [\bar{\partial}^*, L] \alpha &= \bar{\partial}^* (\omega \wedge \alpha) - \omega \wedge \bar{\partial}^* \alpha \\ &= -\sum_l \iota_{\frac{\partial}{\partial \bar{z}^l}} \frac{\partial}{\partial z^l} (\omega \wedge \alpha) + \omega \wedge \sum_l \iota_{\frac{\partial}{\partial \bar{z}^l}} \frac{\partial \alpha}{\partial z^l} \\ &\stackrel{(1)}{=} -\sum_l \iota_{\frac{\partial}{\partial \bar{z}^l}} (\omega \wedge \frac{\partial \alpha}{\partial z^l}) + \omega \wedge \sum_l \iota_{\frac{\partial}{\partial \bar{z}^l}} \frac{\partial \alpha}{\partial z^l} \\ &= -\left\{ \sum_l (\iota_{\frac{\partial}{\partial \bar{z}^l}} \omega) \wedge \frac{\partial \alpha}{\partial z^l} + \omega \wedge \sum_l \iota_{\frac{\partial}{\partial \bar{z}^l}} \frac{\partial \alpha}{\partial z^l} \right\} + \sum_l \omega \wedge \iota_{\frac{\partial}{\partial \bar{z}^l}} \frac{\partial \alpha}{\partial z^l} \\ &= -\sum_l (\iota_{\frac{\partial}{\partial \bar{z}^l}} \omega) \wedge \frac{\partial \alpha}{\partial z^l} \\ &\stackrel{(2)}{=} \sqrt{-1} \sum_l dz^l \wedge \frac{\partial \alpha}{\partial z^l} \\ &= \sqrt{-1} \partial \alpha \end{aligned}$$

¹¹You can try to understand this thing in a following way: operators $d, d^*, \partial, \partial^*, \bar{\partial}, \bar{\partial}^*$ take derivatives, but L and Λ are linear operators.

where

(1) holds from ω is a closed $(1, 1)$ -form.

(2) holds from Proposition 6.1.4, that is $\omega = \sqrt{-1} \sum_{i=1}^n dz^i \wedge d\bar{z}^i$.

□

Theorem 9.1.1. Let (X, ω) be a compact Kähler manifold. Then

$$\Delta_d = 2\Delta_\partial = 2\Delta_{\bar{\partial}}$$

Proof. Since

$$\Delta_d = (\partial + \bar{\partial})(\partial^* + \bar{\partial}^*) + (\partial^* + \bar{\partial}^*)(\partial + \bar{\partial})$$

By the fourth Kähler identity, one has

(1) The first term can be computed as

$$\begin{aligned} (\partial + \bar{\partial})(\partial^* + \bar{\partial}^*) &= (\partial + \bar{\partial})(\partial^* - \sqrt{-1}\Lambda\partial + \sqrt{-1}\partial\Lambda) \\ &= \partial\partial^* - \sqrt{-1}\partial\Lambda\partial + \bar{\partial}\partial^* - \sqrt{-1}\cdot\bar{\partial}\Lambda\partial + \sqrt{-1}\cdot\bar{\partial}\partial\Lambda \end{aligned}$$

(2) The second term can be computed as

$$\begin{aligned} (\partial^* + \bar{\partial}^*)(\partial + \bar{\partial}) &= (\partial^* - \sqrt{-1}\Lambda\partial + \sqrt{-1}\partial\Lambda)(\partial + \bar{\partial}) \\ &= \partial^*\partial + \sqrt{-1}\partial\Lambda\partial + \partial^*\bar{\partial} - \sqrt{-1}\Lambda\partial\bar{\partial} + \sqrt{-1}\partial\Lambda\bar{\partial} \end{aligned}$$

By the third Kähler identity, one has

$$\partial^* = \sqrt{-1}[\Lambda, \bar{\partial}] = \sqrt{-1}\Lambda\bar{\partial} - \sqrt{-1}\bar{\partial}\Lambda$$

then

$$\begin{aligned} \bar{\partial}\partial^* &= \bar{\partial}(\sqrt{-1}\Lambda\bar{\partial} - \sqrt{-1}\cdot\bar{\partial}\Lambda) = \sqrt{-1}\cdot\bar{\partial}\Lambda\bar{\partial} \\ \partial^*\bar{\partial} &= (\sqrt{-1}\Lambda\bar{\partial} - \sqrt{-1}\cdot\bar{\partial}\Lambda)\bar{\partial} = -\sqrt{-1}\cdot\bar{\partial}\Lambda\bar{\partial} = -\bar{\partial}\partial^* \end{aligned}$$

Now we have

$$\begin{aligned} \Delta_d &= \Delta_\partial - \sqrt{-1}\cdot\bar{\partial}\Lambda\partial - \sqrt{-1}\Lambda\partial\bar{\partial} + \sqrt{-1}\bar{\partial}\partial\Lambda + \sqrt{-1}\partial\Lambda\bar{\partial} \\ &= \Delta_\partial + \sqrt{-1}(\Lambda\bar{\partial}\partial - \bar{\partial}\Lambda\partial) + \sqrt{-1}(\partial\Lambda\bar{\partial} - \bar{\partial}\partial\Lambda) \\ &= \Delta_\partial + \sqrt{-1}[\Lambda, \bar{\partial}]\partial + \sqrt{-1}\partial[\Lambda, \bar{\partial}] \\ &= \Delta_\partial + \partial^*\partial + \partial\partial^* \\ &= 2\Delta_\partial \end{aligned}$$

□

Corollary 9.1.1. On a compact Kähler manifold, Δ_d -harmonic is equivalent to Δ_∂ -harmonic, and is equivalent to $\Delta_{\bar{\partial}}$ -harmonic.

Corollary 9.1.2. Let (X, ω) be a Kähler manifold and α be a (p, q) -form. Then $\Delta_d\alpha$ is still a (p, q) -form.

Proof. It's clear to see $\Delta_\partial\alpha$ is still a (p, q) -form.

□

Exercise 9.1.1. Show that for compact Kähler manifold we have

$$[\Delta_d, L] = 0$$

$$[L, \Lambda] = (k - n) \text{id} \quad \text{on } C^\infty(X, \Omega_{X, \mathbb{C}}^k)$$

Proof. For the first equation, we have $\Delta_d = 2\Delta_\partial = 2(\partial\partial^* + \partial^*\partial)$. Thus

$$[\Delta_d, L] = 2([\partial\partial^*, L] + [\partial^*\partial, L]) = 2(\partial[\partial^*, L] + [\partial^*, L]\partial)$$

The last equality holds by the fact that L commutes with ∂ since ω is ∂ -closed. Now we use the identity $[\partial^*, L] = -\sqrt{-1} \cdot \bar{\partial}$, which anticommutes with ∂ to conclude.

For the second equation, without lose of generality it suffices to check on $U \subseteq \mathbb{C}^n$ equipped with standard Hermitian metric since we are considering operators of order zero. Suppose $\varphi = \varphi_{IJ} dz^I d\bar{z}^J$ is a k -form with type (p, q) . A direct computation shows

$$\begin{aligned} L\Lambda\varphi &= L \left(\sqrt{-1} \sum_{i=1}^n \varphi_{IJ} \iota_i \circ \iota_{\bar{i}} (dz^I \wedge d\bar{z}^J) \right) \\ &= L \left(\sqrt{-1} \sum_{i=1}^n (-1)^p \varphi_{IJ} \iota_i dz^I \wedge \iota_{\bar{i}} d\bar{z}^J \right) \\ &= (\sqrt{-1})^2 \sum_{i,j=1}^n (-1)^{2p-1} \varphi_{IJ} dz^j \wedge \iota_i dz^I \wedge d\bar{z}^j \iota_{\bar{i}} d\bar{z}^J \\ \Lambda L\varphi &= \Lambda \left(\sqrt{-1} \sum_{j=1}^n (-1)^p \varphi_{IJ} dz^j \wedge dz^I \wedge d\bar{z}^j \wedge d\bar{z}^J \right) \\ &= (\sqrt{-1})^2 \sum_{i,j=1}^n (-1)^p \varphi_{IJ} \iota_i \circ \iota_{\bar{i}} (dz^j \wedge dz^I \wedge d\bar{z}^j \wedge d\bar{z}^J) \\ &= (\sqrt{-1})^2 \sum_{i,j=1}^n (-1)^{2p+1} \varphi_{IJ} \iota_i (dz^j \wedge dz^I \wedge \iota_{\bar{i}} (d\bar{z}^j \wedge d\bar{z}^J)) \\ &= (\sqrt{-1})^2 \sum_{i,j=1}^n (-1)^{2p+1} \varphi_{IJ} \iota_i \left(dz^j \wedge dz^I \wedge \underbrace{(\delta_{\bar{i}}^{\bar{j}} d\bar{z}^J - d\bar{z}^j \wedge \iota_{\bar{i}} d\bar{z}^J)}_A \right) \\ &= (\sqrt{-1})^2 \sum_{i,j=1}^n (-1)^{2p+1} \varphi_{IJ} (\delta_i^j dz^I \wedge A - dz^j \wedge \iota_i dz^I \wedge A) \end{aligned}$$

Then

$$\begin{aligned} L\Lambda\varphi - \Lambda L\varphi &= (\sqrt{-1})^2 \sum_{j=1}^n \varphi_{IJ} \left(dz^I \wedge d\bar{z}^J - dz^I \wedge d\bar{z}^j \wedge \iota_{\bar{j}} dz^J - dz^j \wedge \iota_j dz^I \wedge d\bar{z}^J \right) \\ &= (k - n)\varphi \end{aligned}$$

□

9.2. Hodge decomposition.

Theorem 9.2.1. Let (X, h) be a compact Kähler manifold, $\alpha = \sum_{p+q=k} \alpha^{p,q}$. Then α is harmonic if and only if $\alpha^{p,q}$ is harmonic, that is

$$\mathcal{H}^k \otimes_{\mathbb{R}} \mathbb{C} = \bigoplus_{p+q=k} \mathcal{H}^{p,q}$$

with $\overline{\mathcal{H}^{p,q}} = \mathcal{H}^{q,p}$.

Proof. It follows from Δ_d preserves bidegree. □

Theorem 9.2.2 (Hodge decomposition). Let (X, h) be a compact Kähler manifold. Then

$$H^k(X, \mathbb{C}) \cong \bigoplus_{p+q=k} H^{p,q}(X)$$

with $\overline{H^{p,q}(X)} = H^{q,p}(X)$.

Proof. It follows from there are natural isomorphisms $H^k(X, \mathbb{C}) \cong \mathcal{H}^k \otimes \mathbb{C}$ and $H^{p,q}(X) \cong \mathcal{H}^{p,q}$. □

Corollary 9.2.1. Let (X, h) be a compact Kähler manifold. Then

$$b_k = \sum_{p+q=k} h^{p,q}$$


with $h^{p,q} = h^{q,p}$, where $b_k = \dim H^k(X, \mathbb{C})$ and $h^{p,q} = \dim H^{p,q}(X)$.

Corollary 9.2.2. b_k is even when k is odd.

Corollary 9.2.3. $b_k \neq 0$ when k is even.

Proof. $h^{k,k} \neq 0$ since $0 \neq \omega^k \in H^{k,k}(X)$. □

There are many relations between $h^{p,q}$, and we can draw a picture as follows, called Hodge diamond since it has the same symmetry as a diamond.



Example 9.2.1.

$$H^{p,q}(\mathbb{C}\mathbb{P}^n) = \begin{cases} \mathbb{C} & 0 \leq p = q \leq n \\ 0 & \text{otherwise} \end{cases}$$

Proof. It's known to all that the singular cohomology of $\mathbb{C}\mathbb{P}^n$ with complex coefficient is

$$H^k(\mathbb{C}\mathbb{P}^n, \mathbb{C}) = \begin{cases} \mathbb{C} & k \text{ is even} \\ 0 & k \text{ is odd} \end{cases}$$

Thus it's clear to compute Dolbeault cohomology of $\mathbb{C}\mathbb{P}^n$ using the symmetry of Hodge diamond. \square

9.3. Bott-Chern cohomology. In the proof Hodge decomposition, we used the Kähler metric. A natural question is to consider (in)dependence of the Kähler metric. In this section we will show our decomposition is independent of the choice of Kähler metric, by using Bott-Chern cohomology.

Definition 9.3.1 (Bott-Chern cohomology). Let X be a complex manifold. The Bott-Chern cohomology is defined as

$$H_{\text{BC}}^{p,q}(X) := \frac{Z_{\text{BC}}^{p,q} := \{\alpha \in C^\infty(X, \Omega_X^{p,q}) \mid d\alpha = 0\}}{\partial\bar{\partial}C^\infty(X, \Omega_X^{p-1,q-1})}$$

Remark 9.3.1. There is a natural map

$$Z_{\text{BC}}^{p,q}(X) \rightarrow H^{p+q}(X, \mathbb{C})$$

which descends to

$$H_{\text{BC}}^{p,q}(X) \rightarrow H^{p+q}(X, \mathbb{C})$$

since $\partial\bar{\partial}\beta = d\bar{\partial}\beta$. On the other hand, there is also a natural map

$$Z_{\text{BC}}^{p,q}(X) \rightarrow H^{p,q}(X)$$

which descends to

$$H_{\text{BC}}^{p,q}(X) \rightarrow H^{p,q}(X)$$

since $\partial\bar{\partial}\beta = -\bar{\partial}\partial\beta$. So if we can prove there are isomorphisms between

$$\begin{aligned} H_{\text{BC}}^{p,q}(X) &\cong H^{p,q}(X) \\ \bigoplus_{p+q=k} H_{\text{BC}}^{p,q}(X) &\cong H^k(X, \mathbb{C}) \end{aligned}$$

then Hodge decomposition is canonical, that is independent of choice of Kähler metric since Bott-Chern cohomology is independent of the choice of Kähler metric.

Lemma 9.3.1 ($\partial\bar{\partial}$ -lemma). Let (X, ω) be a compact Kähler manifold and α be a d-closed (p, q) -form. If α is $\bar{\partial}$ -exact or ∂ -exact, then there exists a $(p-1, q-1)$ -form such that

$$\alpha = \partial\bar{\partial}\beta$$

Proof. Suppose α is $\bar{\partial}$ -exact. Then $\alpha = \bar{\partial}\gamma$ for some $(p, q - 1)$ -form γ , and Hodge's theorem implies γ has decomposition

$$\gamma = a + \partial b + \partial^* c$$

where a is Δ_{∂} -harmonic, and b, c are forms with appropriate degrees. Direct computation shows

$$\begin{aligned} \alpha &= \bar{\partial}\gamma = \bar{\partial}a + \bar{\partial}\partial b + \bar{\partial}\partial^* c \\ &= -\partial\bar{\partial}b + \bar{\partial}\partial^* c \\ &= -\partial\bar{\partial}b - \partial^*\bar{\partial}c \end{aligned}$$

Now it suffices to show $-\partial^*\bar{\partial}c = 0$. A trick here is to note that

$$0 = \partial\alpha = -\partial\partial^*\bar{\partial}c \implies \partial^*\bar{\partial}c \in \ker \partial \cap \text{im } \partial^* = 0 \implies \partial^*\bar{\partial}c = 0$$

So we have

$$\alpha = \partial\bar{\partial}(-b)$$

as desired. \square

Corollary 9.3.1. Let (X, ω) be a compact Kähler manifold. Then

- (1) $H_{\text{BC}}^{p,q}(X) \rightarrow H^{p,q}(X)$ is an isomorphism.
- (2) $\bigoplus_{p+q=k} H_{\text{BC}}^{p,q}(X) \rightarrow H^k(X, \mathbb{C})$ is an isomorphism.

Proof. Here we only prove the first isomorphism. From Remark 9.3.1, there is a canonical map $H_{\text{BC}}^{p,q}(X) \rightarrow H^{p,q}(X)$, and if we choose a Kähler metric, we have $H^{p,q}(X) \cong \mathcal{H}^{p,q}$, we will show our canonical map is both surjective and injective via this chosen metric.

- (1) To see surjectivity: For element in $H^{p,q}(X)$ we choose a $\Delta_{\bar{\partial}}$ -harmonic representative. Since Δ_{∂} -harmonic is equivalent to Δ_{d} -harmonic, so this representative is also d-closed.
- (2) To see injectivity: Suppose we have $[\alpha] \in H_{\text{BC}}^{p,q}(X)$ such that α is trivial in $H^{p,q}(X)$, that is $\bar{\partial}$ -exact. Then Lemma 9.3.1, that is $\partial\bar{\partial}$ -lemma implies it's trivial in Bott-Chern cohomology. \square

Corollary 9.3.2. A Hermitian metric ω is Kähler if and only if it can be written locally as

$$\omega = \sqrt{-1}\partial\bar{\partial}f$$

where f is a real-valued smooth function.

Proof. It's clear if ω is locally written as $\sqrt{-1}\partial\bar{\partial}f$, then it gives a Kähler metric. Conversely, a Kähler metric ω is an element in $H^{1,1}(X)$, and we have already shown that $H^{1,1}(X) = H_{\text{BC}}^{1,1}(X)$, and Dolbeault lemma implies Dolbeault cohomology vanishes on open subset which is sufficiently small, this completes the proof. \square

10. SERRE DUALITY

10.1. Operators on bundle valued forms. Let (E, h) be a Hermitian holomorphic vector bundle on Hermitian n -manifold (X, g) , what we have done can be generalized to bundle valued forms. More explicitly, for $\varphi, \psi \in C^\infty(X, \Omega_X^{p,q} \otimes E)$, locally written as $\varphi = \varphi^\alpha e_\alpha, \psi = \psi^\beta e_\beta$, then local inner product is given by

$$\langle \varphi, \psi \rangle := h_{\alpha\bar{\beta}} \langle \varphi^\alpha, \psi^\beta \rangle,$$

where $\langle \varphi^\alpha, \psi^\beta \rangle$ is induced by the Hermitian metric g .

(1) The inner product on $C^\infty(X, \Omega_X^{p,q} \otimes E)$ is given by

$$(\varphi, \psi) := \int_X \langle \varphi, \psi \rangle \frac{\omega^n}{n!},$$

where $\varphi, \psi \in C^\infty(X, \Omega_X^{p,q} \otimes E)$.

(2) The Hodge star operator is the operator

$$\star_E: C^\infty(X, \Omega_X^{p,q} \otimes E) \rightarrow C^\infty(X, \Omega_X^{n-q, n-p} \otimes E)$$

such that

$$(\varphi, \psi) = \int_X \varphi \wedge \star_E \bar{\psi},$$

where $\varphi, \psi \in C^\infty(X, \Omega_X^{p,q} \otimes E)$.

Let ∇ be the Chern connection of (E, h) . By definition one has $\nabla^{0,1} = \bar{\partial}_E$, and if we set $\nabla^{1,0} = \partial_E$, then

$$\begin{aligned} \Theta_h &= \nabla^2 \\ &= \partial_E^2 + \partial_E \bar{\partial}_E + \bar{\partial}_E \partial_E + \bar{\partial}_E^2 \\ &= [\partial_E, \bar{\partial}_E]. \end{aligned}$$

Exercise 10.1.1. Give formulas of ∂_E^* and $\bar{\partial}_E^*$ in terms of Hodge star \star_E .

Proof. □

Exercise 10.1.2. Give formulas of Laplacians $\Delta_{\partial_E}, \Delta_{\bar{\partial}_E}$ in terms of Hodge star \star_E .

Proof. □

Then there is also a Hodge decomposition given by

$$C^\infty(X, \Omega_X^{p,q} \otimes E) = \mathcal{H}^{p,q}(X, E) \oplus \text{im } \bar{\partial}_E \oplus \text{im } \bar{\partial}_E^*,$$

and

$$H^{p,q}(X, E) \cong \mathcal{H}^{p,q}$$

Lefschetz operator is defined as follows

$$\begin{aligned} L: C^\infty(X, \Omega_X^{p,q} \otimes E) &\rightarrow C^\infty(X, \Omega_X^{p+1, q+1} \otimes E) \\ \alpha &\mapsto \omega \wedge \alpha \end{aligned}$$

and Λ is the formal adjoint of L . If (X, ω) is also a Kähler n -manifold, then there are also Kähler identities

$$\begin{aligned} [\bar{\partial}_E^*, L] &= \sqrt{-1} \partial_E \\ [\partial_E^*, L] &= -\sqrt{-1} \cdot \bar{\partial}_E \\ [\Lambda, \bar{\partial}_E] &= -\sqrt{-1} \partial_E^* \\ [\Lambda, \partial_E] &= \sqrt{-1} \cdot \bar{\partial}_E^*, \end{aligned}$$

and

$$[L, \Lambda] = (p + q - n) \text{id}$$

holds on E -valued (p, q) -forms.

10.2. Serre duality.

Theorem 10.2.1 (Serre duality). Let X be a compact complex n -manifold and E be a holomorphic vector bundle. Then there exists a non-degenerate \mathbb{C} -linear pairing

$$\begin{aligned} H^{p,q}(X, E) \times H^{n-p, n-q}(X, E^*) &\rightarrow \mathbb{C} \\ ([\alpha], [\beta]) &\mapsto \int_X \alpha \wedge \beta \end{aligned}$$

In particular, we have

$$H^{p,q}(X, E) = H^{n-p, n-q}(X, E^*)^*$$

Sketch of the proof. Let h be a Hermitian metric on E . Firstly prove that

$$\Delta_{\bar{\partial}_E^*} \circ \bar{\kappa}_E = \bar{\kappa}_E \circ \Delta_{\bar{\partial}_E}.$$

This give an isomorphism

$$\bar{\kappa}_E: \mathcal{H}^{p,q}(X, E) \xrightarrow{\cong} \mathcal{H}^{n-p, n-q}(X, E^*).$$

On the other hand, the Hodge theorem shows that

$$\mathcal{H}^{p,q}(X, E) \cong H^{p,q}(X, E).$$

For any $\alpha \in \mathcal{H}^{p,q}(X, E)$ and $\beta \in \mathcal{H}^{n-p, n-q}(X, E^*)$, there exists some $\gamma \in \mathcal{H}^{p,q}(X, E)$ such that $\beta = \bar{\kappa}_E \gamma$, and thus

$$\int_X \alpha \wedge \beta = \int_X \alpha \wedge \bar{\kappa}_E \gamma = \langle \alpha, \gamma \rangle$$

is non-degenerate. □

Corollary 10.2.1. Let X be a compact complex n -manifold and E be a holomorphic vector bundle. Then

$$H^{p,q}(X) = H^{n-p, n-q}(X)^*.$$

Proof. Consider $E = \mathcal{O}_X$ in Serre duality, and then desired result holds from the fact $\mathcal{O}_X^* = \mathcal{O}_X$. □

Remark 10.2.1. This recovers Corollary 8.2.6.

Corollary 10.2.2. Let X be a compact complex n -manifold and E be a holomorphic vector bundle. Then

$$H^q(X, E) = H^{n-q}(X, K_X \otimes E^*)^*$$

Proof. Set $p = 0$ in Serre duality one has

$$H^{0,q}(X, E) \cong H^{n,n-q}(X, E^*)^*$$

which gives desired result. \square

11. LEFSCHETZ DECOMPOSITION

11.1. Lefschetz decomposition.

Proposition 11.1.1. Let (X, ω) be a Kähler n -manifold. Then $L^{n-k}: C^\infty(X, \Omega_{X, \mathbb{R}}^k) \rightarrow C^\infty(X, \Omega_{X, \mathbb{R}}^{2n-k})$ is an isomorphism for $k \leq n$.

Proof. In fact, we will prove that L^r are injective for all $1 \leq r \leq n - k$. As a consequence, L^{n-k} is an isomorphism since $\Omega_{X, \mathbb{R}}^k$ has the same rank as $\Omega_{X, \mathbb{R}}^{2n-k}$. In Exercise 9.1.1 we have shown that

$$[L, \Lambda]\alpha = (k - n)\alpha, \quad \forall \alpha \in C^\infty(X, \Omega_{X, \mathbb{R}}^k).$$

Then

$$\begin{aligned} [L^r, \Lambda] &= L^r \Lambda - \Lambda L^r \\ &= L(L^{r-1} \Lambda - \Lambda L^{r-1}) + L \Lambda L^{r-1} - \Lambda L L^{r-1} \\ &= L[L^{r-1}, \Lambda] + [L, \Lambda] L^{r-1}. \end{aligned}$$

By induction it's easy to show the following identity

$$[L^r, \Lambda]\alpha = (r(k - n) + r(r - 1))L^{r-1}\alpha, \quad \forall \alpha \in C^\infty(X, \Omega_{X, \mathbb{R}}^k).$$

For $\alpha \in C^\infty(X, \Omega_{X, \mathbb{R}}^k)$, if $L^r \alpha = 0, r \leq n - k$, then

$$\begin{aligned} L^r \Lambda \alpha &= [L^r, \Lambda]\alpha \\ &= (r(k - n) + r(r - 1))L^{r-1}\alpha. \end{aligned}$$

In other words, we have

$$(11.1) \quad L^{r-1}(L \Lambda \alpha - (r(k - n) + r(r - 1))\alpha) = 0.$$

Now let's prove L^r is injective by induction on r : It's clear L is injective, and suppose L^{r-1} is injective. Then by (11.1) one has

$$L \Lambda \alpha = (r(k - n) + r(r - 1))\alpha.$$

If we denote $\beta = \Lambda \alpha$, and apply L^r to both side of above equation, then we have

$$L^{r+1}\beta = (r(k - n) + r(r - 1))L^r \alpha = 0,$$

where $\beta \in C^\infty(X, \Omega_{X, \mathbb{R}}^{k-2})$. It's clear $\beta = 0$ if β is a smooth function. Then by induction on k , we have $\beta = 0$, and thus $\alpha = 0$. \square

Definition 11.1.1 (primitive form). Let (X, ω) be a Kähler n -manifold. A k -form α is called primitive if $L^{n-k+1}\alpha = 0$.

Exercise 11.1.1. A k -form α is primitive if and only if $\Lambda \alpha = 0$.

Proof. For an n -form α , α is primitive if and only if $L \alpha = 0$. On the other hand, the Exercise 9.1.1 implies that

$$[L, \Lambda] = (k - n) \text{id}, \quad \text{on } C^\infty(X, \Omega_{X, \mathbb{R}}^k).$$

This shows if $n = k$, then L and Λ commutes. Thus we have α is primitive if and only if $\Lambda\alpha = 0$, since

$$\Lambda\alpha = 0 \iff L\Lambda\alpha = 0 \iff \Lambda L\alpha = 0 \iff L\alpha = 0$$

and the first and last equality we use the fact that L is injective on $\Omega_{X,\mathbb{R}}^k$, $k \leq n$ and Λ is injective on $\Omega_{X,\mathbb{R}}^{n+2}$. In general case, we have

$$[L^r, \Lambda]\alpha = (r(k-n) + r(r-1))L^{r-1}\alpha$$

and in particular for $r = n - k + 1$ where k is the degree of α , we have

$$[L^r, \Lambda]\alpha = 0$$

The argument can be repeated to conclude. \square

Proposition 11.1.2. For any k -form α , there exists a unique decomposition

$$\alpha = \sum_r L^r \alpha_r,$$

where α_r is primitive $(k - 2r)$ -form.

Proof. Firstly let's prove the uniqueness: If $\sum_r L^r \alpha_r = 0$ with primitive α_r , we need to show $\alpha_r = 0$. If not, then take the largest r_m such that $\alpha_{r_m} \neq 0$. By the choice of α_{r_m} , L^{n-k+r_m} kills everything in $\sum_r L^r \alpha_r$ but $L^{r_m} \alpha_{r_m}$. Then

$$0 = L^{n-k+r_m} \left(\sum_r L^r \alpha_r \right) = L^{n-k+r_m} (L^{r_m} \alpha_{r_m}) \neq 0,$$

which is a contradiction.

Now let's prove the existence: Since $L^{n-k+2}: C^\infty(X, \Omega_{X,\mathbb{R}}^{k-2}) \rightarrow C^\infty(X, \Omega_{X,\mathbb{R}}^{2n-k+2})$ is an isomorphism, then there exists $\beta \in C^\infty(X, \Omega_{X,\mathbb{R}}^{k-2})$ such that

$$L^{n-k+1}\alpha = L^{n-k+2}\beta.$$

Then $\alpha - L\beta$ is primitive a primitive k -form, that is

$$\alpha = (\alpha - L\beta) + L\beta.$$

By induction on k , we have primitive decomposition for $\beta \in C^\infty(X, \Omega_{X,\mathbb{R}}^{k-2})$. and this completes the proof. \square

Remark 11.1.1. If we define $H = [L, \Lambda]$, then (L, H, Λ) generates an \mathfrak{sl}_2 -action on $\bigoplus_k C^\infty(X, \Omega_{X,\mathbb{R}}^k)$.

In fact, the Lefschetz operator also defines a map between cohomology groups

$$\begin{aligned} L: H^k(X, \mathbb{R}) &\rightarrow H^{k+2}(X, \mathbb{R}) \\ [\alpha] &\mapsto [\omega \wedge \alpha]. \end{aligned}$$

Now let's see it's well-defined:

(1) If α is closed, then

$$d(\omega \wedge \alpha) = d\omega \wedge \alpha + \omega \wedge d\alpha = 0.$$

(2) If $\alpha = d\beta$, then

$$\omega \wedge d\beta = d\omega \wedge \beta + \omega \wedge d\beta = d(\omega \wedge \beta).$$

Theorem 11.1.1 (hard Lefschetz theorem¹²). Let (X, ω) be a compact Kähler n -manifold. Then

$$L^{n-k}: H^k(X, \mathbb{R}) \rightarrow H^{2n-k}(X, \mathbb{R})$$

is an isomorphism for $1 \leq k \leq n$.

Proof. In Exercise 9.1.1 we have shown $[\Delta_d, L] = 0$, so the Lefschetz operator induces a map between harmonic forms as follows

$$L^{n-k}: \mathcal{H}^k \rightarrow \mathcal{H}^{2n-k}.$$

By Proposition 11.1.1 L^{n-k} is injective and $\mathcal{H}^k, \mathcal{H}^{2n-k}$ have the same dimension, we obtain the desired result. \square

Definition 11.1.2 (primitive form). Let (X, ω) be a compact Kähler n -manifold. For $[\alpha] \in H^k(X, \mathbb{R})$, it's called primitive, if $L^{n-k+1}[\alpha] = 0$.

Notation 11.1.1. $H^k(X, \mathbb{R})_{\text{prim}}$ denotes the set of all primitive forms.

Corollary 11.1.1 (Lefschetz decomposition). There is the following decomposition

$$H^k(X, \mathbb{R}) = \bigoplus_r L^r H^{k-2r}(X, \mathbb{R})_{\text{prim}}.$$

Remark 11.1.2. If $[\omega] \in H^2(X, \mathbb{Z})$, such as ω comes from a positive holomorphic line bundle, then we can state theorem and corollary for $H^k(X, \mathbb{Q})$.

Moreover, we have the following isomorphism

$$L^{n-k}: H^{p,q}(X) \rightarrow H^{n-q, n-p}(X)$$

for $k = p + q \leq n$.

Corollary 11.1.2. Let (X, ω) be a compact Kähler n -manifold. Then for $2 \leq k \leq n$, one has $b_{k-2} \leq b_k$ and $h^{p-1, q-1} \leq h^{p, q}$ with $k = p + q$.

11.2. Hodge index.

11.2.1. Surface case.

Example 11.2.1. For open subset $U \subseteq \mathbb{C}^2$ equipped with canonical Kähler form

$$\omega = \sqrt{-1} (dz^1 \wedge d\bar{z}^1 + dz^2 \wedge d\bar{z}^2).$$

The volume form is given by

$$\text{vol} = \frac{\omega^2}{2!} = - (dz^1 \wedge d\bar{z}^1 \wedge dz^2 \wedge d\bar{z}^2).$$

¹²Though proof of this theorem is quite easy using tools we have, but it's quite hard for Lefschetz, since during his time, there is no Hodge theorem. Here we use L to denote Lefschetz operator, in order to honor Lefschetz.

Suppose α is a $(2, 0)$ form written as

$$\alpha = adz^1 \wedge dz^2.$$

Now we're going to compute $\star\bar{\alpha}$, which is also a $(2, 0)$ -form. Suppose $\star\bar{\alpha} = bdz^1 \wedge dz^2$. Then by definition, for an arbitrary $(0, 2)$ -form β one has

$$\langle \beta, \alpha \rangle \text{vol} = \beta \wedge \star\bar{\alpha}.$$

In particular, if we choose $\beta = d\bar{z}^1 \wedge d\bar{z}^2$, then

$$\begin{aligned} \beta \wedge \star\bar{\alpha} &= -bdz^1 \wedge d\bar{z}^1 \wedge dz^2 \wedge d\bar{z}^2 \\ \{\beta, \alpha\} \text{vol} &= \{adz^1 \wedge dz^2, d\bar{z}^1 \wedge d\bar{z}^2\} \times -dz^1 \wedge d\bar{z}^1 \wedge dz^2 \wedge d\bar{z}^2 \\ &= -adz^1 \wedge d\bar{z}^1 \wedge dz^2 \wedge d\bar{z}^2. \end{aligned}$$

This shows $\alpha = \star\bar{\alpha}$. By the same computation one has $\alpha = \star\bar{\alpha}$ holds for a $(0, 2)$ -form α . On the other hand, it's clear $(2, 0)$ -form and $(0, 2)$ -form are automatically primitive.

Now we're going to see if a $(1, 1)$ -form α is primitive, what's the relation between α and $\star\bar{\alpha}$. For $(1, 1)$ -form α , written as

$$\alpha = a_{11}dz^1 \wedge d\bar{z}^1 + a_{22}dz^2 \wedge d\bar{z}^2 + a_{12}dz^1 \wedge d\bar{z}^2 + a_{21}dz^2 \wedge d\bar{z}^1.$$

A direct computation shows that

$$\star\bar{\alpha} = a_{22}dz^1 \wedge d\bar{z}^1 + a_{11}dz^2 \wedge d\bar{z}^2 - a_{12}dz^1 \wedge d\bar{z}^2 - a_{21}dz^2 \wedge d\bar{z}^1.$$

On the other hand,

$$L\alpha = \omega \wedge \alpha = \sqrt{-1}(a_{11} + a_{22})dz^1 \wedge d\bar{z}^1 \wedge dz^2 \wedge d\bar{z}^2.$$

Then

$$L\alpha = 0 \iff a_{11} + a_{22} = 0 \iff \star\bar{\alpha} = -\alpha.$$

Lemma 11.2.1. Let (X, ω) be a Kähler surface. If (p, q) -form α is primitive 2-form, then

$$\star\bar{\alpha} = (-1)^p \alpha.$$

Proof. By taking normal coordinate, it suffices to consider $U \subseteq \mathbb{C}^2$, and that's exactly what we have done in Example 11.2.1. \square

Let X be a compact Kähler surface. The Poincaré duality and Stokes theorem imply that we have the following well-defined non-degenerate pairing

$$\begin{aligned} Q: H^2(X, \mathbb{R}) \times H^2(X, \mathbb{R}) &\rightarrow \mathbb{R} \\ ([\alpha], [\beta]) &\mapsto \int_X \alpha \wedge \beta \end{aligned}$$

Then we obtain a Hermitian form by considering

$$H([\alpha], [\beta]) = Q([\alpha], \overline{[\beta]}).$$

Lemma 11.2.2. The Lefschetz decomposition $H^2(X, \mathbb{R}) = H^2(X, \mathbb{R})_{\text{prim}} \oplus \mathbb{R} \cdot [\omega]$ is orthonormal with respect to Q .

Proof.

$$Q([\omega], [\alpha]) = \int_X \omega \wedge \alpha = \int_X L\alpha = 0$$

for α is primitive and harmonic. \square

Theorem 11.2.1. $H^2(X, \mathbb{C})_{\text{prim}} = \bigoplus_{p+q=2} H^{p,q}(X)_{\text{prim}}$ is orthonormal with respect to H , and $(-1)^p H$ is positive definite on $H^{p,q}(X)_{\text{prim}}$.

Proof. It's clear above decomposition is orthonormal, since

$$\int_X \alpha = 0$$

if α is not a $(2, 2)$ -form. To see $(-1)^p H$ is positive definite on $H^{p,q}(X)_{\text{prim}}$, we take a harmonic representative α for any non-zero primitive cohomology class in $H^{p,q}(X)_{\text{prim}}$. Then

$$\begin{aligned} (-1)^p H([\alpha], [\alpha]) &= (-1)^p \int_X \alpha \wedge \bar{\alpha} \\ &= (-1)^{p+q} \int_X \alpha \wedge \star \bar{\alpha} \\ &= \|\alpha\|^2 > 0. \end{aligned}$$

This shows $(-1)^p H$ is positive definite on $H^{p,q}(X)_{\text{prim}}$. \square

Corollary 11.2.1 (Hodge index). The index of H defined on $H^2(X, \mathbb{C}) \cap H^{1,1}(X)$ is $(1, h^{1,1} - 1)$.

Proof. Note that there is the following decomposition

$$H^2(X, \mathbb{C}) \cap H^{1,1}(X) = H^{1,1}(X)_{\text{prim}} \oplus \mathbb{C}[\omega],$$

and we have already shown that H is negative definite on $H^{1,1}(X)_{\text{prim}}$. Then the index for H on $H^2(X, \mathbb{C}) \cap H^{1,1}(X)$ is $(1, h^{1,1} - 1)$. \square

11.2.2. *General case.* In this section we will introduce a more general case: Let (X, ω) be a compact Kähler n -manifold. Then by Lefschetz decomposition we have

$$H^k(X, \mathbb{R}) = \bigoplus_r L^r H^{k-2r}(X, \mathbb{R})_{\text{prim}}, \quad k \leq n,$$

and by Hodge decomposition we have a more explicit decomposition

$$H^k(X, \mathbb{C})_{\text{prim}} = \bigoplus_{p+q=k} H^{p,q}(X)_{\text{prim}}.$$

As we have seen in the case of surface, H will be positive definite or negative definite in these (p, q) components. Now we introduce some symbols, in order to get a neater result.

Consider

$$Q: H^k(X, \mathbb{R}) \times H^k(X, \mathbb{R}) \rightarrow \mathbb{R}$$

$$([\alpha], [\beta]) \mapsto (-1)^{\frac{k(k-1)}{2}} \int_X \omega^{n-k} \wedge \alpha \wedge \beta.$$

Then Q is a bilinear form, and it is symmetric when k is even and anti-symmetric when k is odd.

Definition 11.2.1 (Weil operator). The Weil operator $\mathfrak{C}: H^k(X, \mathbb{C}) \rightarrow H^k(X, \mathbb{C})$ is defined by $\mathfrak{C}|_{H^{p,q}(X)} \mapsto \sqrt{-1}^{p-q} \text{id}$.

Remark 11.2.1. The Weil operator \mathfrak{C} maps $H^k(X, \mathbb{R})$ to $H^k(X, \mathbb{R})$ in fact:

$$\mathfrak{C}|_{\overline{H^{p,q}(X)}} = \mathfrak{C}|_{H^{q,p}(X)} = \sqrt{-1}^{q-p} \text{id} = \overline{\sqrt{-1}^{p-q}} \text{id} = \overline{\mathfrak{C}|_{H^{p,q}(X)}}.$$

Now we define

$$H: H^k(X, \mathbb{C}) \times H^k(X, \mathbb{C}) \rightarrow \mathbb{C}$$

$$([\alpha], [\beta]) \mapsto Q(\mathfrak{C}[\alpha], \overline{[\beta]}).$$

In other words, we have

$$H([\alpha], [\beta]) = (-1)^{\frac{k(k-1)}{2}} \sqrt{-1}^{p-q} \int_X \omega^{n-k} \wedge \alpha \wedge \overline{\beta}, \quad \alpha, \beta \in H^{p,q}(X).$$

Exercise 11.2.1. H is a Hermitian form on $H^{p,q}(X)$.

Proof. For $[\alpha], [\beta] \in H^{p,q}(X)$, one has

$$\begin{aligned} \overline{H([\alpha], [\beta])} &= (-1)^{\frac{k(k-1)}{2}} (-1)^{p-q} \sqrt{-1}^{p-q} \int_X \omega^{n-k} \wedge \overline{\alpha} \wedge \beta \\ &= (-1)^{\frac{k(k-1)}{2}} (-1)^{p-q} \sqrt{-1}^{p-q} (-1)^{(p+q)^2} \int_X \omega^{n-k} \wedge \beta \wedge \overline{\alpha}. \end{aligned}$$

Note that

$$(p+q)^2 - p - q = 2pq + p(p-1) + q(q-1)$$

is always even, this completes the proof. \square

Lemma 11.2.3. Let α be a primitive (p, q) -form with $p+q=k$. Then

$$\star \alpha = (-1)^{\frac{k(k+1)}{2}} \sqrt{-1}^{p-q} \frac{L^{n-k} \alpha}{(n-k)!}.$$

Theorem 11.2.2 (Hodge-Riemann bilinear relations).

- (1) $H^k(X, \mathbb{R}) = \bigoplus_r L^r H^{k-2r}(X, \mathbb{R})_{\text{prim}}$ is orthonormal with respect to Q .
- (2) $H^k(X, \mathbb{C})_{\text{prim}} = \bigoplus_{p+q=k} H^{p,q}(X)_{\text{prim}}$ is orthonormal with respect to H .
- (3) H is positive definite on $H^{p,q}(X)_{\text{prim}}$.

Proof. For (1). For $r < s$, note that

$$\omega^{n-k} \wedge L^r \gamma \wedge L^s \delta = (L^{n-k+r+s} \gamma) \wedge \delta = 0$$

since $L^{n-k+2r+1} \gamma = 0$ and $r < s$.

For (2). If α is a (p, q) -form, and β is (p', q') -form, and $(p, q) \neq (p', q')$, then $\omega^{n-k} \wedge \alpha \wedge \bar{\beta}$ is not a (n, n) -form.

For (3). To see H is positive definite on $H^{p,q}(X)_{\text{prim}}$, we take a harmonic representative α for any non-zero primitive cohomology class in $H^{p,q}(X)_{\text{prim}}$. Then

$$H([\alpha], [\alpha]) = (-1)^{\frac{k(k-1)}{2}} \sqrt{-1}^{p-q} \int_X \omega^{n-k} \wedge \alpha \wedge \bar{\alpha}$$

By Lemma 11.2.3 one has

$$\begin{aligned} \star \bar{\alpha} &= (-1)^{\frac{k(k+1)}{2}} \sqrt{-1}^{q-p} \frac{L^{n-k} \bar{\alpha}}{(n-k)!} \\ &= (-1)^{\frac{k(k-1)}{2}} \sqrt{-1}^{p-q} \frac{L^{n-k} \bar{\alpha}}{(n-k)!}. \end{aligned}$$

Then

$$H([\alpha], [\alpha]) = (n-k)! \int_X \alpha \wedge \star \bar{\alpha} = (n-k)! \|\alpha\|^2 > 0.$$

□

Corollary 11.2.2 (Hodge index theorem). Let X be a compact Kähler n -manifold with n is even¹³. Then $\int_X \alpha \wedge \beta$ on $H^n(X, \mathbb{R})$ is of signature

$$\sum_{p,q} (-1)^p h^{p,q}$$

where summation runs over all p, q .

Proof. Note that the signature of $\int_X \alpha \wedge \beta$ on $H^n(X, \mathbb{R})$ is the same as the signature of $\int_X \alpha \wedge \bar{\beta}$ on $H^n(X, \mathbb{C})$. We write

$$H^n(X, \mathbb{C}) = \bigoplus_{\substack{p+q+2r=n \\ r,p,q \in \mathbb{Z}_{\geq 0}}} L^r H^{p,q}(X)_{\text{prim}}$$

Then Hodge-Riemann bilinear theorem implies that $\int_X \alpha \wedge \bar{\beta}$ is $(-1)^p$ -definite on $L^r H^{p,q}(X)_{\text{prim}}$, where we used the fact n is even. Then we have the signature is

$$\sum_{p+q+2r=n} (-1)^p h_{\text{prim}}^{p,q}.$$

But $h_{\text{prim}}^{p,q} = h^{p,q} - h^{p-1,q-1}$, so

$$\sum_{p+q+2r=n} (-1)^p (h^{p,q} - h^{p-1,q-1}).$$

Note that $p+q = n$ counted once and $p+q < n$ counted twice, so rewrite it as

$$\sum_{p+q \text{ even}} (-1)^p h^{p,q},$$

¹³In this case $\int_X \alpha \wedge \beta$ is symmetric on $H^n(X, \mathbb{R})$.

since $h^{p,q} = h^{n-p,n-q}$. And this is also equivalent to sum all p, q , since

$$\sum_{p+q \text{ odd}} (-1)^p h^{p,q} = 0$$

This completes the proof. □

Example 11.2.2. For surface, we have

$$H^2(X, \mathbb{C}) = H^{2,0}(X) \oplus H^{1,1}(X)_{\text{prim}} \oplus \mathbb{C}[\omega] \oplus H^{0,2}(X)$$

Then this corollary implies

$$h^{0,0} + h^{2,0} - h^{1,1} + h^{0,2} + h^{2,2} = h^{2,0} + h^{0,2} + (1 - (h^{1,1} - 1)),$$

which recovers what we have done in the case of surface.

Part 4. Positivity and vanishing theorems

12. POSITIVITY

12.1. Positivity of line bundle. Let (L, h) be a Hermitian holomorphic line bundle over a complex manifold X . Then $\frac{\sqrt{-1}}{2\pi}\Theta_h$ gives a real $(1, 1)$ -form, so it corresponds to a Hermitian form on TX .

Definition 12.1.1 (positive line bundle). Let L be a holomorphic line bundle over X . L is called positive if it admits a Hermitian metric h such that the Hermitian form corresponding to $\frac{\sqrt{-1}}{2\pi}\Theta_h$ is positive definite.

Remark 12.1.1. The Kodaira embedding theorem implies positive line bundle is exactly ample divisor in algebraic geometry.

Remark 12.1.2 (local form). Locally, one has

$$\frac{\sqrt{-1}}{2\pi}\Theta_h = -\frac{\sqrt{-1}}{2\pi}\partial\bar{\partial}\log h = \frac{\sqrt{-1}}{2\pi}\frac{\partial^2\varphi}{\partial z^i\partial\bar{z}^j}dz^i \wedge d\bar{z}^j$$

where $\varphi = -\log h$. Thus L is positive if and only if the Hermitian matrix $(\frac{\partial^2\varphi}{\partial z^i\partial\bar{z}^j})$ is positive definite everywhere.

Proposition 12.1.1. If X admits a positive holomorphic line bundle, then X is Kähler.

Proof. The first Chern class of (L, h) gives its Kähler form. \square

Proposition 12.1.2. L is positive if and only if $L^{\otimes m}$ is positive for some $m \in \mathbb{N}_{\geq 0}$.

Proof. For a line bundle L locally we have the Hermitian metric corresponding to its curvature looking like

$$\left(\frac{\partial^2\varphi}{\partial z^i\partial\bar{z}^j}\right)$$

and for $L^{\otimes m}$, $m \in \mathbb{N}_{\geq 0}$ we have

$$\left(m \cdot \frac{\partial^2\varphi}{\partial z^i\partial\bar{z}^j}\right)$$

It's clear L is positive if and only if $L^{\otimes m}$ is. \square

Exercise 12.1.1. Let X be a compact complex manifold and L be a positive line bundle. For any holomorphic line bundle L' , there exists $N_0 \in \mathbb{N}$ such that $L' \otimes L^{\otimes N}$ positive for $N \geq N_0$.

Proof. The proof is quite similar to above exercise, we need to check locally, but compactness is necessary here. Over an open subset U_1 , locally we have the Hermitian metric corresponding to $L' \otimes L^m$ looking like

$$\left(\frac{\partial^2\varphi_{L'}}{\partial z^i\partial\bar{z}^j} + m \cdot \frac{\partial^2\varphi_L}{\partial z^i\partial\bar{z}^j}\right)$$

So we can choose suffices large N_1 such that $M \otimes L^{\otimes N_1}$ is positive on U . Since X is compact, we can take a finite open covering $\{U_i\}$ of X and choose the largest N_i to be N we desired. \square

12.2. Positivity of vector bundles. Let (E, h) be a Hermitian holomorphic vector bundle of rank r over a complex n -manifold X with Chern connection ∇ . In local frame, its Chern curvature is given by

$$\Theta_h = \Theta_{i\bar{j}\alpha\bar{\beta}}^\beta dz^i \wedge d\bar{z}^j \otimes e^\alpha \otimes e_\beta$$

Definition 12.2.1 (positivity).

- (1) (E, h) is said to be Griffiths positive, if for any non-zero $(u^i) \in \mathbb{C}^n$ and $(v^\alpha) \in \mathbb{C}^r$

$$\Theta_{i\bar{j}\alpha\bar{\beta}} u^i \bar{u}^j v^\alpha \bar{v}^\beta > 0$$

- (2) (E, h) is said to be Nakano positive, if for any non-zero matrix $(u^{i\alpha})$

$$\Theta_{i\bar{j}\alpha\bar{\beta}} u^{i\alpha} \bar{u}^{j\beta} > 0$$

- (3) (E, h) is said to be dual Nakano positive, if for any non-zero matrix $(u^{i\alpha})$

$$\Theta_{i\bar{j}\alpha\bar{\beta}} u^{i\beta} \bar{u}^{j\alpha} > 0$$

Remark 12.2.1. The semi-positivity and negativity can be defined in the same way.

Proposition 12.2.1.

- (1) If (E, h) is Nakano positive or dual Nakano positive, then (E, h) is Griffiths positive.
(2) (E, h) is Nakano positive if and only if (E^*, h^*) is dual Nakano negative.

Proof. For (1). If (E, h) is Nakano positive, then for non-zero $(u^i) \in \mathbb{C}^n$ and $(v^\alpha) \in \mathbb{C}^r$, consider matrix $(u^{i\alpha})$ defined by $u^{i\alpha} := u^i v^\alpha$, then

$$\Theta_{i\bar{j}\alpha\bar{\beta}} u^{i\alpha} \bar{u}^{j\beta} v^\alpha \bar{v}^\beta = \Theta_{i\bar{j}\alpha\bar{\beta}} u^{i\alpha} \bar{u}^{j\beta} > 0$$

The same argument holds for the case (E, h) is dual Nakano positive. (2) follows from the relation between curvature form of (E, h) and (E^*, h^*) , see Section 4.3 of [Liu23]. \square

Proposition 12.2.2. Let

$$0 \rightarrow S \rightarrow E \rightarrow Q \rightarrow 0$$

be an exact sequence of holomorphic vector bundles.

- (1) If (E, h) is Griffiths or dual Nakano positive, then so is (Q, h^Q) .
(2) If (E, h) is Griffiths or Nakano negative, then so is (S, h^S) .

Proposition 12.2.3. If Hermitian holomorphic vector bundle (E, h) is Griffiths positive, then $(E \otimes \det E, h \otimes \det h)$ is Nakano positive and dual Nakano positive.

13. VANISHING THEOREMS

13.1. Kodaira vanishing theorem.

Theorem 13.1.1 (Bochner-Kodaira-Nakano identity). Let (X, ω) be a compact Kähler manifold and (E, h) a Hermitian holomorphic vector bundle. Then

$$\Delta_{\bar{\partial}_E} = [\sqrt{-1}\Theta_h, \Lambda] + \Delta_{\partial_E}$$

Proof. Direct computation shows

$$\begin{aligned} \Delta_{\bar{\partial}_E} &= [\bar{\partial}_E, \bar{\partial}_E^*] \\ &= -\sqrt{-1}[\bar{\partial}_E, [\Lambda, \partial_E]] \\ &= -\sqrt{-1}[\Lambda, [\partial_E, \bar{\partial}_E]] - \sqrt{-1}[\partial_E, [\bar{\partial}_E, \Lambda]] \\ &= -\sqrt{-1}[\Lambda, \Theta_h] - \sqrt{-1}[\partial_E, \sqrt{-1}\partial_E^*] \\ &= [\sqrt{-1}\Theta_h, \Lambda] + \Delta_{\partial_E} \end{aligned}$$

□

Corollary 13.1.1 (Bochner-Kodaira-Nakano inequality). Let (X, ω) be a compact Kähler manifold and (E, h) a Hermitian holomorphic vector bundle. Then for $\alpha \in C^\infty(X, \Omega_X^{p,q} \otimes E)$, one has

$$([\sqrt{-1}\Theta_h, \Lambda]\alpha, \alpha) \leq (\Delta_{\bar{\partial}_E}\alpha, \alpha)$$

In particular, if α is $\Delta_{\bar{\partial}_E}$ -harmonic, then $([\sqrt{-1}\Theta_h, \Lambda]\alpha, \alpha) \leq 0$.

Proof. Direct computation shows

$$\begin{aligned} (\Delta_{\bar{\partial}_E}\alpha, \alpha) - ([\sqrt{-1}\Theta_h, \Lambda]\alpha, \alpha) &= (\Delta_{\partial_E}\alpha, \alpha) \\ &= \|\partial_E\alpha\|^2 + \|\partial_E^*\alpha\|^2 \geq 0 \end{aligned}$$

□

Corollary 13.1.2. Let X be a complex manifold. If Hermitian holomorphic vector bundle (E, h) is Griffiths positive, then $(\det E, \det h)$ is a positive holomorphic line bundle. In particular, X is Kähler.

Theorem 13.1.2 (Kodaira-Akizuki-Nakano vanishing). Let X be a compact n -manifold, (L, h) a positive Hermitian holomorphic line bundle. Then

$$H^{p,q}(X, L) = 0$$

for $p + q > n$.

Proof. Let X be endowed with the Kähler metric ω given by Chern curvature of L . Then there is an isomorphism $H^{p,q}(X, L) \cong \mathcal{H}^{p,q}(X, L)$. For $\alpha \in \mathcal{H}^{p,q}(X, L)$, by Corollary 13.1.1 one has

$$[\sqrt{-1}\Theta_h, \Lambda]\alpha \leq 0.$$

On the other hand,

$$([\sqrt{-1}\Theta_h, \Lambda]\alpha, \alpha) = 2\pi(p + q - n)\|\alpha\|^2 \geq 0.$$

Thus if $p + q > n$, one has $\alpha = 0$. This completes the proof. \square

Corollary 13.1.3 (Kodaira vanishing). Let X be a compact n -manifold and (L, h) be a positive holomorphic line bundle over X . Then

$$H^q(X, K_X \otimes L) = 0$$

for $q > 0$.

Proof. Just note that

$$H^q(X, K_X \otimes L) = H^{n,q}(X, L)$$

\square

Corollary 13.1.4. Let (X, ω) be a compact Kähler n -manifold. If (L, h) is a semi-positive line bundle and $\text{rk } \Theta_h \geq k$, then

$$H^{p,q}(X, L) = 0$$

for $p + q \geq 2n - k + 1$.

Exercise 13.1.1. Compute all $H^q(\mathbb{C}\mathbb{P}^n, \mathcal{O}_{\mathbb{C}\mathbb{P}^n}(k))$ for all k, q .

Definition 13.1.1 (Fano). A Fano manifold is a compact Kähler manifold with positive anti-canonical bundle $K_X^* = \det TX$.

Proposition 13.1.1. Let X be a Fano manifold, then

$$H^q(X, \mathcal{O}_X) = 0$$

for all $q > 0$.

Proof. Note that $\mathcal{O}_X = K_X \otimes K_X^*$. \square

Theorem 13.1.3 (Serre vanishing). Let X be a compact complex n -manifold and (L, h) be a positive holomorphic line bundle over X . For any holomorphic vector bundle E on X , there exists a constant m_0 such that for all $m \geq m_0$

$$H^q(X, E \otimes L^{\otimes m}) = 0$$

for $q > 0$.

Proof. If X is endowed with Kähler metric ω given by Chern curvature of L and E is endowed with a Hermitian metric h , then $H^{p,q}(X, E \otimes L^{\otimes m}) \cong \mathcal{H}^{p,q}(X, E \otimes L^{\otimes m})$. For $\alpha \in \mathcal{H}^{p,q}(X, E \otimes L^{\otimes m})$, one has

$$\begin{aligned} ([\sqrt{-1}\Theta_h, \Lambda]\alpha, \alpha) + 2\pi m(p + q - n)\|\alpha\|^2 &\stackrel{(1)}{=} ([\sqrt{-1}\Theta_{E \otimes L^{\otimes m}}, \Lambda]\alpha, \alpha) \\ &\stackrel{(2)}{\leq} 0 \end{aligned}$$

where

(1) holds from $\Theta_{E \otimes L^{\otimes m}} = \Theta^E \otimes \text{id} + m(\text{id} \otimes \Theta^L)$.

(2) holds from Corollary 13.1.1, that is Bochner-Kodaira-Nakano inequality.

On the other hand, Cauchy inequality implies that

$$([\sqrt{-1}\Theta_h, \Lambda]\alpha, \alpha) \geq -C\|\alpha\|^2$$

where constant C is the norm of $[\sqrt{-1}\Theta_h, \Lambda]$. So if we have $2\pi m(p + q - n) - C > 0$, the argument in proof of Kodaira vanishing theorem implies $\alpha = 0$. Consider $p = n, q > 0, m_0 \geq \frac{C}{2\pi}$, then for all $m \geq m_0$ and $q > 0$, one has

$$H^{n,q}(X, E \otimes L^{\otimes m}) = 0$$

that is to say $H^q(X, K_X \otimes E \otimes L^{\otimes m}) = 0$. So in order to show $H^q(X, E \otimes L^{\otimes m}) = 0$, it suffices to consider $K_X^* \otimes E$ at beginning, and then we will obtain

$$H^q(X, K_X \otimes K_X^* \otimes E \otimes L^{\otimes m}) = H^q(X, E \otimes L^{\otimes m}) = 0$$

This completes the proof. \square

13.2. Nakano vanishing.

Theorem 13.2.1 (Nakano vanishing). Let X be a compact complex manifold and (E, h) a Hermitian holomorphic vector bundle over X .

(1) If (E, h) is Nakano positive, then

$$H^{n,q}(X, E) = 0$$

for $q \geq 1$.

(2) If (E, h) is dual Nakano positive, then

$$H^{p,n}(X, E) = 0$$

for $p \geq 1$.

Proof. Here we only give the proof of the first one, the second can be derived from the same argument. Suppose $\{\frac{\partial}{\partial z^1}, \dots, \frac{\partial}{\partial z^n}\}$ is a holomorphic local frame which is orthonormal at point $p \in X$, and $\varphi \in H^{n,q}(X, E)$ which is locally written as $\varphi = \varphi_I^\alpha dz^1 \wedge \dots \wedge dz^n \wedge d\bar{z}^I \otimes e_\alpha$ where $|I| = q$. Direct computation shows

$$\begin{aligned} \langle \sqrt{-1}[\Theta_h, \Lambda]\varphi, \varphi \rangle &= \langle \sqrt{-1}\Theta_h \Lambda \varphi, \varphi \rangle \\ &\stackrel{(1)}{=} -\langle \Theta_{k\bar{l}\alpha}^\gamma dz^k \wedge d\bar{z}^l \sum_m \iota_m \iota_{\bar{m}} \varphi_I^\alpha dz^1 \wedge \dots \wedge dz^n \wedge d\bar{z}^I \otimes e_\gamma, \varphi_J^\beta dz^1 \wedge \dots \wedge dz^n \wedge d\bar{z}^J \otimes e_\beta \rangle \\ &= -\Theta_{k\bar{l}\alpha\bar{\beta}} \varphi_I^\alpha \overline{\varphi_J^\beta} \langle dz^k \wedge d\bar{z}^l \wedge \sum_m \iota_m \iota_{\bar{m}} dz^1 \wedge \dots \wedge dz^n \wedge d\bar{z}^I, dz^1 \wedge \dots \wedge dz^n \wedge d\bar{z}^J \rangle \\ &= \sum_k \Theta_{k\bar{l}\alpha\bar{\beta}} \varphi_I^\alpha \overline{\varphi_J^\beta} \langle d\bar{z}^l \wedge \iota_{\bar{k}} d\bar{z}^I, d\bar{z}^J \rangle \\ &= \sum_{I, J} \left\{ \Theta_{k\bar{l}\alpha\bar{\beta}} \left(\sum_{k \in I} \varphi_I^\alpha \right) \overline{\left(\sum_{l \in J} \varphi_J^\beta \right)} \right\} \\ &\stackrel{(2)}{\geq} 0 \end{aligned}$$

where

(1) holds from Proposition 8.1.4 since by Corollary 13.1.2 one has X is Kähler.

(2) holds from (E, h) is Nakano positive.

But by Corollary 13.1.1, that is Bochner-Kodaira-Nakano inequality, one has $\langle \sqrt{-1}[\Theta_h, \Lambda]\varphi, \varphi \rangle \leq 0$. This shows $\varphi = 0$ as before. \square

Corollary 13.2.1. Let X be a compact complex n -manifold with $n \geq 2$. Then there doesn't exist metric h such that (TX, h) is Nakano positive.

Proof. Suppose (TX, h) is Nakano positive, then

$$\begin{aligned} H^{1,1}(X) &= H^{0,1}(X, T^*X) \\ &\stackrel{(1)}{=} H^{n,n-1}(X, TX) \\ &\stackrel{(2)}{=} 0 \end{aligned}$$

where

(1) holds from Serre duality.

(2) holds from Nakano vanishing theorem.

However, by Proposition 12.2.1 and Corollary 13.1.2 one has X is a Kähler manifold, which leads to a contradiction since $H^{1,1}(X) \neq 0$ for a Kähler manifold. \square

Corollary 13.2.2. Let X be a compact complex n -manifold with $n \geq 2$. Then there doesn't exist metric h such that (T^*X, h) is dual Nakano positive.

Proof. Note that

$$H^{n-1,n}(X, T^*X) = H^{1,0}(X, TX) = H^0(X, \text{End}(TX)) \neq 0$$

\square

Example 13.2.1. If $n \geq 2$, $(\mathbb{C}\mathbb{P}^n, \omega_{\text{FS}})$ is dual Nakano positive and Nakano semi-positive, but not Nakano positive.

Proof. By Theorem 7.4.1 one has the curvature of Fubini-Study metric is

$$\Theta_{i\bar{j}k\bar{l}} = h_{i\bar{j}}h_{k\bar{l}} + h_{i\bar{l}}h_{k\bar{j}}$$

For $p \in X$, if we consider normal coordinate at p , then for any non-zero matrix (u^{ij}) , a direct computation shows

$$\begin{aligned} \Theta_{i\bar{j}k\bar{l}}u^{il}\bar{u}^{jk} &= h_{i\bar{j}}h_{k\bar{l}}u^{il}\bar{u}^{jk} + h_{i\bar{l}}h_{k\bar{j}}u^{il}\bar{u}^{jk} \\ &= \sum_{i,j} |u^{ij}|^2 + u^{ii}\bar{u}^{jj} \\ &= \sum_{i \neq j} |u^{ij}|^2 + \frac{1}{2} \sum_{i,j} |u^{ii} + u^{jj}|^2 \\ &> 0 \end{aligned}$$

which implies $(\mathbb{C}\mathbb{P}^n, \omega_{FS})$ is dual Nakano positive.

$$\begin{aligned} \Theta_{i\bar{j}k\bar{l}} u^{ik} \bar{u}^{jl} &= h_{i\bar{j}} h_{k\bar{l}} u^{ik} \bar{u}^{jl} + h_{i\bar{l}} h_{k\bar{j}} u^{ik} \bar{u}^{jl} \\ &= \sum_{i,j} |u^{ij}|^2 + u^{ij} \bar{u}^{ji} \\ &= \frac{1}{2} \sum_{i,j} |u^{ij} + u^{ji}|^2 \\ &\geq 0 \end{aligned}$$

which implies $(\mathbb{C}\mathbb{P}^n, \omega_{FS})$ is Nakano semi-positive, and it's not Nakano positive by Corollary 13.2.1. \square

13.3. Griffiths vanishing.

Theorem 13.3.1 (Griffiths vanishing). If (E, h) is Griffiths positive, then


- (1) $H^{n,n}(X, E) = 0$.
- (2) $H^{p,q}(X, E) = 0$ for $p + q \geq n + \text{rk } E$.

14. KODAIRA EMBEDDING AND CHOW THEOREM

14.1. **Blow-up.** In this section we will introduce a technical tool we need in the proof of Kodaira embedding, which is blow-up at a point.

Definition 14.1.1 (blow-up). $U \times \mathbb{C}\mathbb{P}^{n-1} \supset \tilde{U} := \{((x_1, \dots, x_n), (y_1 : \dots : y_n)) \mid x_i y_j = x_j y_i\}$

Remark 14.1.1. The most vivid way to understand blow-up is to consider the fibres of projection $\tilde{U} \rightarrow U$: If $x \neq 0$, then the fibre of x is just a point since the ratio of y_i is uniquely determined. But for $x = 0$, there is no restriction for y_i , so you get the whole projective space $\mathbb{C}\mathbb{P}^{n-1}$. Thus as you can imagine, there is nothing happening except the origin, sounds like a boom. For example, the following figure shows the case of $n = 2$



Lemma 14.1.1. $\tilde{U} \subseteq U \times \mathbb{C}\mathbb{P}^{n-1}$ is a submanifold of dimension n .

Since blow-up is a local operation, so it can be done on a complex manifold. If X is a complex manifold with dimension n with $x \in X$, and $\{U_i\}$ is an open covering such that $x \in U_1$ and $x \notin U_i, i \neq 1$, then we can show that $\tilde{U}_1 \cup (\bigcup_{i \neq 1} U_i)$ glue together a new complex manifold with dimension n . This is called blow-up of X at point x , and it's denoted by \tilde{X} . Similarly there is a natural projection $\pi: \tilde{X} \rightarrow X$ and $\pi^{-1}(x)$ is biholomorphic to $\mathbb{C}\mathbb{P}^{n-1}$, which is called exceptional divisor and denote it by E .

Exercise 14.1.1. If X is compact Kähler manifold, then \tilde{X} is also a compact Kähler manifold.

Exercise 14.1.2. The idea sheaf of exceptional divisor $\mathcal{I}_E \cong \mathcal{O}_{\tilde{X}}(E)^*$

Proposition 14.1.1. The canonical bundle $K_{\tilde{X}}$ of the blow-up \tilde{X} is isomorphic to $\pi^* K_X \otimes \mathcal{O}_{\tilde{X}}(E)^{\otimes n-1}$.

Corollary 14.1.1. For the exceptional divisor $E = \mathbb{C}\mathbb{P}^n \subseteq \tilde{X} \rightarrow X$ one has $\mathcal{O}(E)|_E \cong \mathcal{O}_{\mathbb{C}\mathbb{P}^{n-1}}(-1)$.

Proof. The adjunction formula implies

$$K_E \cong (K_{\tilde{X}} \otimes \mathcal{O}_{\tilde{X}}(E))|_E$$

Then by Proposition 14.1.1 and $K_E = \mathcal{O}_{\mathbb{C}\mathbb{P}^{n-1}}(-n)$ one has

$$\mathcal{O}_{\mathbb{C}\mathbb{P}^{n-1}}(-n) \cong (\pi^* K_X \otimes \mathcal{O}_{\tilde{X}}(E)^{\otimes n})|_E$$

Note that E is a fibre of π , so $\pi^* K_X|_E$ is trivial. Thus

$$\mathcal{O}_{\mathbb{C}\mathbb{P}^{n-1}}(-n) \cong \mathcal{O}_{\tilde{X}}(E)^{\otimes n}|_E$$

On the other hand, the only possible line bundles on $\mathbb{C}\mathbb{P}^{n-1}$ take the form $\mathcal{O}_{\mathbb{C}\mathbb{P}^{n-1}}(k)$, $k \in \mathbb{Z}$. Thus $\mathcal{O}_{\tilde{X}}(E)|_E \cong \mathcal{O}_{\mathbb{C}\mathbb{P}^{n-1}}(-1)$. \square

The main reason we need blow-up here is that the sections after blow-up is the “same” as the one before.

Proposition 14.1.2. For a holomorphic line bundle L , one has

$$H^0(X, L^{\otimes m}) = H^0(\tilde{X}, \pi^* L^{\otimes m})$$

holds for arbitrary $m \in \mathbb{Z}_{\geq 0}$.

Proof. If X is one-dimensional, then $\pi: \tilde{X} \rightarrow X$ is an isomorphism and thus $H^0(X, L^{\otimes m}) = H^0(\tilde{X}, \pi^* L^{\otimes m})$.

Now let’s consider the case $\dim_{\mathbb{C}} X \geq 2$. Given an element $s \in H^0(X, L^{\otimes m})$, we can get an element in $H^0(\tilde{X}, \pi^* L^{\otimes m})$ by composing projection π . Conversely, for $\tilde{s} \in H^0(\tilde{X}, \pi^* L^{\otimes m})$, it can be restricted to $\tilde{X} \setminus E = X \setminus \{x\}$, and then extended to a global section of $L^{\otimes m}$ by Hartogs theorem. \square

14.2. Kodaira embedding.

Theorem 14.2.1 (Kodaira embedding). Let X be a compact complex manifold. The following statements are equivalent:

- (1) There exists a holomorphic embedding $\varphi: X \hookrightarrow \mathbb{C}\mathbb{P}^N$.
- (2) There exists an integral Kähler form ω on X , that is, $[\omega] \in \text{im}\{H^2(X, \mathbb{Z}) \rightarrow H^2(X, \mathbb{R})\}$
- (3) There exists a positive holomorphic line bundle on X .

Remark 14.2.1. (1) clearly implies (2), and (2) implies (3) is Lefschetz (1, 1)-theorem, so the heart of the proof is (3) to (1).

Sketch. Use holomorphic global sections $H^0(X, L^{\otimes m})$ for sufficiently large m to construct $\varphi: X \hookrightarrow \mathbb{C}\mathbb{P}^N$. We need to show the following three things:

- (1) For sufficiently large m , $L^{\otimes m}$ is globally generated, which means for all $x \in X$, there exists a global section $s \in H^0(X, L^{\otimes m})$ such that $s(x) \neq 0$. Then for all $x \in X$, $H_x = \{s \in H^0(X, L^{\otimes m}) \mid s(x) = 0\}$ is a hypersurface. Thus we get a holomorphic map $\varphi: X \rightarrow \mathbb{C}\mathbb{P}(H^0(X, L^{\otimes m})^*)$, defined by $x \mapsto H_x$. Indeed since any hypersurface in $H^0(X, L^{\otimes m})$ is a

line in $H^0(X, L^{\otimes m})^*$, that is an element in $\mathbb{C}\mathbb{P}(H^0(X, L^{\otimes m})^*)$. And you will find φ is holomorphic since we're using holomorphic sections.

- (2) For more sufficiently large¹⁴ m , $L^{\otimes m}$ separates points, that is for all $x, y \in X$, there exists $s \in H^0(X, L^{\otimes m})$ such that $s(x) \neq 0, s(y) = 0$. Thus in this case our φ is injective.
- (3) For more more sufficiently large m , $L^{\otimes m}$ separates tangent vectors, that is, for all $x \in X, u \in T_{X,x}$ there exists $s \in H^0(X, L^{\otimes m})$ such that $s(x) = 0$ and $ds(u) \neq 0$. Thus in this case our φ is an immersion, together with X is compact we have φ is an embedding.

Remark 14.2.2. We can also describe φ more explicitly. Locally around x_0 , choose a basis s_0, \dots, s_N of $H^0(X, L^{\otimes m})$ such that $s_0(x_0) \neq 0$. Then there exists a neighborhood U of x_0 such that $s_0(x) \neq 0$ for all $x \in U$. Then

$$\frac{s_1}{s_0}, \dots, \frac{s_N}{s_0} \in H^0(U, \mathcal{O}_U)$$

So we can define

$$\begin{aligned} \varphi|_U: U &\rightarrow \mathbb{C}\mathbb{P}^N \\ x &\mapsto (1, \frac{s_1}{s_0}(x), \dots, \frac{s_N}{s_0}(x)) \end{aligned}$$

And you can check it's same as what we have defined without choosing a basis.

Here we only give a sketch proof of the first statement, the proofs for second and third are similar, but more complicated.

We want to detect the value of a section at a point is zero or not. Sheaves give us a good way to describe such thing. For $x \in X$, consider ideal sheaf of x

$$\mathcal{I}_x = \{s \in \mathcal{O}_X \mid s(x) = 0\} \subseteq \mathcal{O}_X$$

Then sections we are searching for is $H^0(X, \mathcal{I}_x \otimes L^{\otimes m})$. For computation, we have exact sequence of sheaf

$$0 \rightarrow \mathcal{I}_x \otimes L^{\otimes m} \rightarrow L^{\otimes m} \rightarrow L^{\otimes m}|_x \rightarrow 0$$

And using Čech cohomology we can derive a long exact sequence

$$0 \rightarrow H^0(X, \mathcal{I}_x \otimes L^{\otimes m}) \rightarrow H^0(X, L^{\otimes m}) \rightarrow \mathbb{C} \rightarrow \check{H}^1(X, \mathcal{I}_x \otimes L^{\otimes m}) \rightarrow \dots$$

Our goal is to show $H^0(X, \mathcal{I}_x \otimes L^{\otimes m}) \neq 0$. If $\check{H}^1(X, \mathcal{I}_x \otimes L^{\otimes m}) = 0$ for sufficiently large m , then we can get desired result.

For Čech cohomology we know a little, but we know quite a lot for Dolbeault cohomology. So an ideal is to turn idea sheaf into a line bundle and use Dolbeault cohomology to compute.

Similarly, we have

$$H^0(X, \mathcal{I}_x \otimes L^{\otimes m}) = H^0(\tilde{X}, \mathcal{I}_E \otimes \pi^* L^{\otimes m}) = H^0(\tilde{X}, \mathcal{O}_{\tilde{X}}(E)^* \otimes \pi^* L^{\otimes m})$$

And that's why blow-up works since $\mathcal{I}_x \otimes L^{\otimes m}$ is just a sheaf, and it's a little difficult for us to compute the cohomology of sheaf, but after blow-up,

¹⁴Larger than m is step one.

we make it to a line bundle $\mathcal{O}_{\tilde{X}}(E)^* \otimes \pi^* L^{\otimes m}$, and Dolbeault cohomology comes into its place.

Consider the following short exact sequence of sheaves on \tilde{X} :

$$0 \rightarrow \mathcal{I}_E \otimes \pi^* L^{\otimes m} \rightarrow \pi^* L^{\otimes m} \rightarrow \pi^* L^{\otimes m}|_E \rightarrow 0$$

So we get a long exact sequence

$$0 \rightarrow H^0(\tilde{X}, \mathcal{I}_E \otimes \pi^* L^{\otimes m}) \rightarrow H^0(\tilde{X}, \pi^* L^{\otimes m}) \rightarrow \mathbb{C} \rightarrow \check{H}^1(\tilde{X}, \mathcal{I}_E \otimes \pi^* L^{\otimes m}) \rightarrow \dots$$

But

$$\check{H}^1(\tilde{X}, \mathcal{I}_E \otimes \pi^* L^{\otimes m}) = H_{Dol}^1(\tilde{H}, \mathcal{I}_E \otimes \pi^* L^{\otimes m})$$

Claim $H_{Dol}^1(\tilde{H}, \mathcal{I}_E \otimes \pi^* L^{\otimes m}) = 0$, when m is sufficiently large. Indeed, note that

$$\mathcal{I}_E \otimes \pi^* L^{\otimes m} \cong K_{\tilde{X}} \otimes \{\mathcal{O}_{\tilde{X}}(E)^{* \otimes n} \otimes \pi^*(K_X^* \otimes L^{\otimes m})\}$$

So by Kodaira vanishing, it suffices to show the following line bundle is positive when m is sufficiently large:

$$\mathcal{O}_{\tilde{X}}(E)^{* \otimes n} \otimes \pi^*(K_X^* \otimes L^{\otimes m})$$

In fact, $K_X^* \otimes L^{\otimes m}$ will be positive on X when m is sufficiently large. But when we pull it back something bad may happen since $\pi^*(K_X^* \otimes L^{\otimes m})$ is positive except along E . However, $\mathcal{O}_{\tilde{X}}(E)^{* \otimes n}|_E = \mathcal{O}_{\mathbb{C}\mathbb{P}^{n-1}}(n)$, so two parts work together to give a positive line bundle. To be more explicit, take any Hermitian metric on $\mathcal{O}_{\tilde{X}}(E)^{* \otimes n}$ extending (Fubini-Study) $^{\otimes n}$, then its positive on E , but may not be positive otherwise. However we can choose m sufficiently large to offset its negative impact.

This completes the proof of first part of Kodaira embedding, for second and third, arguments are similar, but we need more blow-ups and things become complicated. \square

Corollary 14.2.1. Let (X, ω) be a compact Kähler manifold such that $H^{2,0}(X) = H^{0,2}(X) = 0$. Then X is a projective manifold.

Proof. Hodge decomposition implies that $H^{2,0}(X) = H^{0,2}(X) = 0$, and thus $H^2(X, \mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{C} = H^2(X, \mathbb{C}) = H^{1,1}(X)$. Let $[\alpha_1], \dots, [\alpha_n] \in H^2(X, \mathbb{Q})$ be a basis such that α_i is harmonic and of type $(1, 1)$. Since the Kähler form ω is real, harmonic¹⁵ and of type $(1, 1)$. Then

$$\omega = \sum_i \lambda_i \alpha_i, \quad \lambda_i \in \mathbb{R}$$

For $\mu_i \in \mathbb{Q}$ sufficiently close to λ_i , one still has $\sum_i \mu_i \alpha_i$ is positive. Thus $\sum_i \mu_i \alpha_i$ gives a Kähler form. Take N sufficiently large such that $[N \sum_i \mu_i \alpha_i] \in \text{im}(H^2(X, \mathbb{Z}) \rightarrow H^2(X, \mathbb{R}))$. Applying Kodaira embedding to complete the proof. \square

Corollary 14.2.2. Fano manifold is projective.

Proof. Since for Fano manifold, all $H^{0,p}(X) = 0, q > 0$. \square

¹⁵It's harmonic since $[\Delta_d, L] = 0$.

14.3. Chow's theorem. Since we already embed a compact complex manifold into projective space, there is no reason for us to avoid Chow's theorem. A wonderful theorem lies in the intersection of algebraic and analytic.

Theorem 14.3.1 (Chow). Every closed complex submanifold $X \subseteq \mathbb{C}\mathbb{P}^n$ is algebraic, that is, defined by polynomial equations.

Remark 14.3.1. Finally, for holomorphic line on compact complex manifolds: positive is equivalent to ample.

Although Chow's theorem can be derived from GAGA proved by Serre, in an elegant way using sheaf theory, here we give a sketch of a classical proof of Chow's theorem.

We need to show every closed complex submanifold X of $\mathbb{C}\mathbb{P}^n$ is algebraic, our ideal is to construct an analytic hypersurface in a Grassmannian manifold $Gr(r, n)$ with Plücker embedding $\varphi: Gr(r, n) \hookrightarrow \mathbb{C}\mathbb{P}(\bigwedge^r \mathbb{C}^n)$, and use some facts about it:

- (1) $\text{Pic}(Gr(r, n)) = \mathbb{Z} \cdot \varphi^* \mathcal{O}_{\mathbb{C}\mathbb{P}}(1)$
- (2) Every closed analytic hypersurface of $Gr(r, n)$ is algebraic.

If the analytic hypersurface W we construct in Grassmannian manifold can determine X algebraically, that is W is algebraic implies X is, then we complete the proof.

The philosophy here is to convert a submanifold with arbitrary codimension in $\mathbb{C}\mathbb{P}^n$ to a an hypersurface, the cost we pay is that we need to consider Grassmannian manifold rather than $\mathbb{C}\mathbb{P}^n$. But it do works!

Let's be more explicit:

Definition 14.3.1 (analytic subset). A closed analytic subset in $\mathbb{C}\mathbb{P}^n$ is a closed subset, locally defined by some holomorphic equations.

Remark 14.3.2. We can replace closed complex submanifold by closed analytic subset in Chow's theorem since we can not avoid singularity, and it doesn't matter in fact.

However, although we allow singularities, singularities won't be too much: Let $X \subseteq \mathbb{C}\mathbb{P}^n$ be an irreducible closed analytic subset of dimension r . Then there exists closed analytic subset $X_{\text{sing}} \subseteq X$ such that $X \setminus X_{\text{sing}}$ is smooth and dense. Furthermore, $X \setminus X_{\text{sing}}$ is a submanifold of $\mathbb{C}\mathbb{P}^n$ of dimension r .

Now fix X , an irreducible closed analytic subset of dimension r in $\mathbb{C}\mathbb{P}^n$. Let $V \in Gr(n-r, n+1)$, then $\mathbb{C}\mathbb{P}(V) \subseteq \mathbb{C}\mathbb{P}^n$ with dimension $n-r-1$. So as you can imagine, an object with dimension r and an object with dimension $n-r-1$ may fail to intersect with each other.

Let

$$W = \{V \in Gr(n-r, n+1) \mid \mathbb{C}\mathbb{P}(V) \cap X \neq \emptyset\}$$

Claim:

- (1) W is a closed analytic hypersurface of $Gr(n-r, n+1)$.
- (2) W determines X algebraically.

Here we give a sketch of proof of Claims:

For (1). Consider the following diagram

$$\begin{array}{ccc} \mathbb{C}\mathbb{P}(E) & \xrightarrow{q} & \mathbb{C}\mathbb{P}^n \supset X \\ \downarrow p & & \\ Gr(n-r, n+1) & & \end{array}$$

where $E \rightarrow Gr(n-r, n+1)$ is tautological bundle of Grassmannian manifold. Then we can write¹⁶

$$W = p(q^{-1}(X))$$

Since q is holomorphic and X is closed and analytic, then $q^{-1}(X)$ is also closed and analytic. But the difficulty is $p(q^{-1}(X))$ is also closed and analytic, and this holds from the following fact.

Theorem 14.3.2. p is holomorphic and proper¹⁷.

Now we show that W is a hypersurface, and that's just a computation for dimension: We already know the dimension of $Gr(n-r, n+1)$ is $(n-r)(n+1-(n-r)) = (n-r)(r+1)$, so we need to show the dimension of W is $(n-r)(r+1) - 1$.

First, let's consider the fibre of q : it consists of subspaces $V \subseteq \mathbb{C}^{n+1}$ of dimension $n-r$ containing a given line l , and that's another Grassmannian manifold $Gr(n-r-1, n)$, if we consider $V \mapsto V/l$, and its dimension is $(n-r-1)(r+1)$. So the dimension of $q^{-1}(X)$ is $r + (n-r-1)(r+1) = (n-r)(r+1) - 1 = \dim Gr(n-r, n+1) - 1$.

So we may desire the property of p is not too bad so that we will obtain $\dim p(q^{-1}(X)) = \dim q^{-1}(X)$ as we desired. It suffices to show that there exists a dense open subset $U \subseteq q^{-1}(X)$, such that $p|_U$ has finite fibres. In fact, we will show it's one to one correspondence.

Consider fibre of $p_X : q^{-1}(X) \rightarrow Gr(n-r, n+1)$ over given $V \in Gr(n-r, n+1)$, and that's $\mathbb{C}\mathbb{P}(V) \cap X$. So we may desire almost every V such that this intersection is just a point. It suffices to show that the complement of

$$\{(V, x) \in q^{-1}(X) \mid \mathbb{C}\mathbb{P}(V) \cap X \text{ has only one smooth point } x\}$$

is closed analytic of dimension less than $\dim q^{-1}(X)$. There are three cases:

- (1) $\mathbb{C}\mathbb{P}(V) \cap X$ contains $x \in X_{\text{sing}}$, singular locus of X . But $\dim_{\mathbb{C}} q^{-1}(X_{\text{sing}}) < \dim_{\mathbb{C}} q^{-1}(X)$.
- (2) $\mathbb{C}\mathbb{P}(V) \cap X$ has at least two points.
- (3) $\mathbb{C}\mathbb{P}(V) \cap X$ not transverse intersection at x .

Then $W = p(q^{-1}(X)) \subseteq Gr(n-r, n+1)$ is a closed analytic hypersurface.

For (2). Consider $q_W : p^{-1}(W) \rightarrow \mathbb{C}\mathbb{P}^n$. Claim

$$X = \{x \in \mathbb{C}\mathbb{P}^n \mid q_W^{-1}(x) = q^{-1}(x)\}$$

¹⁶Why?

¹⁷Proper means: For any $Y \subseteq \mathbb{C}\mathbb{P}(E)$ closed and analytic, then $p(Y)$ is closed and analytic.

And there are some equivalent descriptions:

$$\begin{aligned} q_W^{-1}(x) = q^{-1}(x) &\iff p^{-1}(W) \cap q^{-1}(x) = q^{-1}(x) \\ &\iff q^{-1}(x) \subseteq p^{-1}(W) \end{aligned}$$

Clearly, if $x \in X$, then $q^{-1}(x) \in p^{-1}(W)$ since $W = p(q^{-1}(X))$. For the other direction: we can translate it as: If $y \notin X$, then we need to find $V \in Gr(n-r, n+1)$ containing $l = \langle y \rangle$, but $\mathbb{C}\mathbb{P}(V) \cap X = \emptyset$.

To see this: Use projection from y , that is $\mathbb{C}\mathbb{P}^n \xrightarrow{\pi_y} \mathbb{C}\mathbb{P}^{n-1}$. Since $y \notin X$, then $\pi_y|_X$ is well-defined and has finite fibres. Note that $\mathbb{C}\mathbb{P}(V) \cap X \neq \emptyset$ if and only if $\mathbb{C}\mathbb{P}(V/\langle y \rangle) \cap \pi_y(X) = \emptyset$. From computation before, we know it's a condition for hypersurface. So it's easy to choose V we desire.

Conclusion (from analytic to algebraic): p and q is algebraic, and W is algebraic, so we obtain $p^{-1}(W)$ is also algebraic. So q_W is also algebraic. Thus X is algebraic.

15. MORE BOCHNER TECHNIQUES

15.1. Obstruction to holomorphic vector fields. In Riemannian geometry, there is the following theorem which shows there are obstructions of Killing vector fields or harmonic 1-forms.

Theorem 15.1.1. Let (M, g) be a compact Riemannian manifold.

- (1) If $\text{Ric}(g) < 0$, then there is no non-trivial Killing vector field.
- (2) If $\text{Ric}(g) > 0$, then $b_1(M) = 0$.

In the realm of complex geometry, analogous obstructions exist, drawing a parallel between Killing fields and holomorphic vector fields. However, it is important to note that not every Killing vector field is a holomorphic vector field.

Theorem 15.1.2. On a compact Kähler manifold, a Killing vector field X is holomorphic if and only if

- (1) $\text{div} X = 0$.
- (2) $\nabla^* X = 0$.

Lemma 15.1.1. Let (M, g) be a compact Riemannian manifold. A vector field X is Killing if and only if $\text{div} X = 0$ and $\nabla^* \nabla X = (\text{Ric}(X))^\sharp$.

Lemma 15.1.2. Let (M, ω) be a compact Kähler manifold. A vector field X is holomorphic if and only if $\nabla^* \nabla X = (\text{Ric}(X))^\sharp$.

Theorem 15.1.3. Let (M, ω) be a compact Kähler manifold with $\text{Ric}(\omega) < 0$. Then there is no non-trivial holomorphic vector field.

Proof. Let X be a non-trivial holomorphic vector field locally given by $X = X^i \frac{\partial}{\partial z^i}$ and $\omega = \sqrt{-1} h_{i\bar{j}} dz^i \wedge d\bar{z}^j$. If we define $u = |X|^2$, a direct computation shows

$$\begin{aligned} \sqrt{-1} \partial \bar{\partial} u &= \sqrt{-1} (\partial \{ \nabla^{0,1} X, X \} + \{ X, \nabla^{1,0} X \}) \\ &\stackrel{(1)}{=} \sqrt{-1} \partial \{ X, \nabla^{1,0} X \} \\ &= \sqrt{-1} (\{ \nabla^{1,0} X, \nabla^{1,0} X \} + \{ X, \nabla^{0,1} \nabla^{1,0} X \}) \\ &\stackrel{(2)}{=} \sqrt{-1} (\{ \nabla X, \nabla X \} + \{ X, \Theta_h X \}) \end{aligned}$$

where (1) and (2) holds from X is a holomorphic vector field. Note that

$$\begin{aligned} \text{tr}_\omega \sqrt{-1} \{ \nabla X, \nabla X \} &= \text{tr}_\omega (\sqrt{-1} \nabla_i X^j \overline{\nabla_k X^l} h_{j\bar{l}} dz^i \wedge d\bar{z}^k) \\ &= \nabla_i X^j \overline{\nabla_k X^l} h_{j\bar{l}} h^{i\bar{k}} \\ &= |\nabla X|^2 \\ \text{tr}_\omega \sqrt{-1} \{ X, \Theta_h X \} &= -\text{tr}_\omega \{ \sqrt{-1} \Theta_h X, X \} \\ &= -\text{tr}_\omega (\sqrt{-1} \Theta_{i\bar{j}\alpha}^\beta X^\alpha \overline{X^\gamma} h_{\beta\bar{\gamma}} dz^i \wedge d\bar{z}^j) \\ &= -h^{i\bar{j}} \Theta_{i\bar{j}\alpha\bar{\gamma}} X^\alpha \overline{X^\gamma} \\ &= -\text{Ric}(X, X) \end{aligned}$$

This shows

$$\operatorname{tr}_\omega \sqrt{-1} \partial \bar{\partial} u = |\nabla X|^2 - \operatorname{Ric}(X, X) > 0$$

If u obtains its maximum at some point $p \in M$, then $\operatorname{tr}_\omega \sqrt{-1} \partial \bar{\partial} u(p) \leq 0$, a contradiction. \square

Corollary 15.1.1. Let (M, ω) be a compact Kähler manifold. If $\operatorname{Ric}(\omega) \leq 0$, then the following statements are equivalent:

- (1) X is parallel.
- (2) X is Killing.
- (3) X is holomorphic.

Proof. As shown in Theorem 15.1.3, if $\operatorname{Ric}(\omega) \leq 0$, for a non-trivial holomorphic vector field X one has

$$\operatorname{tr}_\omega \sqrt{-1} \partial \bar{\partial} u = |\nabla X|^2 - \operatorname{Ric}(X, X) \geq 0$$

Then by the same argument one has $\nabla X = 0$, that is every holomorphic vector field is parallel, and it's clear every parallel vector field is Killing. \square

Corollary 15.1.2. Let (M, ω) be a compact complex manifold with negative holomorphic sectional curvature. Then there is no non-trivial holomorphic vector field.

Proof. Let X be a non-trivial holomorphic vector field locally given by $X = X^i \frac{\partial}{\partial z^i}$ and $u = |X|^2$. The same argument shows

$$\sqrt{-1} \partial \bar{\partial} u(X, X) = |\nabla_X X|^2 - \Theta_{i\bar{j}k\bar{l}} X^i \bar{X}^j X^k \bar{X}^l$$

Then maximum principle completes the proof. \square

15.2. Obstruction to holomorphic 1-form.

Theorem 15.2.1. Let (M, ω) be a compact Kähler manifold. If $\operatorname{Ric}(\omega) > 0$. Then there is no non-trivial holomorphic 1-form.

Corollary 15.2.1. Let (M, ω) be compact Kähler manifold and $\operatorname{Ric}(\omega) \geq 0$. Then any holomorphic 1-form is parallel.

15.3. Rigidity of complex projective space. The rigidity of complex projective space is a fascinating subject in complex geometry. It involves determining whether a geometric object satisfies certain conditions to be $\mathbb{C}\mathbb{P}^n$ or not, and it has produced numerous interesting results. Yau demonstrated that any Kähler manifold, which is homeomorphic to $\mathbb{C}\mathbb{P}^n$, is also biholomorphic to $\mathbb{C}\mathbb{P}^n$. He also resolved a conjecture put forth by Frankel in 1961, which stated that a compact Kähler manifold with positive bisectional curvature must be biholomorphic to $\mathbb{C}\mathbb{P}^n$. Additionally, there exists an even stronger result.

Theorem 15.3.1 ([FLW17]). Let (X, ω) be a compact Kähler manifold with positive orthogonal bisectional curvature. Then X is biholomorphic to $\mathbb{C}\mathbb{P}^n$.

In this section we try to use Bochner technique to show the following easier result.

Proposition 15.3.1. Let (X, ω) be a compact Kähler manifold with positive orthogonal bisectional curvature. Then $H^{1,1}(X) = \mathbb{C}$.

16. SCHWARZ LEMMAS

In this section, we will present the powerful formulas called Schwarz calculation, which extend the Bochner techniques discussed earlier.

16.1. Schwarz lemmas for holomorphic bisectional curvature.

Lemma 16.1.1. Let $f: (M, g) \rightarrow (N, h)$ be a holomorphic map between Hermitian manifolds and ∇ be the induced connection on $T^*X \otimes f^*(TN)$ by Chern connections. Then in the local holomorphic coordinates $\{z^i\}$ and $\{w^\alpha\}$ on M and N respectively, one has

$$\partial\bar{\partial}u = \{\nabla df, \nabla df\} + \left((\Theta_g)_{i\bar{j}k\bar{l}} g^{k\bar{q}} g^{p\bar{l}} h_{\alpha\bar{\beta}} f_p^\alpha \overline{f_q^\beta} - (\Theta_h)_{\alpha\bar{\beta}\gamma\bar{\delta}} (f_i^\alpha \overline{f_j^\beta}) (g^{p\bar{q}} f_p^\gamma \overline{f_q^\delta}) \right) dz^i \wedge d\bar{z}^j$$

and

$$\mathrm{tr}_\omega \sqrt{-1} \partial\bar{\partial}u = |\nabla df|^2 + (g^{i\bar{j}} (\Theta_g)_{i\bar{j}k\bar{l}}) g^{k\bar{q}} g^{p\bar{l}} h_{\alpha\bar{\beta}} f_p^\alpha \overline{f_q^\beta} - (\Theta_h)_{\alpha\bar{\beta}\gamma\bar{\delta}} (g^{i\bar{j}} f_i^\alpha \overline{f_j^\beta}) (g^{p\bar{q}} f_p^\gamma \overline{f_q^\delta})$$

where $u = \mathrm{tr}_g(f^*\omega_h) = |\nabla f|^2$, $f = f_i^\alpha dz^i \otimes e_\alpha$ and $e_\alpha = f^*\left(\frac{\partial}{\partial w^\alpha}\right)$. Furthermore, one has

$$\mathrm{tr}_\omega \sqrt{-1} \partial\bar{\partial} \log u \geq \frac{1}{u} \left((g^{i\bar{j}} (\Theta_g)_{i\bar{j}k\bar{l}}) g^{k\bar{q}} g^{p\bar{l}} h_{\alpha\bar{\beta}} f_p^\alpha \overline{f_q^\beta} - (\Theta_h)_{\alpha\bar{\beta}\gamma\bar{\delta}} (g^{i\bar{j}} f_i^\alpha \overline{f_j^\beta}) (g^{p\bar{q}} f_p^\gamma \overline{f_q^\delta}) \right)$$

holds outside the set of critical points of f .

Corollary 16.1.1. Let g, h be two Hermitian metrics on a complex manifold M . Then

$$\mathrm{tr}_g \sqrt{-1} \partial\bar{\partial} \log \mathrm{tr}_g \omega_h \geq \frac{1}{\mathrm{tr}_g \omega_h} \left((g^{i\bar{j}} (\Theta_g)_{i\bar{j}k\bar{l}}) g^{k\bar{q}} g^{p\bar{l}} h_{p\bar{q}} - (\Theta_h)_{i\bar{j}p\bar{q}} g^{i\bar{j}} g^{p\bar{q}} \right)$$

and

$$\mathrm{tr}_h \sqrt{-1} \partial\bar{\partial} \log \mathrm{tr}_g \omega_h \geq \frac{1}{\mathrm{tr}_g \omega_h} \left((h^{i\bar{j}} (\Theta_g)_{i\bar{j}k\bar{l}}) g^{k\bar{q}} g^{p\bar{l}} h_{p\bar{q}} - (h^{i\bar{j}} (\Theta_h)_{i\bar{j}p\bar{q}}) g^{p\bar{q}} \right)$$

Corollary 16.1.2. Let $f: (M, g) \rightarrow (N, h)$ be a holomorphic map between Hermitian manifolds. Suppose

- (1) the second Chern-Ricci curvature $\mathrm{Ric}^{(2)}(\omega_g) \geq a\omega_g$ for some a ;
- (2) the holomorphic bisectional curvature (N, h) is bounded from above by b , that is

$$\Theta_h(w, \bar{w}, v, \bar{v}) \leq b|w|^2|v|^2$$

Then

$$\mathrm{tr}_g \sqrt{-1} \partial\bar{\partial}u \geq au - bu^2$$

where $u = \mathrm{tr}_g(f^*\omega_h) = |\nabla f|^2$. Furthermore, outside the set of critical points of f , one has

$$\mathrm{tr}_g \sqrt{-1} \partial\bar{\partial} \log u \geq a - bu$$

Proof. Note that $\text{Ric}^{(2)}(\omega_g) \geq a\omega_g$ implies $(\Theta_g)_{i\bar{j}k\bar{l}}g^{k\bar{l}} \geq ag_{i\bar{j}}$. Thus

$$\begin{aligned} (g^{i\bar{j}}(\Theta_g)_{i\bar{j}k\bar{l}})g^{k\bar{q}}g^{p\bar{l}}h_{\alpha\bar{\beta}}f_p^\alpha\overline{f_q^\beta} &= g^{i\bar{j}}(\Theta_g)_{i\bar{j}k\bar{l}}\frac{g^{k\bar{l}}g_{k\bar{l}}}{n}g^{k\bar{q}}g^{p\bar{l}}h_{\alpha\bar{\beta}}f_p^\alpha\overline{f_q^\beta} \\ &\geq \frac{ag^{i\bar{j}}g_{i\bar{j}}}{n}g_{k\bar{l}}g^{k\bar{q}}g^{p\bar{l}}h_{\alpha\bar{\beta}}f_p^\alpha\overline{f_q^\beta} \\ &= ag^{p\bar{l}}f_p^\alpha\overline{f_l^\beta} \\ &= au \end{aligned}$$

On the other hand, by using normal coordinate on M it's easy to see

$$(\Theta_h)_{\alpha\bar{\beta}\gamma\bar{\delta}}(g^{i\bar{j}}f_i^\alpha\overline{f_j^\beta})(g^{p\bar{q}}f_p^\gamma\overline{f_q^\delta}) \leq b(f_i^\alpha\overline{f_j^\beta}g^{i\bar{j}}h_{\alpha\bar{\beta}})(f_p^\gamma\overline{f_q^\delta}g^{p\bar{q}}h_{\gamma\bar{\delta}}) = bu^2$$

Then by Lemma 16.1.1 this completes the proof. \square

Lemma 16.1.2 ([Yau75]). Let (M, g) be a complete Riemannian manifold with Ricci curvature bounded from below and $f \in C^2(M, \mathbb{R})$ be bounded from above. Then there exists $\{p_k\} \subseteq M$ such that

- (1) $\lim_k |\nabla f(p_k)| = 0$.
- (2) $\limsup_k \Delta f(p_k) \leq 0$.
- (3) $\lim_k f(p_k) = \sup f$.

Theorem 16.1.1 ([Yau78a]). Let (M, ω_g) be a complete Kähler manifold with $\text{Ric}(\omega_g) \geq a\omega_g$ and (N, ω_h) be a Hermitian manifold with holomorphic bisectional curvature $\leq b < 0$. If $f: M \rightarrow N$ is a non-constant holomorphic map, then $a < 0$ and

$$f^*\omega_h \leq \frac{a}{b}\omega_g$$

Proof. Let Δ_g denotes the operator $\text{tr}_g \sqrt{-1}\partial\bar{\partial}$. By Corollary 16.1.2 one has

$$\Delta_g u \geq au - bu^2$$

where $u = \text{tr}_g(f^*\omega_h)$. It suffices to show $\sup_M u \leq a/b$. Now let's consider the following two cases:

- (1) If $\sup_M u < \infty$, then by Lemma 16.1.2 there exists a sequence $\{p_k\}$ such that $\limsup \Delta u(p_k) \leq 0$, and $\lim u(p_k) = \sup_M u$ and so

$$\Delta_g u(p_k) \geq au(p_k) - bu^2(p_k)$$

By taking \limsup , we deduce that

$$0 \geq a \sup_M u - b(\sup_M u)^2$$

Since $\sup_M u > 0$ and $b < 0$, one has $a < 0$ and $\sup_M u \leq a/b$.

- (2) If $\sup_M u = \infty$, consider

$$v = \frac{1}{\sqrt{u+c}}$$

for some constant $c > 0$. Direct computation shows

$$\begin{aligned}\Delta_g v &= -\frac{1}{2}v^3 \Delta_g u + \frac{3}{v}|\nabla v|^2 \\ \Delta_g u &= \frac{2}{v^3}(\frac{3}{v}|\nabla v|^2 - \Delta_g v) \geq au - bu^2\end{aligned}$$

that is

$$6|\nabla v|^2 - 2v\Delta_g v \geq auv^4 - bu^2v^4$$

By Lemma 16.1.2 again there exists a sequence $\{p_k\}$ such that $\lim_k(-v)(p_k) = \sup_M(-v) = 0$ and $\nabla v(p_k) = 0$ and

$$\limsup_k \Delta_g v(p_k) \leq 0$$

Then

$$0 \geq \limsup(6|\nabla v|^2 - 2v\Delta_g v)(p_k) \geq \limsup(auv^4 - bu^2v^4)(p_k) = -b$$

which is a contradiction to $b > 0$.

In a word, only the first case will happen, and this completes the proof. \square

Corollary 16.1.3. Let $f: (\mathbb{D}, \omega) \rightarrow (\mathbb{D}, \omega)$ be a holomorphic map between unit disk with Poincaré metric. Then

$$f^*(\omega) \leq \omega$$

Proof. It's clear since unit disk with Poincaré metric has constant holomorphic bisectional curvature -1 and $\text{Ric}(\omega) = -\omega$. \square

Corollary 16.1.4. Let $f: (M, g) \rightarrow (N, h)$ be a holomorphic map between two Hermitian manifolds. If

- (1) (M, g) is complete Kähler with $\text{Ric}(g) \geq 0$.
- (2) (N, h) has negative holomorphic bisectional curvature.

Then f is constant.

Proof. Suppose the holomorphic bisectional curvature of N is bounded above by $b < 0$. If there exists a non-constant holomorphic map from (M, g) to (N, h) , then it contradicts to $\text{Ric}(\omega_g) \geq 0$ since by Theorem 16.1.1 one has if $\text{Ric}(\omega_g) \geq a\omega_g$, then $a < 0$. \square

Corollary 16.1.5. Let $f: (M, g) \rightarrow (N, h)$ be a holomorphic map between two Hermitian manifolds such that M is compact with $\text{Ric}^{(2)}(\omega_g) \geq 0$ and N has non-positive holomorphic bisectional curvature. If one of the following statements holds, then f is a constant.

- (1) $\text{Ric}^{(2)}(\omega_g) > 0$ at some point.
- (2) N has negative holomorphic bisectional curvature at some point.

Proof. Let ω_G be a Gauduchon metric of M and $u = \text{tr}_g(f^*\omega_h)$. By Lemma 16.1.1 the integration

$$\int_M \text{tr}_\omega \sqrt{-1} \partial \bar{\partial} u \cdot \omega_G^n = 0$$

implies $\nabla df = 0$ and

$$(g^{i\bar{j}}(\Theta_g)_{i\bar{j}k\bar{l}})g^{k\bar{q}}g^{p\bar{l}}h_{\alpha\bar{\beta}}f_p^\alpha\overline{f_q^\beta} = (\Theta_h)_{\alpha\bar{\beta}\gamma\bar{\delta}}(g^{i\bar{j}}f_i^\alpha\overline{f_j^\beta})(g^{p\bar{q}}f_p^\gamma\overline{f_q^\delta}) = 0$$

Hence $|df|$ is a constant, and it's clear from the second equation that (1) or (2) implies $df = 0$, that is f is a constant. \square

16.2. Schwarz lemmas for holomorphic sectional curvature.

Lemma 16.2.1. There is the following identity

$$\int_{\mathbb{C}\mathbb{P}^{n-1}} \frac{\xi^i \bar{\xi}^j \xi^k \bar{\xi}^l}{|\xi|^4} \omega_{FS}^{n-1} = \frac{\delta_{ij}\delta_{kl} + \delta_{il}\delta_{kj}}{n(n+1)}$$

where $[\xi^1 : \dots : \xi^n]$ are homogenous coordinates on $\mathbb{C}\mathbb{P}^{n-1}$ and ω_{FS} is the Fubini-Study metric.

Theorem 16.2.1. Let (M, g) be a Hermitian n -manifold, (N, h) be a Kähler n -manifold and $f: (M, g) \rightarrow (N, h)$ be a non-constant holomorphic map. Suppose that

- (1) $\text{Ric}^{(2)}(g) \geq -\lambda\omega_g + \mu f^*\omega_h$ for continuous functions λ, μ with $\mu \geq 0$.
- (2) holomorphic sectional curvature of h is bounded from above by a continuous functions $-\kappa \leq 0$.

Then

$$\text{tr}_g \sqrt{-1}\partial\bar{\partial}u \geq -\lambda u + \left(\frac{(r+1)f^*\kappa}{2r} + \frac{\mu}{n} \right) u^2$$

where r is maximal rank of df and $u = \text{tr}_g(f^*\omega_h)$. Furthermore, outside the critical points of f one has

$$\text{tr}_g \sqrt{-1}\partial\bar{\partial} \log u \geq -\lambda + \left(\frac{(r+1)f^*\kappa}{2r} + \frac{\mu}{n} \right) u$$

Proof. Let Δ_g denote the operator $= \text{tr}_g \sqrt{-1}\partial\bar{\partial}$. Then as computation in Lemma 16.1.1 one has

$$\Delta_g u = |\nabla df|^2 + \underbrace{(g^{i\bar{j}}(\Theta_g)_{i\bar{j}k\bar{l}})g^{k\bar{q}}g^{p\bar{l}}h_{\alpha\bar{\beta}}f_p^\alpha\overline{f_q^\beta}}_I - \underbrace{(\Theta_h)_{\alpha\bar{\beta}\gamma\bar{\delta}}(g^{i\bar{j}}f_i^\alpha\overline{f_j^\beta})(g^{p\bar{q}}f_p^\gamma\overline{f_q^\delta})}_{\text{II}}$$

For part I , one has

$$\begin{aligned} I &\geq g^{i\bar{j}}(\Theta_h)_{i\bar{j}k\bar{l}} \frac{g^{k\bar{l}}g_{k\bar{l}}}{n} g^{k\bar{q}}g^{p\bar{l}}h_{\alpha\bar{\beta}}f_p^\alpha\overline{f_q^\beta} \\ &= (-\lambda g_{k\bar{l}} + \mu g^{i\bar{j}}g_{k\bar{l}}h_{\gamma\bar{\delta}}f_i^\gamma\overline{f_j^\delta})g^{k\bar{q}}g^{p\bar{l}}h_{\alpha\bar{\beta}}f_p^\alpha\overline{f_q^\beta} \\ &= (-\lambda g_{k\bar{l}} + \mu h_{\gamma\bar{\delta}}f_k^\gamma\overline{f_l^\delta})g^{k\bar{q}}g^{p\bar{l}}h_{\alpha\bar{\beta}}f_p^\alpha\overline{f_q^\beta} \\ &= -\lambda\mu + \frac{\mu}{n}\mu^2 \end{aligned}$$

For part *II*, by taking normal coordinate at $p \in M$ and $f(p) \in N$ we may assume $f_i^\alpha = \lambda_i \delta_i^\alpha$ with $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_{r(p)} > \lambda_{r(p)+1} = 0$, where $r(p)$ is rank of $df(p)$. Then $\text{tr}_g(f^* \omega_h) = \sum_{i=1}^n \lambda_i^2$. Hence

$$\begin{aligned}
II &= (\Theta_h)_{\alpha\bar{\beta}\gamma\bar{\delta}}(g^{i\bar{j}} f_i^\alpha \overline{f_j^\beta})(g^{p\bar{q}} f_p^\gamma \overline{f_q^\delta}) \\
&= \sum_{i,k=1}^n (\Theta_h)_{\alpha\bar{\beta}\gamma\bar{\delta}} f_i^\alpha \overline{f_i^\beta} f_k^\gamma \overline{f_k^\delta} \\
&\stackrel{(a)}{=} \sum_{i,j,k,l=1}^n (\Theta_h)_{\alpha\bar{\beta}\gamma\bar{\delta}} f_i^\alpha \overline{f_j^\beta} f_k^\gamma \overline{f_l^\delta} \left(\frac{\delta_{ij}\delta_{kl} + \delta_{il}\delta_{kj}}{2} \right) \\
&\stackrel{(b)}{=} \frac{n(n+1)}{2} (\Theta_h)_{\alpha\bar{\beta}\gamma\bar{\delta}} f_i^\alpha \overline{f_j^\beta} f_k^\gamma \overline{f_l^\delta} \int_{\mathbb{C}\mathbb{P}^{n-1}} \frac{\xi^i \bar{\xi}^j \xi^k \bar{\xi}^l}{|\xi|^4} \omega_{FS}^{n-1} \\
&= \frac{n(n+1)}{2} \int_{\mathbb{C}\mathbb{P}^{n-1}} (\Theta_h)_{\alpha\bar{\beta}\gamma\bar{\delta}} \frac{(f_i^\alpha \xi^i) \overline{(f_j^\beta \xi^j)} (f_k^\gamma \xi^k) \overline{(f_l^\delta \xi^l)}}{|\xi|^4} \omega_{FS}^{n-1} \\
&\stackrel{(c)}{\leq} -\frac{\kappa(f(p))n(n+1)}{2} \int_{\mathbb{C}\mathbb{P}^{n-1}} \frac{(|f_i^\alpha \xi^i|^2)^2}{|\xi|^4} \omega_{FS}^{n-1}
\end{aligned}$$

where

(a) holds from (N, h) is Kähler.

(b) holds from Lemma 16.2.1.

(c) holds from holomorphic sectional curvature of h is bounded from above by $-\kappa$.

Since $f_i^\alpha = \lambda_i \delta_i^\alpha$, one has

$$\begin{aligned}
\int_{\mathbb{C}\mathbb{P}^{n-1}} \frac{(|f_i^\alpha \xi^i|^2)^2}{|\xi|^4} \omega_{FS}^{n-1} &= \frac{1}{n(n+1)} \sum_{i,j,\alpha,\beta} \lambda_i \lambda_j \lambda_k \lambda_l \delta_i^\alpha \delta_j^\alpha \delta_k^\beta \delta_l^\beta (\delta_{ij}\delta_{kl} + \delta_{il}\delta_{kj}) \\
&= \frac{1}{n(n+1)} \left(\left(\sum_{\alpha} \lambda_{\alpha}^2 \right)^2 + \sum_{\alpha} \lambda_{\alpha}^4 \right) \\
&\stackrel{(d)}{\geq} \frac{1}{n(n+1)} \cdot \frac{r+1}{r} \left(\sum_{\alpha} \lambda_{\alpha}^2 \right)^2
\end{aligned}$$

where (d) holds since r is the maximal numbers of non-zero elements of λ_{α} . Hence one has

$$II \leq -\frac{(r+1)f^* \kappa}{2r} \left(\sum_{\alpha} \lambda_{\alpha}^2 \right)^2 = -\frac{(r+1)f^* \kappa}{2r} u^2$$

This completes the proof. \square

Corollary 16.2.1. Let (M, ω) be a complete Kähler manifold with Ricci curvature bounded from below by a positive constant. Then M is compact, and there is no non-trivial holomorphic map from M into a Kähler manifold with non-positive holomorphic sectional curvature.

Corollary 16.2.2. Let (N, h) be a Hermitian manifold with non-positive holomorphic sectional curvature. Then any holomorphic map from $\mathbb{C}\mathbb{P}^n$ to N is constant. In particular, N contains no rational curves.

Corollary 16.2.3. Let (N, h) be a Hermitian manifold with negative holomorphic sectional curvature. Then any holomorphic map from torus \mathbb{T}^2 to N is constant.

Part 5. Topics

17. CALABI-YAU THEOREM

17.1. Introduction. The investigation into the existence of Kähler-Einstein metrics is a compelling and extensive topic that traces back to 1954. In which year Calabi proposed the famous Calabi conjecture, which was finally solved by Yau in 1976. Firstly, Calabi proved the uniqueness of the solution and laid out the program of proving the existence by the method of continuity and also pointed out the openness and the need of a priori estimates in [Cal57]. The very easy a priori zeroth order estimate for the case of negative first Chern class was firstly given by Aubin in [Aub78], but he did not apply his a priori estimate to the continuity method. Instead he used the method of variation which is rather difficult to comprehend.

Yau made the important contribution of using Moser's method of integration by parts and iteration by Sobolev inequality to get a priori zeroth order estimate.

Theorem 17.1 ([MR054],[Yau78b]). Let (X, ω_g) be a compact Kähler manifold. If Ω is a real $(1, 1)$ -form which represents $2\pi c_1(X)$, then there exists a unique metric $\omega \in [\omega_g]$ such that $\text{Ric}(\omega) = \Omega$.

This remarkable result establishes several related results which are of fundamental importance in the study of complex manifolds.

Corollary 17.1.1. Let (X, ω_g) be a compact Kähler manifold with $c_1(X) = 0$. Then there exists a unique Ricci flat metric.

Corollary 17.1.2. Let (X, ω_g) be a compact Kähler manifold with $c_1(X) > 0$. Then M is simply-connected.

Corollary 17.1.3 ([Yau77]). Every complex surface which is homotopic equivalent to $\mathbb{C}\mathbb{P}^2$ is biholomorphic to $\mathbb{C}\mathbb{P}^2$

17.2. The Monge-Ampère equation and priori estimates.

17.2.1. The reformulation of Calabi conjecture in Monge-Ampère equation. The Calabi conjecture can be reduced to a problem of fully non-linear partial differential equations. By Lemma 9.3.1, that is $\partial\bar{\partial}$ -lemma, one has

$$\omega = \omega_g + \sqrt{-1}\partial\bar{\partial}\varphi$$

where φ is the smooth function we desire. Again by $\partial\bar{\partial}$ -lemma one also has

$$\text{Ric}(\omega_g) = \Omega + \sqrt{-1}\partial\bar{\partial}F$$

where F is a smooth function which is unique up to a constant. If we consider the normalization

$$\int_X e^F \omega_g^n = \int_X \omega_g^n$$

then F is unique. Suppose $\text{Ric}(\omega) = \Omega$. Then

$$\text{Ric}(\omega_g) - \text{Ric}(\omega) = \sqrt{-1}\partial\bar{\partial}F$$

that is

$$\sqrt{-1}\partial\bar{\partial}\log\frac{\omega^n}{\omega_g^n} = \sqrt{-1}\partial\bar{\partial}F$$

which is equivalent to the following equation

$$\omega^n = e^{F+C}\omega_g^n$$

and by normalization of F one has $C = 0$. Thus in order to solve Calabi conjecture, it suffices to solve the following complex Monge-Ampère equation:

$$(17.1) \quad \begin{cases} (\omega_g + \sqrt{-1}\partial\bar{\partial}\varphi)^n = e^F\omega_g^n \\ \int_X e^F\omega_g^n = 1 \end{cases}$$

Theorem 17.2.1 ([Cal57]). The solution of (17.1) is unique.

Proof. Suppose φ_1, φ_2 are two solutions. Then

$$\begin{aligned} 0 &= \omega_1^n - \omega_2^n = (\omega_1 - \omega_2)(\omega_1^{n-1} + \cdots + \omega_2^{n-1}) \\ &= \sqrt{-1}\partial\bar{\partial}(\varphi_1 - \varphi_2)(\omega_1^{n-1} + \cdots + \omega_2^{n-1}) \end{aligned}$$

If we define $\psi = \varphi_1 - \varphi_2$, then

$$\begin{aligned} -\int_X \psi\sqrt{-1}\partial\bar{\partial}\psi \wedge (\omega_1^{n-1} + \cdots + \omega_2^{n-1}) &\stackrel{(1)}{=} \int_X \partial\psi \wedge \bar{\partial}\psi \wedge (\omega_1^{n-1} + \cdots + \omega_2^{n-1}) \\ &\stackrel{(2)}{\geq} \frac{1}{n}|\partial\psi|^2\omega_1^n \end{aligned}$$

where

(1) holds from integration by parts and ω_1, ω_2 are Kähler forms.

(2) holds from (2) of Proposition 6.3.3 and positivity of ω_1, ω_2 .

This shows $\partial\psi = 0$, and thus ψ is a constant, which completes the proof. \square

To solve the existence of solution of (17.1), Yau used the continuity method. Consider a sequence of equations as follows

$$(17.2) \quad \det(g_{i\bar{j}} + \varphi_{i\bar{j}}) = \frac{e^{tF}}{\int_X e^{tF}\omega_g^n} \det(g_{i\bar{j}})$$

It's clear (17.2) is solvable at $t = 0$, and thus consider the following set is non-empty

$$I = \{t \in [0, 1] \mid \text{equation (17.2) is solvable}\}$$

The openness of I can be relatively easily demonstrated through the application of the inverse function theorem. However, the true challenge lies in establishing its closedness. To accomplish this, a series of rigorous a priori estimates are essential.

17.2.2. C^0 -estimate.

Theorem 17.2.2 (C^0 -estimate). Suppose φ is a solution of (17.1). Then there exists constant C depending on X, ω_g, F such that

$$\sup \varphi - \inf \varphi \leq C$$

Proof. Without lose of generality we may assume $\sup \varphi = -1$. Then Green formula says

$$\varphi(x) = \frac{1}{V} \int \varphi(y) \omega_g^n - \frac{1}{V} \int G(x, y) \Delta \varphi(y) \omega_g^n$$

and $\omega - \omega_g = \sqrt{-1} \partial \bar{\partial} \varphi$ implies

$$\text{tr}_g \sqrt{-1} \partial \bar{\partial} \varphi = \text{tr}_g \omega - n \geq n$$

Assume $G \geq 0$, if $\varphi(p) = -1$, then

$$\begin{aligned} -1 &= \varphi(p) \\ &= \frac{1}{V} \int \varphi(y) \omega_g^n - \frac{1}{V} \int G(p, y) \Delta \varphi(y) \omega_g^n \\ &\leq \frac{1}{V} \int \varphi(y) \omega_g^n + \frac{1}{V} \int n G(p, y) \omega_g^n \end{aligned}$$

this shows

$$\|\varphi\|_{L^1} = - \int \varphi \omega_g^n \leq C_1$$

For L_2 -estimate, note that

$$\begin{aligned} \int \varphi (e^F - 1) \omega_g^n &= \int \varphi (\omega^n - \omega) \\ &= \int \varphi (\sqrt{-1} \partial \bar{\partial} \varphi) (\omega^{n-1} + \dots + \omega_g^{n-1}) \\ &\leq -\frac{1}{n} \int |\partial \varphi|^2 \omega_g^n \end{aligned}$$

that is

$$\int |\partial \varphi|^2 \omega_g^n \leq n \sup |e^F - 1| \int |\varphi| \omega_g^n \leq C_2$$

that is $\|\nabla \varphi\|_{L^2}^2 \leq C_3$, and by Poincaré inequality one has

$$\|\varphi - \bar{\varphi}\|_{L^2} \leq C_4 \|\nabla \varphi\|_{L^2}$$

that is $\|\varphi\|_{L^2} \leq C_5$.

For $p \geq 2$, set $\psi = -\varphi$, one has

$$\begin{aligned}
\int \psi^{p-1}(e^F - 1)\omega_g^n &= - \int \psi^{p-1}\sqrt{-1}\partial\bar{\partial}\psi \wedge (\omega^{n-1} + \dots + \omega_g^{n-1}) \\
&= \int (p-1)\psi^{p-2}\sqrt{-1}\partial\psi \wedge \bar{\partial}\psi \wedge (\omega^{n-1} + \dots + \omega_g^{n-1}) \\
&\geq \int (p-1)\psi^{p-2}\sqrt{-1}\partial\psi \wedge \bar{\partial}\psi \wedge \omega_g^{n-1} \\
&= \frac{4(p-1)}{p^2} \int \sqrt{-1}\partial\psi^{\frac{p}{2}} \wedge \bar{\partial}\psi^{\frac{p}{2}} \wedge \omega_g^{n-1} \\
&= \frac{4(p-1)}{np^2} \int |\partial\psi^{\frac{p}{2}}|^2 \omega_g^n
\end{aligned}$$

This gives

$$\int |\partial\psi^{\frac{p}{2}}|^2 \omega_g^n \leq C_6 p \int |\varphi|^{p-1} \omega_g^n$$

that is

$$\|\nabla\psi\|_{L^2}^2 \leq C_6 p \|\psi\|_{L^{p-1}}^{p-1}$$

The Sobolev inequality shows for all $f \in W^{1,q}(X, \omega_g)$, one has

$$\|f\|_{L^{\frac{2nq}{2n-q}}} \leq C_7 \|f\|_{W^{1,q}}$$

Set $f = \psi^{\frac{p}{2}}$ and $q = 2$, one has

$$\begin{aligned}
\left(\int \psi^{\frac{np}{n-1}} \omega_g^n\right)^{\frac{n-1}{n}} &= \|\psi\|_{L^{\frac{np}{n-1}}}^p = \|\psi\|_{L^{\frac{2n}{n-1}}}^2 \\
&\leq C_8 \left(\int |\nabla\psi^{\frac{p}{2}}|^2 \omega_g^n + \int \psi \omega_g^n\right) \\
&\leq C_9 (p \|\psi\|_{L^{p-1}}^{p-1} + \|\psi\|_{L^p}^p) \\
&\leq C_{10} p \|\psi\|_{L^p}^p
\end{aligned}$$

This gives

$$\|\psi\|_{L^{\frac{np}{n-1}}} \leq C_{10}^{\frac{1}{p}} p^{\frac{1}{p}} \|\psi\|_{L^p}$$

Set $p_k = (\frac{n}{n-1})^k p$, then

$$\|\psi\|_{L^{p_{k+1}}} \leq (C_{10} p_k)^{\frac{1}{p_k}} \|\psi\|_{L^{p_k}} \leq \prod_{j=0}^k (C_{10} p_j)^{\frac{1}{p_j}} \|\psi\|_{L^p}$$

This gives

$$\|\psi\|_{L^\infty} \leq C \|\psi\|_{L^p}$$

□

17.2.3. C^2 -estimate.

Theorem 17.2.3. Suppose ω_h is a solution of (17.1). Then there exists a constant c depending on X, ω_g, F such that

$$c^{-1}\omega_g \leq \omega_h \leq c\omega_g$$

Proof. It suffices to prove $\text{tr}_g \omega_h \leq c$ since linear algebra yields the following inequality

$$\text{tr}_h \omega_g \leq \frac{1}{n-1} (\text{tr}_g \omega_h)^{n-1} \frac{\omega_g^n}{\omega_h^n}$$

By Corollary 16.1.1, that is Schwarz computation, one has

$$\Delta_h \log \text{tr}_g \omega_h \geq \frac{1}{\text{tr}_g \omega_h} \left(h^{i\bar{j}} (\Theta_g)_{i\bar{j}k\bar{l}} g^{k\bar{q}} g^{p\bar{l}} h_{p\bar{q}} - h^{i\bar{j}} (\Theta_h)_{i\bar{j}p\bar{q}} g^{p\bar{q}} \right)$$

where $\Delta_h = \text{tr}_h \sqrt{-1} \partial \bar{\partial}$. Since (X, g) is a given Kähler manifold, one has its curvature is bounded from below by $-B$ as follows

$$(\Theta_g)_{i\bar{j}k\bar{l}} \geq -B(g_{i\bar{j}}g_{k\bar{l}} + g_{i\bar{l}}g_{k\bar{j}})$$

Then

$$\begin{aligned} h^{i\bar{j}} (\Theta_g)_{i\bar{j}k\bar{l}} g^{k\bar{q}} g^{p\bar{l}} h_{p\bar{q}} &\geq -B(g_{i\bar{j}}g_{k\bar{l}} + g_{i\bar{l}}g_{k\bar{j}}) g^{k\bar{q}} g^{p\bar{l}} h^{i\bar{j}} h_{p\bar{q}} \\ &= -B \left(1 + \frac{1}{n} \right) \delta_i^p \delta_j^q h^{i\bar{j}} h_{p\bar{q}} \\ &= -(n+1)B \end{aligned}$$

On the other hand, one has

$$h^{i\bar{j}} (\Theta_h)_{i\bar{j}p\bar{q}} g^{p\bar{q}} = \left(g^{i\bar{j}} (\Theta_g)_{i\bar{j}p\bar{q}} - F_{p\bar{q}} \right) g^{p\bar{q}} = s - F_{p\bar{q}} g^{p\bar{q}}$$

where $\text{Ric}(\omega_g) = \Omega + \sqrt{-1} \partial \bar{\partial} F$ and s is the scalar curvature of (X, g) . Then by $\text{tr}_h \omega_g \text{tr}_g \omega_h \geq n^2$ one has

$$\begin{aligned} \Delta_h \log \text{tr}_g \omega_h &\geq -\frac{1}{n^2} ((n+1)B + c_0) \text{tr}_h \omega_g \\ &\geq -2B \text{tr}_h \omega_g - \frac{c_0}{n^2} \text{tr}_g \omega_h \end{aligned}$$

where $c_0 = (s - F_{p\bar{q}} g^{p\bar{q}})$. Note that

$$\Delta_h \varphi = h^{j\bar{k}} \varphi_{j\bar{k}} = h^{j\bar{k}} (h_{j\bar{k}} - g_{j\bar{k}}) = n - \text{tr}_h \omega_g$$

Then there exists an appropriate λ such that

$$\Delta_h (\log \text{tr}_g \omega_h + \lambda \varphi) \geq \text{tr}_h \omega_g - C_1$$

If $\log \text{tr}_g \omega_h + \lambda \varphi$ obtain its maximum at $p \in X$, then $\text{tr}_h \omega_g(p) \leq C_1$, and again by trick of linear algebra one has

$$\text{tr}_g \omega_h(p) \leq C_2$$

for some constant C_2 . Since p is the point such that $\log \text{tr}_g \omega_h + \lambda \varphi$ obtains its maximum, then by C^0 -estimate there exists some constant C such that

$$\text{tr}_g \omega_h \leq C$$

This completes the proof. \square

17.2.4. *C³-estimate.* Let $(\Gamma_g)_{ij}^k$ and $(\Gamma_h)_{ij}^k$ denote the Christoffel symbols of g and h respectively, and set

$$S_{ij}^k = (\Gamma_h)_{ij}^k - (\Gamma_g)_{ij}^k$$

Suppose we have the following C^2 -estimate

$$c^{-1}\omega_g \leq \omega_h \leq c\omega_g$$

Lemma 17.2.1. There exists a constant C depending on X, ω_g, F, c such that

$$\Delta_h |S|_h^2 \geq -C|S|_h^2 - C$$

Lemma 17.2.2. There exists a constant C depending on X, ω_g, F, c such that

$$|S| < C$$

17.3. Proof of Calabi-Yau theorem.

Theorem 17.3.1. Let (X, ω_g) be a compact Kähler manifold. Then for any $k \geq 3$ and $F \in C^k(X, \mathbb{R})$, the complex Monge-Ampère equation

$$\begin{cases} \omega_h^n = e^F \omega_g^n \\ \omega_h = \omega_g + \sqrt{-1} \partial \bar{\partial} \varphi \\ \int_X e^F \omega_g^n = \int_X \omega_g^n \end{cases}$$

has a solution $\varphi \in C^{k+1, \alpha}(X)$.

17.4. Aubin-Yau theorem.

17.4.1. *Uniqueness of Kähler-Einstein metric when $c_1(X) < 0$.*

Lemma 17.4.1. Let X be a compact complex manifold. Then there exists at most one Kähler metric ω such that $\text{Ric}(\omega) = -\omega$.

Proof. Suppose that there are two Kähler metrics ω_1 and ω_2 such that

$$\text{Ric}(\omega_1) = -\omega_1$$

$$\text{Ric}(\omega_2) = -\omega_2$$

By $\partial\bar{\partial}$ -lemma, there exists $\varphi \in C^\infty(X, \mathbb{R})$ such that $\omega_1 = \omega_2 + \sqrt{-1} \partial \bar{\partial} \varphi$, so one has

$$\sqrt{-1} \partial \bar{\partial} \varphi = \sqrt{-1} \partial \bar{\partial} \log \frac{\omega_1^n}{\omega_2^n}$$

In other words, there exists a constant c such that

$$\omega_1^n = e^{\varphi+c} \omega_2^n$$

Suppose φ attains its maximum at point p , that is, $\sqrt{-1} \partial \bar{\partial} \varphi(p) \leq 0$. Then

$$\omega_1 = \omega_2 + \sqrt{-1} \partial \bar{\partial} \varphi \leq \omega_2$$

which implies $e^{\varphi(p)+c} \leq 1$. This shows $\varphi + c \leq \varphi(p) + c \leq 0$. Similarly, if φ attains its minimum at q , then $e^{\varphi(q)+c} \geq 1$ and $\varphi + c \geq \varphi(q) + c \geq 0$. Hence $\varphi + c \equiv 0$, and thus $\omega_1 = \omega_2$. \square

17.4.2. *The reformulation of Aubin-Yau theorem in Monge-Ampère equation.*

Theorem 17.4.1 (Aubin-Yau). Let X be a complex manifold with $c_1(X) < 0$. Then there exists a unique Kähler metric $\omega \in 2\pi c_1(X)$ such that $\text{Ric}(\omega) = -\omega$.

Proof. Let ω_g be any Kähler metric in $[-2\pi c_1(X)]$. Then by $\partial\bar{\partial}$ -lemma there exists a smooth function F such that

$$\text{Ric}(\omega_g) = -\omega_g + \sqrt{-1}\partial\bar{\partial}F$$

since $[\text{Ric}(\omega_g)] = 2\pi c_1(X)$. Let ω be another Kähler metric such that $[\omega] = [\omega_g]$. Again by $\partial\bar{\partial}$ -lemma there exists a smooth function φ such that

$$\omega = \omega_g + \sqrt{-1}\partial\bar{\partial}\varphi$$

Then $\text{Ric}(\omega) = -\omega$ is equivalent to

$$\sqrt{-1}\partial\bar{\partial} \log \frac{\omega^n}{\omega_g^n} = \sqrt{-1}\partial\bar{\partial}(F + \varphi)$$

or in other words, there exists a constant C such that

$$\omega^n = e^{F+\varphi+C}\omega_g^n$$

By rescaling F we may assume $C = 0$, so Aubin-Yau is equivalent to solve the following complex Monge-Ampère equation:

$$(17.3) \quad \begin{cases} (\omega_g + \sqrt{-1}\partial\bar{\partial}\varphi)^n = e^{F+\varphi}\omega_g^n \\ \int_X e^{F+\varphi}\omega_g^n = 1 \end{cases}$$

\square

17.4.3. *C^0 -estimate.*

Theorem 17.4.2 (C^0 -estimate). Suppose φ is a solution of (17.3). Then

$$\sup_X |\varphi| \leq \sup_X |F|$$

Proof. As argument in Lemma 17.4.1, at maximum point p of φ , one has $\varphi + F(p) \leq \varphi(p) + F(p) \leq 0$, so

$$\sup_X \varphi \leq -F(p) \leq \sup_X |F|$$

Similarly, at minimum point q of φ , $\varphi + F(q) \geq \varphi(q) + F(q) \geq 0$ and

$$\inf_X \varphi \geq -F(q) \geq -\sup_X |F|$$

Hence $\sup_X |\varphi| \leq \sup_X |F|$. \square

17.4.4. C^2 -estimate.

Theorem 17.4.3 (C^2 -estimate). There exists a uniform constant C depending on X, ω_g, F such that

$$C^{-1}\omega_g \leq \omega \leq C\omega_g$$

Proof. The same as Theorem 17.2.3. □

17.4.5. C^3 -estimate. Given the C^0 -estimate and C^2 -estimate, the C^3 -estimate is very similar to that in the proof of Calabi-Yau theorem. Here we formulate a general setup. Let $(\Gamma_g)_{ij}^k$ and $(\Gamma_h)_{ij}^k$ denote the Christoffel symbols of g and h respectively, and set

$$S_{ij}^k = (\Gamma_h)_{ij}^k - (\Gamma_g)_{ij}^k$$

Suppose we have the following C^2 -estimate

$$c^{-1}\omega_g \leq \omega_h \leq c\omega_g$$

Theorem 17.4.4 (C^3 -estimate).

17.4.6. *Proof of Aubin-Yau theorem.*

18. DEFORMATIONS OF COMPLEX STRUCTURE

18.1. The Maurer-Cartan equation. Recall that a complex structure on a smooth manifold M is encoded by an integrable almost complex structure J , and two complex manifolds (M, J) and (M', J') are isomorphic if there exists a diffeomorphism $F: M \rightarrow M'$ such that $dF \circ J = J' \circ dF$. Thus the set of all complex structures on a fixed smooth manifold M is the quotient of the set

$$\mathcal{A}_c(M) := \{J \mid J \text{ is an integrable almost complex structure on } M\}$$

of all complex structures by the action of the diffeomorphism group. Firstly let's consider the set

$$\mathcal{A}_{ac}(M) := \{J \mid J \text{ is an almost complex structure on } M\}$$

For arbitrary almost complex structure $J \in \mathcal{A}_{ac}(M)$, it's uniquely determined by a decomposition of the complexified tangent bundle $T_{\mathbb{C}}M = T^{1,0}M \oplus T^{0,1}M$ with J is $\sqrt{-1}$ id on $T^{1,0}M$ and $-\sqrt{-1}$ id on $T^{0,1}M$. In fact, giving $T^{0,1}M \subseteq T_{\mathbb{C}}M$ is enough, since we can set $T^{1,0}M = \overline{T^{0,1}M}$. If $J(t)$ is a continuous family of almost complex structure with $J(0) = J$, there is a continuous family of such decompositions $T_{\mathbb{C}}M = T_t^{1,0}M \oplus T_t^{0,1}M$, or equivalently, of subspaces $T_t^{0,1}M \subseteq T_{\mathbb{C}}M$.

Thus for small t the deformations $J(t)$ of J gives a map

$$\phi(t): T^{0,1}M \rightarrow T^{1,0}M$$

with $v + \phi(t)v \in T_t^{1,0}M$. Conversely, if $\phi(t): T^{0,1}M \rightarrow T^{1,0}M$ is given, then one defines for small t

$$T_t^{1,0}M := (\text{id} + \phi(t))(T^{0,1}M)$$

Here the condition t to be "small" has to be imposed in order to ensure that with this definition $T_t^{1,0}M \subseteq T_{\mathbb{C}}M \rightarrow T^{1,0}M$ is an isomorphism. Thus deformations of almost complex structure is encoded by such a map $\phi(t)$. Now for convenience we assume J is an integrable almost complex, and thus we denote (M, J) by X . In particular, there is $\bar{\partial}$ operator on the holomorphic tangent bundle TX , which can be applied to $\phi(t) \in C^\infty(X, \Omega_X^{0,1} \otimes T^{1,0}X)$.

The following proposition shows that the integrability condition for deformation of the almost complex structure can be rephrased in terms of $\phi(t)$.

Proposition 18.1.1. The integrability condition is equivalent to the Maurer-Cartan equation

$$\bar{\partial}\phi(t) + [\phi(t), \phi(t)] = 0$$

Proof. It suffices to compute locally so we write $\phi(t)$ with respect to local coordinate $\{z^1, \dots, z^n\}$ as follows

$$\phi(t) = \phi_i^j(t) d\bar{z}^i \otimes \frac{\partial}{\partial z_j}$$

The integrability condition implies $[T_t^{0,1}X, T_t^{0,1}X] \subseteq T_t^{0,1}X$, so in particular one has

$$\left[\frac{\partial}{\partial \bar{z}_i} + \phi(t) \left(\frac{\partial}{\partial \bar{z}_i} \right), \frac{\partial}{\partial \bar{z}_j} + \phi(t) \left(\frac{\partial}{\partial \bar{z}_j} \right) \right] \in T_t^{0,1}X$$

A direct computation yields

$$\left(\left[\frac{\partial}{\partial \bar{z}^i}, \phi_k^l(t) \frac{\partial}{\partial z^l} \right] + \left[\phi_i^j(t) \frac{\partial}{\partial z^j}, \frac{\partial}{\partial \bar{z}^k} \right] + \left[\phi_i^j(t) \frac{\partial}{\partial z^j}, \phi_k^l(t) \frac{\partial}{\partial z^l} \right] \right) \in T_t^{0,1}X$$

Note that

$$\begin{aligned} \left[\frac{\partial}{\partial \bar{z}^i}, \phi_k^l(t) \frac{\partial}{\partial z^l} \right] &= \frac{\partial \phi_k^l(t)}{\partial \bar{z}^i} \frac{\partial}{\partial z^l} \\ \left[\phi_i^j(t) \frac{\partial}{\partial z^j}, \frac{\partial}{\partial \bar{z}^k} \right] &= -\frac{\partial \phi_i^j(t)}{\partial \bar{z}^k} \frac{\partial}{\partial z^j} \end{aligned}$$

This shows

$$\left[\frac{\partial}{\partial \bar{z}^i}, \phi_k^l(t) \frac{\partial}{\partial z^l} \right] + \left[\phi_i^j(t) \frac{\partial}{\partial z^j}, \frac{\partial}{\partial \bar{z}^k} \right] = \left(\frac{\partial \phi_k^j(t)}{\partial \bar{z}^i} - \frac{\partial \phi_i^j(t)}{\partial \bar{z}^k} \right) \frac{\partial}{\partial z^j} = \bar{\partial} \phi(t) \left(\frac{\partial}{\partial \bar{z}^i}, \frac{\partial}{\partial \bar{z}^k} \right)$$

On the other hand, note that

$$[\phi(t), \phi(t)] = d\bar{z}^i \wedge d\bar{z}^k \left[\phi_i^j(t) \frac{\partial}{\partial z^j}, \phi_k^l(t) \frac{\partial}{\partial z^l} \right]$$

Thus

$$\left[\phi_i^j(t) \frac{\partial}{\partial z^j}, \phi_k^l(t) \frac{\partial}{\partial z^l} \right] = [\phi(t), \phi(t)] \left(\frac{\partial}{\partial \bar{z}^i}, \frac{\partial}{\partial \bar{z}^k} \right)$$

So integrability condition implies

$$\bar{\partial}(t) + [\phi(t), \phi(t)] \left(\frac{\partial}{\partial \bar{z}^i}, \frac{\partial}{\partial \bar{z}^k} \right) \in T_t^{0,1}X$$

This shows $\bar{\partial}(t) + [\phi(t), \phi(t)]$ is a section of $\Omega_X^{0,2} \otimes (T^{1,0}X \cap T_t^{0,1}X)$, but for sufficiently small t one has $T^{1,0}X \cap T_t^{0,1}X = 0$. This shows

$$\bar{\partial}(t) + [\phi(t), \phi(t)] = 0$$

Conversely, if the Maurer-Cartan equation holds, then the integrability condition holds for a local frame of $T_t^{0,1}X$, and thus for all sections of $T_t^{0,1}X$. \square

Now let's consider the power series expansion of a given deformation $\phi(t)$ as follows

$$\phi(t) = \phi_0 + \phi_1 t + \phi_2 t^2 + \dots$$

The Maurer-Cartan equation gives

$$\bar{\partial} \left(\sum_{i=1}^{\infty} \phi_i t^i \right) + \sum_{i,j=1}^{\infty} [\phi_i, \phi_j] t^{i+j} = 0$$

This yields a recursive system of equations

$$\begin{aligned} 0 &= \bar{\partial}\phi_1 \\ 0 &= \bar{\partial}\phi_2 + [\phi_1, \phi_1] \\ &\vdots \\ 0 &= \bar{\partial}\phi_k + \sum_{0 < i < k} [\phi_i, \phi_{k-i}] \end{aligned}$$

In particular, the first-order deformation of the complex structure is determined by a $\bar{\partial}$ -closed $(0, 1)$ -form ϕ_1 with valued in TX . Thus it determines an element $[\phi_1] \in H^1(X, TX)$.

Definition 18.1.1 (Kodaira-Spencer class). The Kodaira-Spencer class of a one-parameter deformation $J(t)$ of a complex structure J is the induced cohomology class $[\phi_1] \in H^1(X, TX)$.

Proposition 18.1.2. Let X be a complex manifold. There is a natural bijection between all first-order deformations of X and elements of $H^1(X, TX)$.

Corollary 18.1.1. A first-order deformations $v \in H^1(X, TX)$ cannot be integrated if $[v, v] \in H^2(X, TX)$ does not vanish.

19. HERMITIAN-YANG-MILLS METRIC

Definition 19.1 (Hermitian-Yang-Mills metric). Let (X, ω) be a compact Kähler manifold and (E, h) be a holomorphic vector bundle over X with Chern curvature Θ_h . The metric h is called an Hermitian-Yang-Mills metric if

$$\mathrm{tr}_\omega \Theta_h = \lambda I$$

for some constant $\lambda \in \mathbb{R}$, where I is the identity operator in $\mathrm{End} E$.

Remark 19.1 (local form). Suppose the Chern curvature Θ_h is locally given by $(\Theta_h)_{\beta i \bar{j}}^\alpha$, where

$$(\Theta_h)_{\beta}^\alpha = \bar{\partial}(h^{\alpha \bar{\gamma}} \partial h_{\beta \bar{\gamma}}).$$

Then the Hermitian-Yang-Mills equation is

$$g^{i \bar{j}} (\Theta_h)_{\beta i \bar{j}}^\alpha = \lambda \delta_\beta^\alpha,$$

where $\omega = \sqrt{-1} g_{i \bar{j}} dz^i \wedge d\bar{z}^j$.

The final goal of this section is to introduce the following two celebrated theorems of existence of Hermitian-Yang-Mills metrics.

Theorem 19.1 ([UY86]). Let (X, ω) be a compact Kähler manifold and E be an ω -stable holomorphic vector bundle over X . Then E admits a unique Hermitian-Yang-Mills metric.

Theorem 19.2 ([Sim88]). Let (X, ω) be a compact Kähler manifold and (E, θ) be an ω -stable Higgs bundle over X . Then (E, θ) admits an Hermitian-Yang-Mills metric.

Along the way, we're going to talk the following topics

- (1) ω -stabilities of holomorphic vector bundle.
- (2)
- (3) Higgs bundles.
- (4)

19.1. Stable bundle. In this section, X always denotes a complex manifold.

Definition 19.1.1 (degree). Let $\pi: E \rightarrow X$ be a holomorphic vector bundle. The degree of E is defined as

$$\mathrm{deg}(E) := \int_X c_1(E).$$

Definition 19.1.2 (slope). Let $\pi: E \rightarrow X$ be a holomorphic vector bundle. The slope of E is defined as

$$\mu(E) := \frac{\mathrm{deg}(E)}{\mathrm{rk}(E)}.$$

Remark 19.1.1. Note that the slope of a holomorphic vector bundle is independent of the holomorphic structure, since both the degree and rank are topological invariants.

Definition 19.1.3 (slope stability). Let $\pi: E \rightarrow X$ be a holomorphic vector bundle.

- (1) E is said to be stable if for every non-trivial holomorphic subbundle F , $\mu(F) < \mu(E)$;
- (2) E is said to be semi-stable if for every non-trivial holomorphic subbundle F , $\mu(F) \leq \mu(E)$;
- (3) E is said to be unstable if it's not semi-stable.

Remark 19.1.2.

- (a) It's clear that all holomorphic line bundles are stable, since they don't have non-trivial subbundles;
- (b) A semi-stable vector bundle with coprime rank and degree is actually stable, since
- (c) The slope stability is not topological invariant, since here we only consider holomorphic subbundles, which depends on the holomorphic structure.

Proposition 19.1.1. Let $\pi: E \rightarrow X$ be a holomorphic vector bundle.

- (1) It's stable if and only if for every non-trivial holomorphic subbundle F , $\mu(E/F) > \mu(E)$;
- (2) It's semi-stable if and only if for every non-trivial holomorphic subbundle F , $\mu(E/F) \geq \mu(E)$.

Proof. Denote r, r', r'' the ranks of $E, F, E/F$ respectively, and d, d', d'' their degrees respectively. From exact sequence

$$0 \rightarrow F \rightarrow E \rightarrow E/F \rightarrow 0$$

one has $r = r' + r''$ and $d = d' + d''$, thus

$$\frac{d'}{r'} < \frac{d' + d''}{r' + r''} \iff \frac{d'}{r'} < \frac{d''}{r''} \iff \frac{d' + d''}{r' + r''} < \frac{d''}{r''}$$

and likewise with the case semi-stable. \square

A philosophy is that semi-stable bundles don't admit too many subbundles, since any subbundle they may have is of slope no greater than their own. This turns out to have many interesting consequences we're going to show, for example, the category of semi-stable bundles is abelian.

Lemma 19.1.1. If $\varphi: E \rightarrow E'$ is a non-zero homomorphism of holomorphic vector bundles over X , then

$$\mu(E/\ker \varphi) \leq \mu(\operatorname{im} \varphi)$$

Proposition 19.1.2. Let E, E' be two semi-stable bundles such that $\mu(E) > \mu(E')$. Then any homomorphism $\varphi: E \rightarrow E'$ is zero.

Proof. If φ is non-zero, since E is semi-stable, then

$$\mu(\operatorname{im} \varphi) \stackrel{(1)}{\geq} \mu(E/\ker \varphi) \stackrel{(2)}{\geq} \mu(E) > \mu(E')$$

where

(1) holds from Lemma 19.1.1;

(2) holds from Proposition 19.1.1.

which contradicts to the semi-stability of E' . \square

Proposition 19.1.3. Let $\varphi: E \rightarrow E'$ be a non-zero homomorphism of semi-stable holomorphic of slope μ . Then $\ker \varphi$ and $\operatorname{im} \varphi$ are semi-stable bundles of slope μ , and the natural map $E/\ker \varphi \rightarrow \operatorname{im} \varphi$ is an isomorphism.

Corollary 19.1.1. The category of semi-stable bundles of slope μ is abelian, and the simple object¹⁸ in this category is the stable bundles of slope μ .

Proof. By Proposition 19.1.3 one has the category of semi-stable bundles of slope μ is abelian. A stable bundle E is simple in this category, since it admits no non-trivial subbundles with slope μ ; Conversely, if a semi-stable bundle E is simple, then any non-trivial subbundle F satisfies $\mu(F) \leq \mu(E)$ since E is semi-stable and $\mu(F) \neq \mu(E)$ since E is simple, this shows E is stable. \square

Proposition 19.1.4. Let E, E' be two stable holomorphic vector bundles over X with same slopes and $\varphi: E \rightarrow E'$ be a non-zero homomorphism. Then φ is an isomorphism.

Proof. Since $\varphi: E \rightarrow E'$ is a non-zero homomorphism between stable bundles with same slopes, then by Proposition 19.1.3 one has $\ker \varphi$ is either 0 or has slope $\mu(E)$, but E is actually stable, then $\ker \varphi$ must be 0, and since φ is strict, this shows φ is injective. Likewise, $\operatorname{im} \varphi \neq 0$ and has slope $\mu(E')$, then it must be E' since E' is stable. Then again by φ is strict, $\operatorname{im} \varphi = E'$ implies φ is surjective. Therefore φ is an isomorphism. \square

Proposition 19.1.5. If E is a stable bundle over X , then $\operatorname{End} E = \mathbb{C}$. In particular, $\operatorname{Aut} E = \mathbb{C}^*$.

Proof. Let φ be a non-zero endomorphism of E , by Proposition 19.1.4 one has φ is an automorphism, so $\operatorname{End} E$ is a field, which contains \mathbb{C} as its subfield of scalar endomorphisms. For any $\varphi \in \operatorname{End} E$, by Cayley-Hamilton theorem one has φ is algebraic over \mathbb{C} , and since \mathbb{C} is algebraically closed, this shows $\operatorname{End} E \cong \mathbb{C}$. \square

Corollary 19.1.2. A stable bundle is indecomposable, that is it's not isomorphic to a direct sum of non-trivial subbundles.

Proof. The automorphism group of $E = E_1 \oplus E_2$ contains $\mathbb{C}^* \times \mathbb{C}^*$, so by Proposition 19.1.5 it can't be stable. \square

¹⁸Recall a simple object in an abelian category is an object with no non-trivial sub-object.

Theorem 19.1.1 (Jordan-Hölder filtration). Any semi-stable bundle of slope μ over X admits a filtration

$$0 = E_0 \subset E_1 \subset \cdots \subset E_k = E$$

by holomorphic subbundles such that for each $1 \leq i \leq k$, one has

- (1) E_i/E_{i-1} is stable;
- (2) $\mu(E_i/E_{i-1}) = \mu(E)$.

Proposition 19.1.6 (Seshadri). Any two Jordan-Hölder filtrations

$$S : 0 = E_0 \subset E_1 \subset \cdots \subset E_k = E$$

and

$$S' : 0 = E'_0 \subset E'_1 \subset \cdots \subset E'_k = E$$

of a semi-stable bundle E have same length, and the associated graded objects

$$\text{gr}(S) : 0 = E_1/E_0 \oplus \cdots \oplus E_k/E_{k-1}$$

and

$$\text{gr}(S') : 0 = E'_1/E'_0 \oplus \cdots \oplus E'_k/E'_{k-1}$$

satisfy $E_i/E_{i-1} \cong E'_i/E'_{i-1}$ for all $1 \leq i \leq k$.

Definition 19.1.4 (poly-stable bundle). A holomorphic vector bundle E over X is called poly-stable if it is isomorphic to a direct sum $E_1 \oplus \cdots \oplus E_k$ of stable bundles of the same slope.

Example 19.1.1. A stable bundle is poly-stable.

Example 19.1.2. The graded object associated to any Jordan-Hölder filtration of a semi-stable bundle E is a poly-stable, and by Proposition 19.1.6, it's unique up to isomorphism, this isomorphic class is denoted by $\text{gr}(E)$.

Part 6. Appendix

20. SHEAF AND COHOMOLOGY

20.1. **Sheaves.** Along this section, X denotes a topological space.

20.1.1. *Definitions and Examples.*

Definition 20.1.1 (sheaf). A presheaf of abelian group \mathcal{F} on X consisting of the following data:

- (1) For any open subset U of X , $\mathcal{F}(U)$ is an abelian group.
- (2) If $U \subseteq V$ are two open subsets of X , then there is a group homomorphism $r_{UV}: \mathcal{F}(U) \rightarrow \mathcal{F}(V)$. Moreover, above data satisfy
 - I $\mathcal{F}(\emptyset) = 0$.
 - II $r_{UU} = \text{id}$.
 - III If $W \subseteq U \subseteq V$ are open subsets of X , then $r_{UW} = r_{VW} \circ r_{UV}$.

Moreover, \mathcal{F} is called a sheaf if it satisfies the following extra conditions

- IV Let $\{V_i\}_{i \in I}$ be an open covering of open subset $U \subseteq X$ and $s \in \mathcal{F}(U)$. If $s|_{V_i} := r_{UV_i}(s) = 0$ for all $i \in I$, then $s = 0$.
- V Let $\{V_i\}_{i \in I}$ be an open covering of open subset $U \subseteq X$ and $s_i \in \mathcal{F}(V_i)$. If $s_i|_{V_i \cap V_j} = s_j|_{V_i \cap V_j}$ for all $i, j \in I$, then there exists $s \in \mathcal{F}(U)$ such that $s|_{V_i} = s_i$ for all $i \in I$.

Example 20.1.1 (constant presheaf). For an abelian group G , the constant presheaf assign each open subset U the group G itself, but in general it's not a sheaf.

Definition 20.1.2 (morphism of presheaves). A morphism $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ between presheaves consisting of the following data:

- (1) For any open subset U of X , there is a group homomorphism $\varphi(U): \mathcal{F}(U) \rightarrow \mathcal{G}(U)$.
- (2) If $U \subseteq V$ are two open subsets of X , then the following diagram commutes

$$\begin{array}{ccc} \mathcal{F}(U) & \xrightarrow{\varphi(U)} & \mathcal{G}(U) \\ \downarrow r_{UV} & & \downarrow r_{UV} \\ \mathcal{F}(V) & \xrightarrow{\varphi(V)} & \mathcal{G}(V) \end{array}$$

Notation 20.1.1. For convenience, for $s \in \mathcal{F}(U)$, we often write $\varphi(s)$ instead of $\varphi(U)(s)$.

Remark 20.1.1. The morphisms between sheaves are defined as morphisms of presheaves.

Definition 20.1.3 (isomorphism). A morphism of presheaves $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ is called an isomorphism if it has two-sided inverse, that is, there exists a morphism of presheaves $\psi: \mathcal{G} \rightarrow \mathcal{F}$ such that $\psi\varphi = \text{id}_{\mathcal{F}}$ and $\varphi\psi = \text{id}_{\mathcal{G}}$.

Remark 20.1.2. A morphism of presheaves $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ is an isomorphism if and only if for every open subset $U \subseteq X$, $\varphi(U): \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ is an isomorphism of abelian groups.

20.1.2. *Stalks.*

Definition 20.1.4 (stalks). For a presheaf \mathcal{F} and $p \in X$, the stalk at p is defined as

$$\mathcal{F}_p = \varinjlim_{p \in U} \mathcal{F}(U)$$

Remark 20.1.3 (alternative definition). In order to avoid language of direct limit, we give a more useful but equivalent description of stalk: For $p \in U \cap V$, $s_U \in \mathcal{F}(U)$ and $s_V \in \mathcal{F}(V)$ are equivalent if there exists $p \in W \subseteq U \cap V$ such that $s_U|_W = s_V|_W$. An element $s_p \in \mathcal{F}_p$, which is called a germ, is an equivalence class $[s_U]$.

Notation 20.1.2.

- (1) For $s \in \mathcal{F}(U)$ and $p \in U$, $s|_p$ denotes the equivalent class it gives.
- (2) For $s_p \in \mathcal{F}_p$, $s \in \mathcal{F}(U)$ denotes the section such that $s|_p = s_p$.

Definition 20.1.5 (morphisms on stalks). Given a morphism of sheaves $\varphi: \mathcal{F} \rightarrow \mathcal{G}$, it induces a morphism of abelian groups $\varphi_p: \mathcal{F}_p \rightarrow \mathcal{G}_p$ as follows:

$$\begin{aligned} \varphi_p: \mathcal{F}_p &\rightarrow \mathcal{G}_p \\ s_p &\mapsto \varphi(s)|_p. \end{aligned}$$

Remark 20.1.4. It's necessary to check the φ_p is well-defined since there are different choices s such that $s|_p = s_p$.

Proposition 20.1.1. Let $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ be a morphism between sheaves. Then φ is an isomorphism if and only if the induced map $\varphi_p: \mathcal{F}_p \rightarrow \mathcal{G}_p$ is an isomorphism for every $p \in X$.

Proof. It's clear if φ is an isomorphism between sheaves, then it induces an isomorphism between stalks. Conversely, it suffices to show $\varphi(U): \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ is an isomorphism for every open subset $U \subseteq X$.

- (1) *Injectivity:* For $s, s' \in \mathcal{F}(U)$ such that $\varphi(s) = \varphi(s')$, by passing to stalks one has $\varphi_p(s|_p) = \varphi_p(s'|_p)$ for every $p \in U$, and thus $s|_p = s'|_p$ since φ_p is an isomorphism. By definition of stalks there exists an open subset $V_p \subseteq U$ containing p such that s agrees with s' on V_p . Then it gives an open covering $\{V_p\}$ of U , and by axiom (IV) one has $s = s'$ on U .
- (2) *Surjectivity:* For $t \in \mathcal{G}(U)$, by passing to stalks there exists $s_p \in \mathcal{F}_p$ such that $\varphi_p(s_p) = t|_p$ for every $p \in U$ since φ_p is surjective. By definition of stalks there exists an open subset $V_p \subseteq U$ containing p and $s \in \mathcal{F}(V_p)$ such that $\varphi(s) = t$ on V_p . This gives a collection of sections defined on an open covering $\{V_p\}$ of U , and by injectivity we proved above one has these sections agree with each other on the intersections. Then by axiom (V) there exists a section $s \in \mathcal{F}(U)$ such that $\varphi(s) = t$.

□

20.1.3. *Sheafification.* In Example 20.1.1, we come across a presheaf that is not a sheaf. To obtain a sheaf from a presheaf, we require a process known as sheafification. One approach to defining sheafification is through its universal property.

Definition 20.1.6 (sheafification). Given a presheaf \mathcal{F} there is a sheaf \mathcal{F}^+ and a morphism $\theta: \mathcal{F} \rightarrow \mathcal{F}^+$ with the property that for any sheaf \mathcal{G} and any morphism $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ there is a unique morphism $\bar{\varphi}: \mathcal{F}^+ \rightarrow \mathcal{G}$ such that the following diagram commutes:

$$\begin{array}{ccc} \mathcal{F} & \xrightarrow{\varphi} & \mathcal{G} \\ \downarrow \theta & \nearrow \bar{\varphi} & \\ \mathcal{F}^+ & & \end{array}$$

The universal property shows that if the sheafification exists, then it's unique up to a unique isomorphism. One way to give an explicit construction of sheafification is to glue stalks together in a suitable way. Let $\mathcal{F}^+(U)$ be a set of functions

$$f: U \rightarrow \coprod_{p \in U} \mathcal{F}_p$$

such that $f(p) \in \mathcal{F}_p$ and for every $p \in U$ there is an open subset $V_p \subseteq U$ containing p and $t \in \mathcal{F}(V_p)$ such that $t|_q = f(q)$ for all $q \in V_p$.

Proposition 20.1.2. \mathcal{F}^+ is the sheafification of \mathcal{F} .

Proof. Firstly let's show \mathcal{F}^+ is a sheaf: It's clear \mathcal{F}^+ is a presheaf, so it suffices to check conditions (IV) and (V) in the definition. Let $U \subseteq X$ be an open subset and $\{V_i\}$ be an open covering of U .

- (1) If $s \in \mathcal{F}^+(U)$ such that $s|_{V_i} = 0$ for all i , then s must be zero: It suffices to show $s(p) = 0$ for all $p \in U$. For any $p \in U$, then there exists an open subset V_i contains p , hence $s(p) = s|_{V_i}(p) = 0$.
- (2) Suppose there exists a collection of sections $\{s_i \in \mathcal{F}^+(V_i)\}_{i \in I}$ such that

$$s_i|_{V_i \cap V_j} = s_j|_{V_i \cap V_j}$$

holds for all $i, j \in I$. Now we construct $s \in \mathcal{F}^+(U)$ as follows: For $p \in U$ and V_i containing p , we define $s(p) = s_i(p)$. This is well-defined since s_i agree on the intersections, so it remains to show $s \in \mathcal{F}^+(U)$. It's clear $s(p) \in \mathcal{F}_p$. For $p \in U$, there exists V_i containing p , and thus there exists $W_i \subseteq V_i$ containing p and $t \in \mathcal{F}(W_i)$ such that $t|_q = s_i(q) = s(q)$ for all $q \in V_p$.

There is a canonical morphism $\theta: \mathcal{F} \rightarrow \mathcal{F}^+$ as follows: For open subset $U \subseteq X$, and $s \in \mathcal{F}(U)$, $\theta(s)$ is defined by

$$\begin{aligned} \theta(s): U &\rightarrow \coprod_{p \in U} \mathcal{F}_p \\ p &\mapsto s|_p. \end{aligned}$$

Note that if \mathcal{F} is a sheaf, the canonical morphism $\theta: \mathcal{F} \rightarrow \mathcal{F}^+$ is an isomorphism.

- (1) Injectivity: If $s \in \mathcal{F}(U)$ such that $s|_p = 0$ for all $p \in U$, then there exists an open covering $\{V_i\}_{i \in I}$ of U such that $s|_{V_i} = 0$, by axiom (IV) of sheaf one has $s = 0$.
- (2) Surjectivity: For $f \in \mathcal{F}^+(U)$ and $p \in U$, there exists $p \in V_p \subseteq U$ and $t \in \mathcal{F}(V_p)$ such that $f|_p = t|_p$ by construction of \mathcal{F}^+ . Then glue these sections together to get our desired s such that $\theta(s) = f$.

Finally let's show \mathcal{F}^+ satisfies the universal property of sheafification. A morphism of presheaves $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ induces a map on stalks

$$\varphi_p: \mathcal{F}_p \rightarrow \mathcal{G}_p.$$

For $f \in \mathcal{F}^+(U)$, the composite of f with the map

$$\prod_{p \in U} \varphi_p: \prod_{p \in U} \mathcal{F}_p \rightarrow \prod_{p \in U} \mathcal{G}_p$$

gives a map $\tilde{\varphi}(f): U \rightarrow \prod_{p \in U} \mathcal{G}_p$, and in fact $\tilde{\varphi}(f) \in \mathcal{G}^+(U)$: For $p \in U$, $\tilde{\varphi}(f)(p) \in \mathcal{G}_p$ since $f|_p \in \mathcal{F}_p$ and $\varphi_p: \mathcal{F}_p \rightarrow \mathcal{G}_p$. If for all $q \in V_p$ we have $t|_q = f|_q$, then

$$\tilde{\varphi}(f)(q) = \varphi_q(f|_q) = \varphi_q(t|_q) = \varphi(t)|_q.$$

Since \mathcal{G} is a sheaf, the canonical morphism $\theta': \mathcal{G} \rightarrow \mathcal{G}^+$ is an isomorphism, so we can define $\bar{\varphi}: = \theta'^{-1} \circ \tilde{\varphi}$. Now let's show $\varphi = \bar{\varphi} \circ \theta = \theta'^{-1} \circ \tilde{\varphi} \circ \theta$. It's easy to show they coincide on each stalk since $\varphi_p = \theta'_p{}^{-1} \circ \tilde{\varphi}_p \circ \theta_p$, and thus $\varphi = \bar{\varphi} \circ \theta$ by Proposition 20.1.1. Furthermore, uniqueness follows from the fact that $\bar{\varphi}_p$ is uniquely determined by φ_p . \square

Remark 20.1.5. From the construction, one can see the stalk of \mathcal{F}^+ at p is exactly \mathcal{F}_p .

Remark 20.1.6. The sheafification can be described in a more fancy language: Since we have sheaf of abelian groups on X as a category, denote it by \underline{Ab}_X , and presheaf is a full subcategory of \underline{Ab}_X , there is a natural inclusion functor ι from category of sheaf to category of presheaf. The sheafification is the adjoint functor of ι .

Example 20.1.2 (constant sheaf). For an abelian group G , the associated constant sheaf \underline{G} is the sheafification of the constant presheaf. By the construction of sheafification, \underline{G} can be explicitly expressed as

$$\underline{G}(U) = \{\text{locally constant function } f: U \rightarrow G\}$$

20.1.4. *Exact sequence of sheaf.* Given a morphism $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ between sheaves of abelian groups, there are the following presheaves

$$\begin{aligned} U &\mapsto \ker \varphi(U) \\ U &\mapsto \operatorname{im} \varphi(U) \\ U &\mapsto \operatorname{coker} \varphi(U), \end{aligned}$$

since $\varphi(U): \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ is a group homomorphism.

Proposition 20.1.3. Kernel of a morphism between sheaves is a sheaf.

Proof. Let $\{V_i\}_{i \in I}$ be an open covering of U .

- (1) For $s \in \ker \varphi(U)$, if $s|_{V_i} = 0$, then $s = 0$ since s is also in $\mathcal{F}(U)$.
- (2) If there exists $s_i \in \ker \varphi(V_i)$ such that $s_i|_{V_i \cap V_j} = s_j|_{V_i \cap V_j}$, then they glue together to get $s \in \mathcal{F}(U)$. Note that

$$\varphi(U)(s)|_{V_i} = \varphi(V_i)(s|_{V_i}) = \varphi(V_i)(s_i) = 0$$

Then $s \in \ker \varphi(U)$. □

But the image of morphism may not be a sheaf. Although we can prove the first requirement in the same way, the proof for the second requirement fails: If there exists $s_i \in \text{im } \varphi(V_i)$, and we can glue them together to get a $s \in \mathcal{G}(U)$, but s may not be the image of some $t \in \mathcal{F}(U)$. The cokernel fails to be a sheaf for the same reason.

Definition 20.1.7 (image and cokernel). Let $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ be a morphism between sheaves of abelian groups. Then the image and cokernel of φ is defined to be the sheafification of the following presheaves

$$\begin{aligned} U &\mapsto \text{im } \varphi(U) \\ U &\mapsto \text{coker } \varphi(U) \end{aligned}$$

respectively.

Definition 20.1.8 (exact). For a sequence of sheaves:

$$\dots \rightarrow \mathcal{F}^{i-1} \xrightarrow{\varphi^{i-1}} \mathcal{F}^i \xrightarrow{\varphi^i} \mathcal{F}^{i+1} \rightarrow \dots$$

It's called exact at \mathcal{F}^i , if $\ker \varphi^i = \text{im } \varphi^{i-1}$. If a sequence is exact at everywhere, then it's an exact sequence of sheaves.

Definition 20.1.9 (short exact sequence). An exact sequence of sheaves is called a short exact sequence if it looks like

$$0 \rightarrow \mathcal{F} \xrightarrow{\varphi} \mathcal{G} \xrightarrow{\psi} \mathcal{H} \rightarrow 0$$

Proposition 20.1.4. Let $\varphi: \mathcal{F} \rightarrow \mathcal{G}$ be a morphism between sheaves of abelian groups. Then for any $p \in X$, one has

$$\begin{aligned} (\ker \varphi)_p &= \ker \varphi_p \\ (\text{im } \varphi)_p &= \text{im } \varphi_p. \end{aligned}$$

Proof. For (1). It's clear $(\ker \varphi)_p \subseteq \ker \varphi_p$. Conversely, if $s_p \in \ker \varphi_p$, then $\varphi_p(s_p) = 0 \in \mathcal{G}_p$. In other words, there exists an open subset U containing p and $s \in \mathcal{F}(U)$ such that $s|_p = s_p$ and $\varphi(s)|_p = 0$, which implies there is another open subset V containing p such that $\varphi(s)|_V = 0$. Hence $\varphi(s|_V) = 0$, that is, $s|_V \in \ker \varphi(V)$. Thus $s_p = (s|_V)|_p \in (\ker \varphi)_p$.

For (2). It's clear $(\text{im } \varphi)_p \subseteq \text{im } \varphi_p$ since the sheafification doesn't change stalk. Conversely, if $s_p \in \text{im } \varphi_p$, then there exists $t_p \in \mathcal{F}_p$ such that $\varphi_p(t_p) = s_p$. Suppose $t \in \mathcal{F}(U)$ is a section of some open subset U containing p such that $t|_p = t_p$. Then $\varphi(t)|_p = \varphi_p(t_p) = s_p$. In other words, s_p is in the stalk of the image presheaf at p , but the sheafification doesn't change stalk, so we have $s_p \in (\text{im } \varphi)_p$. \square

Corollary 20.1.1. The sequence of sheaves

$$\dots \rightarrow \mathcal{F}^{i-1} \xrightarrow{\varphi^{i-1}} \mathcal{F}^i \xrightarrow{\varphi^i} \mathcal{F}^{i+1} \rightarrow \dots$$

is exact if and only if the sequence of abelian groups are exact

$$\dots \rightarrow \mathcal{F}_p^{i-1} \xrightarrow{\varphi_p^{i-1}} \mathcal{F}_p^i \xrightarrow{\varphi_p^i} \mathcal{F}_p^{i+1} \rightarrow \dots$$

for all $p \in X$.

Corollary 20.1.2. The the sequence of sheaves

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{G}$$

is exact if and only if for any open subset U , the following sequence of abelian groups is exact

$$0 \rightarrow \mathcal{F}(U) \rightarrow \mathcal{G}(U).$$

Method one. For any open subset $U \subseteq X$, one has

$$\varphi(U): \mathcal{F}(U) \rightarrow \mathcal{G}(U)$$

is injective, since by definition we have for any open subset $U \subseteq X$, $\ker \varphi(U) = 0$, that is injectivity. \square

Method two. Or from another point of view, for each $p \in U$, we have

$$\varphi_p: \mathcal{F}_p \rightarrow \mathcal{G}_p$$

is injective. That is $\ker \varphi_p = 0$. So we obtain $(\ker \varphi(U))_p = 0$ for all $p \in U$. But for a section $s \in \mathcal{F}(U)$ if we have $s|_p = 0$, then we must have $s = 0$, and thus $\ker \varphi(U) = 0$. \square

Example 20.1.3 (exponential sequence). Let X be a complex manifold and \mathcal{O}_X be its holomorphic function sheaf. Then

$$0 \rightarrow 2\pi\sqrt{-1}\mathbb{Z} \rightarrow \mathcal{O}_X \xrightarrow{\text{exp}} \mathcal{O}_X^* \rightarrow 0$$

is an exact sequence of sheaves, called exponential sequence.

Proof. The difficulty is to show exponential map is surjective on stalks at $p \in X$. That is we need to construct logarithms of functions $g \in \mathcal{O}_X^*(U)$ for U , a neighborhood of p . We may choose U is simply-connected, then define

$$\log g(q) = \log g(p) + \int_{\gamma_q} \frac{dg}{g}$$

for $q \in U$, where γ_q is a path from p to q in U , and the definition of $\log g(q)$ is independent of the choice of γ_q since U is simply-connected. \square

Remark 20.1.7. In fact, U is simply-connected is crucial for constructing logarithm. If we consider $X = \mathbb{C}$ and $U = \mathbb{C} \setminus \{0\}$, then

$$\exp: \mathcal{O}_X(U) \rightarrow \mathcal{O}_X^*(U)$$

cannot be surjective.

20.2. Derived functor formulation of Sheaf Cohomology. The category \underline{Ab}_X : sheaves of abelian groups on X . In this section we will introduce sheaf cohomology by considering it as a derived functor.

Given an exact sequence of sheaf as follows

$$0 \rightarrow \mathcal{F}' \xrightarrow{\phi} \mathcal{F} \xrightarrow{\psi} \mathcal{F}''.$$

By taking its section over open subset U , we obtain a sequence of abelian groups

$$0 \rightarrow \mathcal{F}'(U) \xrightarrow{\phi(U)} \mathcal{F}(U) \xrightarrow{\psi(U)} \mathcal{F}''(U).$$

Above sequence is not only exact at $\mathcal{F}'(U)$, but also is exact at $\mathcal{F}(U)$. In other words, the functor given by taking section over open subset is a left exact functor.

- (1) Firstly let's show $\ker \psi(U) \supseteq \text{im } \phi(U)$. For $s \in \mathcal{F}'(U)$, if we want to show $\psi \circ \phi(s) = 0$, it suffices to show $(\psi \circ \phi(s))|_p = 0$ for all $p \in U$ since \mathcal{F}'' is a sheaf. For any $p \in U$, by considering stalk at p we obtain an exact sequence of abelian groups

$$0 \rightarrow \mathcal{F}'_p \xrightarrow{\phi_p} \mathcal{F}_p \xrightarrow{\psi_p} \mathcal{F}''_p.$$

Then we obtain $\psi_p \circ \phi_p(s|_p) = 0$, which implies $(\psi \circ \phi(s))|_p = 0$.

- (2) Conversely, Given $s \in \ker \psi(U)$, we have $s|_p \in \ker \psi_p$ for any $p \in U$. By exactness of stalks, there exists $t_p \in \mathcal{F}'_p$ such that $\phi_p(t_p) = s|_p$. Thus there exists an open subset V_i containing p and $t_i \in \mathcal{F}'(V_i)$ such that $\phi(t_i) = s|_{V_i}$. Now it suffices to show these t_i can be glued together to obtain $t \in \mathcal{F}(U)$, and since \mathcal{F} is a sheaf, it suffices to check these t_i agree on intersections $V_i \cap V_j$. Note that $\phi(t_i - t_j|_{V_i \cap V_j}) = s|_{V_i \cap V_j} - s|_{V_i \cap V_j} = 0$, then these t_i agree on intersections since ϕ is injective.

Remark 20.2.1. From above argument, we can see that

$$0 \rightarrow \mathcal{F}' \xrightarrow{\phi} \mathcal{F} \xrightarrow{\psi} \mathcal{F}''$$

is exact if and only if for any open subset $U \subseteq X$

$$0 \rightarrow \mathcal{F}'(U) \xrightarrow{\phi(U)} \mathcal{F}(U) \xrightarrow{\psi(U)} \mathcal{F}''(U)$$

is exact.

In homological algebra, we always consider the derived functor of a left or right-exact functor. In particular, the functor of taking global section is a left exact functor, and its right derived functor defines the cohomology of a sheaf. Before we come into the definition of derived functor, firstly let's define the injective resolution of a sheaf.

Definition 20.2.1 (injective). A sheaf \mathcal{I} is injective if $\text{Hom}(-, \mathcal{I})$ is an exact functor.

Definition 20.2.2 (injective resolution). Let \mathcal{F} be a sheaf. An injective resolution of \mathcal{F} is an exact sequence

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{I}^0 \rightarrow \mathcal{I}^1 \rightarrow \mathcal{I}^2 \rightarrow \dots$$

where \mathcal{I}^i are injective for all i .

Theorem 20.2.1. Every sheaf admits an injective resolution.

Theorem 20.2.2. Let $\mathcal{F} \rightarrow \mathcal{I}^\bullet$ and $\mathcal{G} \rightarrow \mathcal{G}^\bullet$ are two resolutions and $\phi: \mathcal{F} \rightarrow \mathcal{G}$ be a morphism of sheaves. Then there exists a morphism $\tilde{\phi}: \mathcal{I}^\bullet \rightarrow \mathcal{G}^\bullet$ which lifts ϕ , which is unique up to homotopy.

Definition 20.2.3 (sheaf cohomology). Let \mathcal{F} be a sheaf of abelian groups. Then

$$H^p(X, \mathcal{F}) := H^p(\mathcal{I}^\bullet(X)).$$

Remark 20.2.2. The Theorem 20.2.2 shows that the definition of sheaf cohomology is independent of the choice of injective resolution.

Example 20.2.1. By definition, the 0-th cohomology is exact the global section

$$H^0(X, \mathcal{F}) := \ker \{ \mathcal{I}^0(X) \rightarrow \mathcal{I}^1(X) \}.$$

Thus $H^0(X, \mathcal{F}) = \mathcal{F}(X)$, the global sections of sheaf.

Example 20.2.2. If \mathcal{F} is a injective sheaf, then $H^i(X, \mathcal{F}) = 0$ for all $i > 0$, since the sheaf cohomology of injective sheaf can be computed by using the following special injective resolution

$$0 \rightarrow \mathcal{F} \xrightarrow{\text{id}} \mathcal{F} \rightarrow 0 \rightarrow 0 \rightarrow \dots$$

Theorem 20.2.3 (zig-zag). If

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{G} \rightarrow \mathcal{H} \rightarrow 0$$

is a short sequence of sheaves, then there is an induced long exact sequence of abelian groups

$$0 \rightarrow H^0(X, \mathcal{F}) \rightarrow H^0(X, \mathcal{G}) \rightarrow H^0(X, \mathcal{H}) \rightarrow H^1(X, \mathcal{F}) \rightarrow H^1(X, \mathcal{G}) \rightarrow \dots$$

Definition 20.2.4 (direct image). Let $f: X \rightarrow Y$ be continuous map between topological spaces and \mathcal{F} be a sheaf of abelian groups on X . The direct image of \mathcal{F} , denoted by $f_*\mathcal{F}$, is a sheaf on Y defined by

$$f_*\mathcal{F}(U) := \mathcal{F}(f^{-1}(U)).$$

Proposition 20.2.1. $f_*: \underline{Ab}_X \rightarrow \underline{Ab}_Y$ is a left exact functor.

Proof. Given an exact sequence of sheaves on X

$$0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}''.$$

Then we need to show

$$0 \rightarrow f_*\mathcal{F}' \rightarrow f_*\mathcal{F} \rightarrow f_*\mathcal{F}''$$

is also an exact sequence of sheaves on Y . By Remark 20.2.1 it suffices to show that for any open subset $V \subseteq Y$, we have the following exact sequence

$$0 \rightarrow f_*\mathcal{F}'(V) \rightarrow f_*\mathcal{F}(V) \rightarrow f_*\mathcal{F}''(V),$$

and that's exactly

$$0 \rightarrow \mathcal{F}'(f^{-1}(V)) \rightarrow \mathcal{F}(f^{-1}(V)) \rightarrow \mathcal{F}''(f^{-1}(V)).$$

Since f is continuous, then $f^{-1}(V)$ is an open subset in X , and thus above sequence of abelian is exact since $0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}''$ is exact. \square

20.3. Acyclic resolution. In practice it may be difficult for us to choose an injective resolution, so we usual other resolutions to compute sheaf cohomology.

Definition 20.3.1 (acyclic sheaf). A sheaf \mathcal{F} is acyclic if $H^i(X, \mathcal{F}) = 0$ for all $i > 0$.

Example 20.3.1. Injective sheaf is acyclic.

Definition 20.3.2 (acyclic resolution). Let \mathcal{F} be a sheaf. An acyclic resolution of \mathcal{F} is an exact sequence

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{A}^0 \rightarrow \mathcal{A}^1 \rightarrow \mathcal{A}^2 \rightarrow \dots$$

where \mathcal{A}^i is acyclic for all i .

Proposition 20.3.1. The cohomology of sheaf \mathcal{F} can be computed using acyclic resolution.

In fact, it's a quite homological techniques, called dimension shifting, so we will state this technique in language of homological algebra. Let's see a baby version of it.

Example 20.3.2. Let \mathcal{F} be a left exact functor and $0 \rightarrow A \rightarrow M_1 \rightarrow B \rightarrow 0$ be an exact sequence with M_1 is \mathcal{F} -acyclic. Then $R^{i+1}\mathcal{F}(A) \cong R^i\mathcal{F}(B)$ for $i > 0$, and $R^1\mathcal{F}(A)$ is the cokernel of $\mathcal{F}(M_1) \rightarrow \mathcal{F}(B)$.

Proof. By considering the long exact sequence induced by $0 \rightarrow A \rightarrow M^1 \rightarrow B \rightarrow 0$, one has

$$R^i\mathcal{F}(M^1) \rightarrow R^i\mathcal{F}(B) \rightarrow R^{i+1}\mathcal{F}(A) \rightarrow R^{i+1}\mathcal{F}(M^1)$$

(1) If $i > 0$, then $R^i\mathcal{F}(M^1) = R^{i+1}\mathcal{F}(M^1) = 0$ since M^1 is \mathcal{F} -acyclic, and thus $R^{i+1}\mathcal{F}(A) \cong R^i\mathcal{F}(B)$ for $i > 0$.

(2) If $i = 0$, then

$$0 \rightarrow \mathcal{F}(M^1) \rightarrow \mathcal{F}(B) \rightarrow R^1\mathcal{F}(A) \rightarrow 0$$

implies $R^1\mathcal{F}(A) = \text{coker}\{\mathcal{F}(M^1) \rightarrow \mathcal{F}(B)\}$.

□

Now let's prove dimension shifting in a general setting.

Lemma 20.3.1 (dimension shifting). If

$$0 \rightarrow A \rightarrow M^1 \rightarrow M^2 \rightarrow \cdots \rightarrow M^m \rightarrow B \rightarrow 0$$

is exact with M^i is \mathcal{F} -acyclic, then $R^{i+m}\mathcal{F}(A) \cong R^i\mathcal{F}(B)$ for $i > 0$, and $R^m\mathcal{F}(A)$ is the cokernel of $\mathcal{F}(M^m) \rightarrow \mathcal{F}(B)$.

Proof. Prove it by induction on m . For $m = 1$, we already see it in Example 20.3.2. Assume it holds for $m < k$, then for $m = k$, let's split $0 \rightarrow A \rightarrow M^1 \rightarrow M^2 \rightarrow \cdots \rightarrow M^k \xrightarrow{d_k} B \rightarrow 0$ into two exact sequences

$$0 \rightarrow A \rightarrow M^1 \rightarrow M^2 \rightarrow \cdots \rightarrow M^{k-1} \rightarrow \ker d_k \rightarrow 0$$

$$0 \rightarrow \ker d_k \rightarrow M^k \xrightarrow{d_k} B \rightarrow 0.$$

Then by induction hypothesis, for $i > 0$ we have

$$R^{i+k-1}\mathcal{F}(A) \cong R^i\mathcal{F}(\ker d_k)$$

$$R^{i+1}\mathcal{F}(\ker d_k) \cong R^i\mathcal{F}(B).$$

Combine these two isomorphisms together we obtain $R^{i+k}\mathcal{F}(A) \cong R^i\mathcal{F}(B)$ for $i > 0$, as desired. For $i = 0$, it suffices to let $i = 1$ in $R^{i+k-1}\mathcal{F}(A) \cong R^i\mathcal{F}(\ker d_k)$, then we obtain

$$R^k\mathcal{F}(A) = R^1\mathcal{F}(\ker d_k) = \text{coker}\{\mathcal{F}(M^k) \rightarrow \mathcal{F}(B)\}.$$

This completes the proof. □

Corollary 20.3.1. If $0 \rightarrow A \rightarrow M^\bullet$ is a \mathcal{F} -acyclic resolution, then $R^i\mathcal{F}(A) = H^i(\mathcal{F}(M^\bullet))$.

Proof. Truncate the resolution as

$$0 \rightarrow A \rightarrow M^0 \rightarrow M^1 \rightarrow \cdots \rightarrow M^{i-1} \rightarrow B \rightarrow 0$$

$$0 \rightarrow B \rightarrow M^i \rightarrow M^{i+1} \rightarrow \cdots$$

Since we already have $R^i\mathcal{F}(A) = \text{coker}\{\mathcal{F}(M^{i-1}) \rightarrow \mathcal{F}(B)\}$, and \mathcal{F} is left exact, one has

$$\mathcal{F}(B) = \ker\{\mathcal{F}(M^i) \rightarrow \mathcal{F}(M^{i+1})\}.$$

Thus we obtain

$$R^i\mathcal{F}(A) = \text{coker}\{\mathcal{F}(M^{i-1}) \rightarrow \ker\{\mathcal{F}(M^i) \rightarrow \mathcal{F}(M^{i+1})\}\} = H^i(\mathcal{F}(M^\bullet)).$$

□

20.4. Examples about acyclic sheaf.

20.4.1. *Flabby sheaf.* First kind of acyclic sheaf is flabby¹⁹ sheaf.

Definition 20.4.1 (flabby). A sheaf \mathcal{F} is flabby if for all open $U \subseteq V$, the restriction map $\mathcal{F}(V) \rightarrow \mathcal{F}(U)$ is surjective.

Now let's see some examples about flabby sheaves.

Example 20.4.1. A constant sheaf on an irreducible topological space is flabby.

Proof. Note that the constant presheaf on a irreducible topological space is a sheaf in fact, and it's easy to see this constant presheaf is flabby. \square

In particular, we have

Example 20.4.2. Let X be an algebraic variety. Then constant sheaf \mathbb{Z}_X is flabby.

Example 20.4.3. If \mathcal{F} is a flabby sheaf on X , and $f: X \rightarrow Y$ is a continuous map, then $f_*\mathcal{F}$ is a flabby sheaf on Y .

Proof. For $V \subseteq W$ in Y , it suffices to show $f_*\mathcal{F}(W) \rightarrow f_*\mathcal{F}(V)$ is surjective, and that's

$$\mathcal{F}(f^{-1}W) \rightarrow \mathcal{F}(f^{-1}V)$$

it's surjective since \mathcal{F} is flabby. \square

Example 20.4.4. An injective sheaf is flabby.

Proof. Let \mathcal{I} be an injective sheaf and $V \subseteq U$ be open subsets. Now we define sheaf \mathbb{Z}_U on X by

$$\mathbb{Z}_U := \begin{cases} \mathbb{Z}(W) & W \subseteq U \\ 0 & \text{otherwise} \end{cases}$$

where \mathbb{Z} is constant sheaf valued in \mathbb{Z} , and similarly we define sheaf \mathbb{Z}_V . By construction one has $\mathbb{Z}_U(W) = \mathbb{Z}_V(W)$ unless $W \subseteq U$ and $W \not\subseteq V$. Thus we obtain an exact sequence

$$0 \rightarrow \mathbb{Z}_V \rightarrow \mathbb{Z}_U.$$

Applying the functor $\text{Hom}(-, \mathcal{I})$, which is exact, we obtain an exact sequence

$$\text{Hom}(\mathbb{Z}_U, \mathcal{I}) \rightarrow \text{Hom}(\mathbb{Z}_V, \mathcal{I}) \rightarrow 0.$$

Now let's explain why we need such a weird sheaf \mathbb{Z}_U . In fact, we will prove $\text{Hom}(\mathbb{Z}_U, \mathcal{I}) = \mathcal{I}(U)$. Indeed since $\varphi: \mathbb{Z}_U \rightarrow \mathcal{I}$ is a sheaf morphism. Then if $W \not\subseteq U$, then $\varphi(U)$ must be zero. If $W = U$, then the group of sections of $\mathbb{Z}_U(U)$ over any connected component is simply \mathbb{Z} and hence $\varphi(U)$ on this connected component is determined by the image of $1 \in \mathbb{Z}$. Thus $\varphi(U)$ can be thought of an element of $\mathcal{I}(U)$. Now on any proper open subset of U , φ is determined by restriction maps. Hence $\text{Hom}(\mathbb{Z}_U, \mathcal{I}) = \mathcal{I}(U)$, as desired.

¹⁹Some authors also call this flasque.

The same argument shows $\text{Hom}(\underline{\mathbb{Z}}_U, \mathcal{I}) = \mathcal{I}(V)$, and thus we obtain an exact sequence

$$\mathcal{I}(U) \rightarrow \mathcal{I}(V) \rightarrow 0,$$

which shows \mathcal{I} is flabby. \square

Our goal is to prove a flabby sheaf is acyclic, but we still need some property of flabby sheaves.

Proposition 20.4.1. If $0 \rightarrow \mathcal{F}' \xrightarrow{\phi} \mathcal{F} \xrightarrow{\psi} \mathcal{F}'' \rightarrow 0$ is an exact sequence of sheaves, and \mathcal{F}' is flabby, then for any open subset U , the sequence

$$0 \rightarrow \mathcal{F}'(U) \xrightarrow{\phi(U)} \mathcal{F}(U) \xrightarrow{\psi(U)} \mathcal{F}''(U) \rightarrow 0$$

is exact.

Proof. It suffices to show $\mathcal{F}(U) \rightarrow \mathcal{F}''(U) \rightarrow 0$ is exact. Here we only gives a sketch of the proof. Since we have exact sequence on stalks for each $p \in U$ as follows

$$0 \rightarrow \mathcal{F}'_p \xrightarrow{\phi_p} \mathcal{F}_p \xrightarrow{\psi_p} \mathcal{F}''_p \rightarrow 0$$

Then for each $s \in \mathcal{F}''(U)$, there exists $t_p \in \mathcal{F}_p$ such that $\psi_p(t_p) = s|_p$, so there exists open subset $V_p \subseteq U$ containing p and $t \in \mathcal{F}(V_p)$ such that $\psi(t) = s|_{V_p}$. If we can glue these t together then we get a section in $\mathcal{F}(U)$ and is mapped to s , which completes the proof. However, they may not equal on the intersection. But things are not too bad, consider another point q and $t' \in \mathcal{F}(V_q)$ such that $\psi(t') = s|_{V_q}$, $(t' - t)|_{V_p \cap V_q} \in \ker \psi|_{V_p \cap V_q} = \text{im } \phi|_{V_p \cap V_q}$. So there exists $t'' \in \mathcal{F}'(V_p \cap V_q)$ such that

$$\phi(t'') = (t' - t)|_{V_p \cap V_q}$$

Now since \mathcal{F}' is flabby, then there exists $t''' \in \mathcal{F}'(V_p)$ such that $t'''|_{V_p \cap V_q} = t''$. And consider $t + \phi(t''') \in \mathcal{F}(V_p)$, which will coincide with t' on $V_p \cap V_q$. After above corrections, we can glue t after correction together. \square

Proposition 20.4.2. If $0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0$ is an exact sequence of sheaves, and if \mathcal{F}' and \mathcal{F} are flabby, then \mathcal{F}'' is flabby.

Proof. Take $V \subseteq U$ and consider the following diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{F}'(U) & \longrightarrow & \mathcal{F}(U) & \longrightarrow & \mathcal{F}''(U) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{F}'(V) & \longrightarrow & \mathcal{F}(V) & \longrightarrow & \mathcal{F}''(V) & \longrightarrow & 0 \end{array}$$

Then the desired result follows from five lemma. \square

Proposition 20.4.3. A flabby sheaf is acyclic.

Proof. Let \mathcal{F} be a flabby sheaf. Since there are enough injective objects in the category of sheaf of abelian groups, there is an exact sequence

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{I} \rightarrow \mathcal{Q} \rightarrow 0$$

with \mathcal{I} is injective. By Example 20.4.4 we have \mathcal{I} is flabby, and thus by Proposition 20.4.2 we have \mathcal{Q} is flabby. Consider the long exact sequence induced from above short exact sequence

$$\mathcal{F}(X) \rightarrow \mathcal{I}(X) \rightarrow \mathcal{Q}(X) \rightarrow H^1(X, \mathcal{F}) \rightarrow H^1(X, \mathcal{I}) \rightarrow \dots$$

Note that $H^1(X, \mathcal{I}) = 0$ since \mathcal{I} is injective, and thus acyclic. Then $H^1(X, \mathcal{F}) = \text{coker}\{\mathcal{I}(X) \rightarrow \mathcal{Q}(X)\}$. But Proposition 20.4.1 shows that $\mathcal{I}(X) \rightarrow \mathcal{Q}(X)$ is surjective since \mathcal{F} is flabby, so $H^1(X, \mathcal{F}) = 0$.

Now let's prove $H^k(X, \mathcal{F}) = 0$ for $k > 0$ by induction on k , and above argument shows it's true for $k = 1$. Assume this holds for $k < n$, and consider

$$\dots \rightarrow H^{n-1}(X, \mathcal{Q}) \rightarrow H^n(X, \mathcal{F}) \rightarrow H^n(X, \mathcal{I}) \rightarrow H^n(X, \mathcal{Q}) \rightarrow \dots$$

By induction hypothesis, we can reduce above sequence to

$$\dots \rightarrow 0 \rightarrow H^n(X, \mathcal{F}) \rightarrow 0 \rightarrow H^n(X, \mathcal{Q}) \rightarrow \dots$$

which implies $H^n(X, \mathcal{F}) = 0$. This completes the proof. \square

20.4.2. *Soft sheaf.* The second kind of acyclic sheaves is called soft sheaves, which is quite similar to flabby.

Definition 20.4.2 (soft). A sheaf \mathcal{F} over X is soft if for any closed subset $S \subseteq X$ the restriction map $\mathcal{F}(X) \rightarrow \mathcal{F}(S)$ is surjective.

Remark 20.4.1. For closed subset S , the section over it is defined by

$$\mathcal{F}(S) := \varinjlim_{S \subseteq U} \mathcal{F}(U)$$

Parallel to Proposition 20.4.1 and Proposition 20.4.2, soft sheaf has the following properties:

Proposition 20.4.4. If $0 \rightarrow \mathcal{F}' \xrightarrow{\phi} \mathcal{F} \xrightarrow{\psi} \mathcal{F}'' \rightarrow 0$ is an exact sequence of sheaves, and \mathcal{F}' is soft, then the following sequence

$$0 \rightarrow \mathcal{F}'(X) \xrightarrow{\phi(X)} \mathcal{F}(X) \xrightarrow{\psi(X)} \mathcal{F}''(X) \rightarrow 0$$

is exact.

Proposition 20.4.5. If $0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0$ is an exact sequence of sheaves, and if \mathcal{F}' and \mathcal{F} are soft, then \mathcal{F}'' is soft.

Proposition 20.4.6. A soft sheaf is acyclic.

So you may wonder, what's the difference between flabby and soft since the definitions are quite similar, and both of them are acyclic. Clearly by definition of sections over a closed subset, we know that every flabby sheaf is soft, but converse fails

Example 20.4.5. The sheaf of smooth functions on a smooth manifold is soft but not flabby.

Lemma 20.4.1. If \mathcal{M} is a sheaf of modules over a soft sheaf of rings \mathcal{R} , then \mathcal{M} is a soft sheaf.

Proof. Let $s \in \mathcal{M}(K)$ for some closed subset $K \subseteq X$. Then s extends to some open neighborhood U of K . Let $\rho \in \mathcal{R}(K \cup (X \setminus U))$ be defined by

$$\rho = \begin{cases} 1, & \text{on } K \\ 0, & \text{on } X \setminus U \end{cases}$$

Since \mathcal{R} is soft, then ρ extends to a section over X , then $\rho \circ s$ is the desired extension of s . \square

20.4.3. *Fine sheaf.* Another important kind of acyclic sheaves, which behaves like sheaf of differential forms Ω_X^k is called fine sheaf. Recall what is a partition of unity: Let $U = \{U_i\}_{i \in I}$ be a locally finite open covering of topological space X . A partition of unity subordinate to U is a collection of continuous functions $f_i: U_i \rightarrow [0, 1]$ for each $i \in I$ such that its support lies in U_i , and for any $x \in X$

$$\sum_{i \in I} f_i(x) = 1.$$

Definition 20.4.3 (fine sheaf). A fine sheaf \mathcal{F} on X is a sheaf of \mathcal{A} -modules, where \mathcal{A} is a sheaf of rings such that for every locally finite open covering $\{U_i\}_{i \in I}$ of X , there is a partition of unity

$$\sum_{i \in I} \rho_i = 1$$

where $\rho_i \in \mathcal{A}(X)$ and $\text{supp}(\rho_i) \subseteq U_i$.

Remark 20.4.2. For a sheaf \mathcal{F} on X and a section $s \in \mathcal{F}(X)$, its support is defined as

$$\text{supp}(s) := \overline{\{x \in X : s|_x \neq 0\}}.$$

Proposition 20.4.7. A fine sheaf is acyclic.

Proof. Let \mathcal{F} be a sheaf of \mathcal{A} -modules and a fine sheaf. And choose a injective resolution

$$0 \rightarrow \mathcal{F} \xrightarrow{d} \mathcal{I}^0 \xrightarrow{d} \mathcal{I}^1 \xrightarrow{d} \mathcal{I}^2 \xrightarrow{d} \dots$$

such that \mathcal{I}^i are injective sheaves of \mathcal{A} -modules. Let $s \in \mathcal{I}^0(X)$ such that $ds = 0$. Then by exactness of injective resolution we have X is covered by open subsets U_i such that for each i there is an element $t_i \in \mathcal{I}^{p-1}(U_i)$ such that $dt_i = s|_{U_i}$. By passing to a refinement we may assume that the cover $\{U_i\}$ is locally finite. Let $\{\rho_i\}$ be a partition of unity subordinate to $\{U_i\}$. Then we have $t = \sum \rho_i t_i \in \mathcal{I}^{p-1}(X)$ such that $dt = s$. This completes the proof. \square

Example 20.4.6. Let M be a smooth manifold and $\pi: E \rightarrow M$ be a vector bundle. Then the sheaf of smooth sections of E is a $C^\infty(M)$ -module sheaf, which is a fine sheaf. In particular, the sheaf of tangent bundle, sheaf of differential forms Ω_M and k -forms Ω_M^k are fine sheaves.

Remark 20.4.3. As a consequence, it's meaningless to compute cohomology of sheaf of differential k -forms, or any other vector bundle over a smooth manifold. But in complex version, something interesting happens: Let (X, \mathcal{O}_X) be a complex manifold and $\pi: E \rightarrow X$ be a holomorphic vector bundle. Then the sheaf of holomorphic sections of E is not a fine sheaf since there is no partition of unity may not be holomorphic, so the cohomology of holomorphic vector bundle is not trivial, and that's what Dolbeault cohomology computes.

For fine sheaf and soft sheaf, we have

Lemma 20.4.2. Fine sheaf is soft.

Proof. Let \mathcal{F} be a fine sheaf, $S \subseteq X$ closed and $s \in \mathcal{F}(S)$. Let $\{U_i\}$ be an open covering of S and $s_i \in \mathcal{F}(U_i)$ such that


$$s_i|_{S \cap U_i} = s|_{S \cap U_i}.$$

Let $U_0 = X - S$, and $s_0 = 0$. Then $\{U_i\} \amalg \{U_0\}$ is an open covering of X . Without loss of generality, we assume this open covering is locally finite and choose a partition of unity $\{\rho_i\}$ subordinate to it. Then

$$\bar{s} := \sum_i \rho_i(s_i)$$

is a section in $\mathcal{F}(X)$ which extends s . □

Remark 20.4.4. Until now, we have shown that soft, fine and flabby sheaves are acyclic. Lemma 20.4.2 shows fine sheaf is soft, and by definition a flabby sheaf is soft. The Example 20.4.5 shows that soft sheaf may not be flabby, and constant sheaf on an irreducible space is flabby but not fine. In a summary, we have the following relations:



20.5. Proof of de Rham theorem using sheaf cohomology. As we already know, for constant sheaf $\underline{\mathbb{R}}$ over a smooth manifold M , we have the following fine resolution

$$0 \rightarrow \underline{\mathbb{R}} \xrightarrow{i} \Omega_M^0 \xrightarrow{d} \Omega_M^1 \xrightarrow{d} \Omega_M^2 \xrightarrow{d} \dots$$

And de Rham cohomology computes the sheaf cohomology of $\underline{\mathbb{R}}$. de Rham theorem implies that de Rham cohomology equals to the singular cohomology with real coefficient. So if we can give constant sheaf another resolution using singular cochains, we may derive the de Rham cohomology.

We state this in a general setting: Let X be a topological manifold, and a constant sheaf \underline{G} over X , where G is an abelian group. Let $S^p(U, G)$ be the group of singular cochains in U with coefficients in G , and let δ denote the coboundary operator.

Let $\mathcal{S}^p(G)$ be the sheaf over X generated by the presheaf $U \mapsto S^p(U, G)$, with induced differential mapping $\mathcal{S}^p(G) \xrightarrow{\delta} \mathcal{S}^{p+1}(G)$.

Similar to Poincaré lemma, we have for a unit ball U in Euclidean space, we have the following sequence

$$\dots \rightarrow S^{p-1}(U, G) \xrightarrow{\delta} S^p(U, G) \xrightarrow{\delta} S^{p+1}(U, G) \rightarrow \dots$$

is exact. So we have the following resolution of the constant sheaf \underline{G}

$$0 \rightarrow \underline{G} \rightarrow \mathcal{S}^0(G) \xrightarrow{\delta} \mathcal{S}^1(G) \xrightarrow{\delta} \mathcal{S}^2(G) \rightarrow \dots$$

Remark 20.5.1. If M is a smooth manifold, then we can consider smooth chains, that is $f: \Delta^p \rightarrow U$, where f is a smooth function. The corresponding results above still hold, and we have a resolution by smooth cochains with coefficients in G :

$$0 \rightarrow \underline{G} \rightarrow \mathcal{S}_\infty^\bullet(G)$$

So if we choose $G = \mathbb{R}$, then it suffices to show $0 \rightarrow \underline{\mathbb{R}} \rightarrow \mathcal{S}_\infty^\bullet(\mathbb{R})$ is an acyclic resolution, then we obtain de Rham theorem.

First, note that \mathcal{S}_∞^p is a \mathcal{S}_∞^0 -module, given by cup product on open subsets. Then by Lemma 20.4.1 and the fact \mathcal{S}_∞^0 is soft we know that it's a soft resolution. This completes the proof.

20.6. Hypercohomology. In homological algebra, the hypercohomology is a generalization of cohomology functor which takes as input not objects in abelian category but instead chain complexes of objects.

One of the motivations for hypercohomology is to generalize the zig-zag lemma, that is, the short exact sequence of sheaves induces a long exact sequence of cohomology groups. It turns out hypercohomology gives techniques for constructing a similar cohomological associated long exact sequence from an arbitrary long exact sequence

$$0 \rightarrow \mathcal{F}_1 \rightarrow \mathcal{F}_2 \rightarrow \dots \rightarrow \mathcal{F}_k \rightarrow 0$$

Now let's give the definition of hypercohomology: Let $\mathcal{F}^\bullet: \dots \rightarrow \mathcal{F}^{i-1} \rightarrow \mathcal{F}^i \rightarrow \mathcal{F}^{i+1} \rightarrow \dots$ be a complex of sheaves of abelian groups, which is

bounded from below, that is, $\mathcal{F}^n = 0$ for $n \ll 0$. Then \mathcal{F}^\bullet admits an injective resolution $\mathcal{F}^\bullet \rightarrow \mathcal{I}^\bullet$. In other words,

$$\begin{array}{ccccccc} \dots & \longrightarrow & \mathcal{F}^{i-1} & \longrightarrow & \mathcal{F}^i & \longrightarrow & \mathcal{F}^{i+1} & \longrightarrow & \dots \\ & & \downarrow & & \downarrow & & \downarrow & & \\ \dots & \longrightarrow & \mathcal{I}^{i-1} & \longrightarrow & \mathcal{I}^i & \longrightarrow & \mathcal{I}^{i+1} & \longrightarrow & \dots \end{array}$$

such that

- (1) All \mathcal{I}^i are injective sheaves.
- (2) The induced homomorphism $H^i(\mathcal{F}^\bullet) \rightarrow H^i(\mathcal{I}^\bullet)$ is an isomorphism.

The hypercohomology of \mathcal{F}^\bullet is defined by

$$H^i(X, \mathcal{F}^\bullet) := H^i(\Gamma(X, \mathcal{I}^\bullet))$$

Definition 20.6.1. For a sheaf \mathcal{F} , $\mathcal{F}^\bullet[n]$ is a sheaf of complex defined by

$$(\mathcal{F}^\bullet[n])^i = \begin{cases} \mathcal{F} & i = n \\ 0 & \text{otherwise.} \end{cases}$$

Example 20.6.1. Let \mathcal{F} be a sheaf and $0 \rightarrow \mathcal{F} \rightarrow \mathcal{I}^0 \rightarrow \mathcal{I}^1 \rightarrow \dots$ be an injective resolution of \mathcal{F} . Then

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{F} & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \dots \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathcal{I}^0 & \longrightarrow & \mathcal{I}^1 & \longrightarrow & \mathcal{I}^2 & \longrightarrow & \dots \end{array}$$

is an injective resolution of $\mathcal{F}^\bullet[0]$. Indeed, \mathcal{I}^i are injective for all $i \geq 0$, and

$$H^i(\mathcal{I}^\bullet) = \begin{cases} \mathcal{F}, & n = 0 \\ 0, & \text{otherwise} \end{cases} = H^i(\mathcal{F}^\bullet[0])$$

So by definition of hypercohomology, we have $H^i(X, \mathcal{F}^\bullet[0]) = H^i(\Gamma(X, \mathcal{I}^\bullet)) = H^i(X, \mathcal{F}^\bullet)$. In general, one has

$$H^i(X, \mathcal{F}^\bullet[n]) \cong H^{i+n}(X, \mathcal{F}).$$

Theorem 20.6.1 (zig-zag). Let $0 \rightarrow \mathcal{F}^\bullet \rightarrow \mathcal{G}^\bullet \rightarrow \mathcal{H}^\bullet \rightarrow 0$ be a short exact sequence of complexes of sheaves which are bounded from below. Then there is an induced long exact sequence

$$\dots \rightarrow H^{i-1}(X, \mathcal{H}^\bullet) \rightarrow H^i(X, \mathcal{F}^\bullet) \rightarrow H^i(X, \mathcal{G}^\bullet) \rightarrow H^i(X, \mathcal{H}^\bullet) \rightarrow H^{i+1}(X, \mathcal{F}^\bullet) \rightarrow \dots$$

21. HODGE THEOREM

REFERENCES

- [Aub78] Thierry Aubin. Équations du type Monge-Ampère sur les variétés kählériennes compactes. *Bull. Sci. Math. (2)*, 102(1):63–95, 1978.
- [Cal57] Eugenio Calabi. On Kähler manifolds with vanishing canonical class. In *Algebraic geometry and topology. A symposium in honor of S. Lefschetz*, pages pp 78–89. Princeton University Press, Princeton, N.J., 1957.
- [Dem12] Jean-Pierre Demailly. *Complex Analytic and Differential Geometry*. Université de Grenoble I, 2012.
- [FLW17] Huitao Feng, Kefeng Liu, and Xueyuan Wan. Compact Kähler manifolds with positive orthogonal bisectional curvature. *Math. Res. Lett.*, 24(3):767–780, 2017.
- [Gau77] Paul Gauduchon. Le théorème de l'excentricité nulle. *C. R. Acad. Sci. Paris Sér. A-B*, 285(5):A387–A390, 1977.
- [Huy05] Daniel Huybrechts. *Complex Geometry An Introduction*. Springer Berlin, Heidelberg, 2005.
- [KN69] Shoshichi Kobayashi and Katsumi Nomizu. *Foundations of differential geometry. Vol. II*, volume Vol. II of *Interscience Tracts in Pure and Applied Mathematics, No. 15*. Interscience Publishers John Wiley & Sons, Inc., New York-London-Sydney, 1969.
- [Liu23] Bowen Liu. Riemannian geometry. https://bowenl-math.github.io/notes/2022Fall/Riemannian_geometry.pdf, 2023.
- [MR054] *Proceedings of the International Congress of Mathematicians, Amsterdam, 1954. Vol. 2*. P. Noordhoff N. V., Groningen; North-Holland Publishing Co., Amsterdam, 1954.
- [Sim88] Carlos T. Simpson. Constructing variations of Hodge structure using Yang-Mills theory and applications to uniformization. *J. Amer. Math. Soc.*, 1(4):867–918, 1988.
- [Tia00] Gang Tian. *Canonical metrics in Kähler geometry*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 2000. Notes taken by Meike Akveld.
- [UY86] K. Uhlenbeck and S.-T. Yau. On the existence of Hermitian-Yang-Mills connections in stable vector bundles. volume 39, pages S257–S293. 1986. *Frontiers of the mathematical sciences: 1985 (New York, 1985)*.
- [Yau75] Shing Tung Yau. Harmonic functions on complete Riemannian manifolds. *Comm. Pure Appl. Math.*, 28:201–228, 1975.
- [Yau77] Shing Tung Yau. Calabi's conjecture and some new results in algebraic geometry. *Proc. Nat. Acad. Sci. U.S.A.*, 74(5):1798–1799, 1977.
- [Yau78a] Shing Tung Yau. A general Schwarz lemma for Kähler manifolds. *Amer. J. Math.*, 100(1):197–203, 1978.
- [Yau78b] Shing Tung Yau. On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation. I. *Comm. Pure Appl. Math.*, 31(3):339–411, 1978.